

**NITROGEN TRANSFORMATION PROCESSES
IN AGRICULTURAL WATERSHED SOILS**

**FINAL REPORT on PROJECT 11
AGRICULTURAL WATERSHED STUDIES**

Task C (Canadian Section) - Activity 1
International Reference Group on Great Lakes
Pollution from Land Use Activities
International Joint Commission

G.C. Kowalenko
Soil Research Institute
Agriculture Canada
Ottawa, Ontario

April, 1978

ACKNOWLEDGEMENTS

The study discussed in this document was carried out as part of the efforts of the International Reference Group on Great Lakes Pollution From Land Use Activities (PLUARG), an organization of the International Joint Commission, established under the Canada - U.S. Great Lakes Water Quality Agreement of 1972. Funding was provided through Agriculture Canada. Findings and conclusions are those of the author and do not necessarily reflect the views of the Reference Group or its recommendations to the Commission.

The technical assistance of D. McDill, C. O'Meara, I. Nicholson and K. Wires, the computer programming of Valerie Malmsten and assistance of other S.R.I. staff is gratefully acknowledged.

TABLE OF CONTENTS

	PAGE
List of Tables	iii
Summary	1
Introduction	2
Data Collection Methods	2
Experimental Results	5
Data Analysis and Interpretation	10
Relationship of Project Results to PLUARG Objectives	11
Supplementary Information	11
Appendix: Rates of gaseous loss, mineralization and plant uptake of nitrogen in soil systems: A Literature Review	53

LIST OF TABLES

TABLE	PAGE
1. Description of surface soil samples from Watershed 5 (IJC-PLUARG Task C - Canada).	13
2. Some characterization analyses of soil samples from Watersheds 1, 5 and 13 (IJC-PLUARG Task C - Canada).	14
3. Inorganic N analyses ($\mu\text{g N/g}$ soil) during incubation experiments examining nitrification and mineralization of watershed soils under combinations of temperature and moisture contents.	15
4. Recovery of added $\text{NH}_4^+\text{-N}$ ($80 \mu\text{g N/g}$) from soil profiles representing watersheds 1, 5 and 13.	34
5. Rate coefficients of nitrification of added $\text{NH}_4^+\text{-N}$ in soils from three (3) watersheds.	35
6. Correlations of nitrification rate coefficients with temperature and moisture in surface samples of watershed soils.	37
7. Weighted average daily rates of mineralization of watershed soils.	38
8. Correlations of rate of N mineralization values with environmental and soil factors of incubation watershed surface soil samples.	40
9. Total surface nitrogen and yearly N mineralized assuming 1% of the total N is mineralized/year.	41
10. Total inorganic N in soil-plant residue incubations.	42
11. Inorganic N analyses ($\mu\text{g N/g}$) during incubation experiments examining denitrification under saturated moisture contents and several temperatures of watershed soils.	43
12. Denitrification rate coefficients of representative soil samples at saturated or near-saturated moisture content.	51
13. Potential N fixation (expressed as μg ethylene produced/g soil/day) in representative soil profile samples.	52

SUMMARY

Soils were characterized with respect to the rates of biological processes important to N leaching considerations. Nitrification was shown to be a relatively fast process which was related to temperature and moisture content conditions. The cross product of temperature x water content was particularly important. A regression equation including these environmental factors was derived for several groups of soils, depending largely on textural differences.

Mineralization was particularly influenced by environmental factors (temperature and moisture content) when soil (%C, %N, pH, etc.) and environmental factors were considered. An estimate of the amount of N produced by mineralization under field conditions suggested considerable N released, depending on the soil type. More precise calculations would need detailed knowledge of field conditions of moisture and temperature and applying the relationships derived in this study to these conditions. Incorporation of plant residues alters the mineralization-immobilization pattern in soils considerably and is dependent on the content of the plant material. Material with a low percentage N results in net immobilization of N. Organic amendments must be considered in the long term for N considerations because of changing mineralization-immobilization relationships.

Denitrification can progress very quickly if warm anaerobic conditions are met. The rates suggest that short but quantitatively large bursts of denitrification may be important in NO_3^- losses and these may occur during or immediately after rains. Calculation of field significance will depend on detailed knowledge of field moisture and temperature conditions.

Asymbiotic N fixation does not appear to be an important source of N in these soils.

Mineralization and denitrification appear to be two key processes to consider in addition to soil N leaching.

INTRODUCTION

The objective of this study was to characterize N transformation rates in a variety of soil types and horizons using laboratory incubation conditions and to evaluate how these conditions influence the rates. Biological transformations influence the sources and sinks of NO_3^- -N to the ground water. These rates are viewed as providing information for a mathematical model of N transport and transformation in the unsaturated soil zone in order to contribute to an understanding of the N storage and release characteristics of selected agricultural soils in the watersheds.

DATA COLLECTION METHODS

Soils

Soils were from three (3) watersheds (#1, #5 and #13) and comprised thirty-one (31) samples. The soils from watersheds 1 and 13 correspond to plot areas of Project 13 (Cameron et al., 1977. Nitrogen Movement in Tile-Drained Clay and Sandy Agricultural Watersheds. Final Report on Project 13, Agricultural Watershed Studies, Task C (Canadian Section) Activity 1, International Reference Group on Great Lakes Pollution from Land Use Activities. I.J.C.) and included surface and subsurface samples, as well as samples from a similar area taken at two different times (spring vs. fall). The soils from watershed #1 are Brookston clay and from watershed #13 are Berrien sand and were sampled in the spring and fall of 1975. Only surface soils were sampled from watershed #5, and they represent five (5) distinct soils (Table 1). All were sampled on September 9, 1975. Some characterizing analyses are given for all the samples (Table 2).

Sample Code Designation

Samples are identified by numbers and/or letters and can be categorized into two main groups. Those with a 2 number code (eg. 1-1, 2-1, 1-3, etc.) designate samples from plot studies (Cameron - Project 13). The first digit designates the plot number (i.e. there are six (6) plots numbered 1, 2, 6). The second digit designates the horizon depths which are as follows:

1	0-10 cm
2	10-20 cm
3	20-30 cm
4	30-45 cm
5	45-60 cm
6	60-75 cm
7	75-90 cm

In four cases, the 2 number code is followed by a letter (eg. 2-1-b) and this letter (b) designates a second sampling date, to evaluate how sampling time affects the results.

Samples designated by N-5 are samples from watershed 5. All samples are surface samples to a depth of 25 cm (except N-5/1 which is to 20 cm). The third digit denotes the soil within the watershed.

Incubation Systems

1) "Aerobic" incubations for:

a) Nitrification and Mineralization

Soil samples were incubated using combinations of addition and no addition of $(\text{NH}_4^+)2\text{SO}_4$ for twenty-one (21) days at 30% and 75% to field capacity water content at 4⁰, 15° and 30°C temperatures. The level of $(\text{NH}_4^+)2\text{SO}_4$ addition was 80µg NH_4^+ -N/g soil. Field capacity moisture content was taken to be 100 cm water tension, performed on a water tension table apparatus. Extraction of inorganic N was done using a sequential extraction using first 0.005 M CaCl_2 then 2N KCl (1:4 soil weight to extractant volume) for hr., centrifuged, and solution filtered and analysed for NH_4^+ -N and NO_3^- -N using autoanalyser. By operational definition, the CaCl_2 extracts all the NO_3^- -N plus all the soluble NH_4^+ -N. The KCl extracts the exchangeable NH_4^+ -N. Corrections for carry-over of soluble NH_4^+ -N and NO_3^- -N were made using a weighing technique. The samples were extracted immediately after addition of solution (to bring the soil to required water content and N addition) and after 5, 14 and 21 days of incubation. Sufficient 5 g samples were prepared to allow incubation and extraction, in duplicate, at each time and with each combination of temperature, water content, and $(\text{NH}_4^+)2\text{SO}_4$ addition. The incubation vessels were closed but aerated every two (2) to three (3) days throughout the incubation period to restrict water loss and maintain aerated conditions. After preliminary trials, some of the incubations were simplified by eliminating the analysis on the fifth day of the incubation, since it was too close to the initial burst of activity after wetting the air dry samples. Also only a KCl extraction was used, eliminating the CaCl_2 extraction, resulting in bulking soluble and exchangeable NH_4^+ -N into a single extractable fraction.

b) Mineralization of Plant Material

Plant material (finely ground) was added to Berrien and Brookston soil samples (equal mixtures of 2-1 and 2-2, and 5-1 and 5-2, respectively) taken to field capacity moisture content and incubated at 30C. Incubation was in beakers, covered with parafilm, having small aeration holes, and moisture maintained by weighing and water additions. The soil was sampled at days 0, 15, 30 and 60 for inorganic nitrogen analyses.

Plant materials added to the soils were from the field plots and were added at a rate corresponding to harvest of dry matter (i.e. 320, 1430, 1790 and 5370 mg/100 g. soil, using rye, soybean, wheat and corn respectively). Nitrogen content of these materials were 2.32, 2.58, 1.18 and 0.99% for rye, soybean, wheat and corn respectively.

2) "Anaerobic" incubations for:

a) Denitrification

Soil samples were incubated using combination of, addition and no addition of NH_4^+NO_3 for five (5) days at 75%, 100% and 120% of saturation water contents at 4°, 15° and 30° temperatures (C). The level of NH_4^+NO_3 addition was 80 μg NO_3^- -N/g soil and 80 μg NH_4^+ -N/g soil. No other amendments were made so the denitrification process was dependent on nutrients and energy sources inherent in the soil sample. Saturation water content was determined by adding water in small measured increments to a know weight of soil to a point just before free water was evident. Extraction of inorganic N was with 2N KCl (1:4 soil weight to extractant volume) for 2 hour, filtered and analysed for NH_4^+ -N and NO_3^- -N using autoanalyser. The samples were extracted immediately after addition of solution (to bring the soil to required water content and level of N added) and after two (2) and five (5) days incubation. Sufficient 5 g samples were prepared to allow incubation and extraction in duplicate at each time and with each combination of temperature, water content and NH_4^+NO_3 addition. Incubation vessels were closed through the period to restrict water loss. Laboratory tests showed that closing or leaving the vessels open made no difference on the N analysis, therefore for convenience the closed system was used. No other control of the gas phase was made. After several trials, the incubations were streamlined by including only one moisture content (saturation).

b) Asymbiotic N fixation

The incubation system was similar to the denitrification experiments, except that no N treatments were included and only one combination of moisture content (saturation) and temperature (30C) was used. Inorganic nitrogen and N fixation was measured by the ability of the system to convert acetylene to ethylene over a three-day period.

Inorganic N analyses: NH_4^+ -N and NO_3^- -N were analysed by autoanalyser using a distillation step (Keay, J. and P.M.A. Menage, 1970, Analyst 95:279-382; Quinn J.R., J.G.A. Boisvert and I. Wood, 1974, Anal. Biochem. 58:609-614). Presence or absence of NO_2^- was first determined by spot test and if present, quantitative analysis was determined by modified Greiss-Ilosvay color reaction (Bremner, J.M., 1965, Inorganic forms of nitrogen *IN* Methods of Soil Analysis Part 2 Chemical and Microbiological Properties, Agronomy Series No. 9 Amer. Soc. Agron. Madison pp. 1219-1224).

EXPERIMENTAL RESULTS

The results of the incubation experiments designed to examine rates of mineralization and nitrification in watershed soils are shown in Table 3. Besides giving information on rates, the results provide some information on other processes such as adsorption and fixation of NH_4^+ -N. Plotting soluble and exchangeable NH_4^+ -N values on a log-log scale, a straight line relationship was perceived. Distinct relationships were evident for the clay and sandy loam soils from watersheds #1 and #13. The following relationships were derived for soil 2-1 (sandy loam) and soil 4-1 (clay) respectively:

$$(\text{NH}_4^+\text{-N})_{\text{exch.}} = 6.8 [(\text{NH}_4^+\text{-N})_{\text{sol.}}]^{0.58} \quad (1)$$

$$(\text{NH}_4^+\text{-N})_{\text{exch.}} = 1.6 [(\text{NH}_4^+\text{-N})_{\text{sol.}}]^{0.76} \quad (2)$$

The relationship shows that a much larger proportion of the NH_4^+ -N goes on to the exchange sites in the clay than in the sandy loam soils. This reflects the available cation exchange sites. This means a much greater potential of leaching NH_4^+ -N in sandy loam soils than in clay soils or more precisely in soils of low cation exchange than with high cation exchange.

The Brookston clay soils contain a large quantity of fixed NH_4^+ -N and the Berrien sandy loam soils contain smaller but measureable amounts (Table 2). Since NH_4^+ -N was added to the soils in the mineralization- nitrification incubations and extracted immediately, the ability of the samples to fixed NH_4^+ -N can be assessed. Table 4 shows that negligible amounts of NH_4^+ -N was fixed in the Berrien sandy loam, whereas the Brookston clay showed some fixation in the subsoils. Small to negligible amounts were fixed in the surface, possibly showing that the fixation sites are almost completely saturated by fertilization, etc. Soils from watershed #5 showed low to negligible ability to fix NH_4^+ -N.

Nitrification

Nitrification is the oxidation of NH_4^+ -N to NO_3^- -N, which in soils, is mediated by micro-organisms. Although it has long been recognized that this is a two step process ($\text{NH}_4^+ \Rightarrow \text{NO}_2^- \Rightarrow \text{NO}_3^-$) with each step being done by a particular genus of autotrophic bacteria, the oxidation is often viewed as a single process. The initial step ($\text{NH}_4^+ \Rightarrow \text{NO}_2^-$) is usually the

rate limiting step and hence NO_2^- rarely accumulates in normal soils due to nitrification. For purposes of this study, the oxidation of $\text{NH}_4^+\text{-N}$ will be discussed as a single process.

Nitrification was measured in these soils by adding a known quantity of $\text{NH}_4^+\text{-N}$ ($80\mu\text{g N/g}$ soil) and monitoring changes in extractable inorganic N. Simultaneous to nitrification, mineralization of organic N also occurs. The end product is $\text{NH}_4^+\text{-N}$, which in turn, is nitrified. $\text{NH}_4^+\text{-N}$ rarely accumulates in a mineralizing soil, (i.e.) all $\text{NH}_4^+\text{-N}$ resulting from mineralization is nitrified immediately. The nitrification rate calculated in this experiment utilizes information from the $\text{NH}_4^+\text{-N}$ treated plots. $\text{NH}_4^+\text{-N}$ was added only at the beginning of the incubations, hence as the experiment progressed, the $\text{NH}_4^+\text{-N}$ may decrease to a point where it becomes limiting. The rate of nitrification can be calculated many ways, but considering the probability of $\text{NH}_4^+\text{-N}$ becoming limiting, a rate coefficient involving loss of $\text{NH}_4^+\text{-N}$ and gain of $\text{NO}_3^-\text{-N}$ was used. The actual calculation assumed a first order relationship (Cameron, D.R. and C.G. Kowalenko, 1976, *Can. J. Soil Sci.*, 56: 71-78).

$$k_n = -\ln (N_t / N_o)/t$$

where k_n = rate coefficient of nitrification (day^{-1});
 N_o = original control $\text{NH}_4^+\text{-N}$ concentration and

$$N_t = (N_t^1 + N_t^{11}) / 2$$

N_t^1 = net loss of fertilizer $\text{NH}_4^+\text{-N}$ over time t and N_t^{11} = net gain of $\text{NO}_3^-\text{-N}$ due to the fertilizer $\text{NH}_4^+\text{-N}$ added. By including both the $\text{NO}_3^-\text{-N}$ gain and $\text{NH}_4^+\text{-N}$ loss results more of the analytical data was used, thus increasing the reliability of the calculation. These rate coefficients were then calculated for 0-14 and 14-21 day intervals and the average of these two gives the accepted rate coefficient.

The rate coefficients calculated in this way are presented in Table 5. Each soil has six (6) rate coefficients from combinations of three (3) temperatures (4, 15, 30 C) and two (2) moisture contents (30 and 75% of field capacity). The rate coefficient changes considerably within a single soil, depending on the moisture and temperature combination. In a few cases, the rate could not be calculated because there was more inorganic N than could be accounted for by fertilizer N (i.e. mineralization was higher in fertilized than control samples). The nature of the soil also influences the coefficient considerably. The surface layers (0-10 and 10-20) of Berrien sandy loam samples (numbers 1, 2 and 3) had an average coefficient of 0.010 day^{-1} , with a range of -0.002 to 0.033 day^{-1} and the corresponding layers of Brookston clay samples (numbers 4, 5 and 6) had an average coefficient of 0.073 day^{-1} , with a range of 0.027 to 0.141 day^{-1} for 15C and 75% of field capacity moisture content, as an example. Comparison of samples (at 15C and 75% of field capacity) taken at two (2) different times (2-1 and 2-1b, 2-2 and 2-2b, 5-1 and 5-2b) show considerable variation. Rate coefficients for subsurface horizons are lower than surface horizons.

Correlation coefficients of nitrification rate coefficients with temperature and moisture

show that the interaction between temperature (T) and moisture content (θ) is the highest in all the surface samples and that a multiple correlation of all factors was not very much higher than the simple correlations. Correlations among factors for the Berrien sandy loam are the lowest of the three (3) groupings of soils. This relationship among the environmental variables and the rate coefficient was similar to previous work (Cameron, D.R. and C.G. Kowalenko, 1976, Can. J. Soil Sci., 56, 71-78). The regression equation relating the nitrification rate coefficient to environmental variables of this study are as follows:

$$\text{Brookston clay: } k_n = 10^{-3} (9.913 + 0.3845T\theta - 3.443T - 1.0603 \theta) \quad (4)$$

$$\text{Berrien sandy Loam: } k_n = 10^{-3} (1.569 + 0.209T\theta - 0.5684T - 0.6769\theta) \quad (5)$$

$$\text{N-5 } k_n = 10^{-3} (-4.515 + 0.4993T\theta - 4.3837T + 0.4811\theta) \quad (6)$$

These relationships are, in general, similar to the previous study (Cameron and Kowalenko, 1976) of an Ottawa area soil which was derived in a slightly different way. That relationship was

$$k_n = 10^{-3} (9.537 + 0.374T\theta - 0.545T - 0.2903\theta) \quad (7)$$

Mineralization

Although in the strictest sense, mineralization is the conversion of organic N to $\text{NH}_4^+\text{-N}$, nitrification is so rapid that it appears as $\text{NO}_3^-\text{-N}$ instead. To measure mineralization in these samples, the accumulation of organic N in a moistened soil (no N added) is taken as the net mineralization. "Net" mineralization is used since both mineralization and immobilization occur simultaneously, and only the net result can be measured. As with nitrification, mineralization rates can be calculated from the results in a number of ways. It was finally decided to use a weighted average daily rate. A rate coefficient was not attempted since one must measure or assume a decline in organic N. Organic N in a soil is a complex component, with fractions having variable degrees of availability toward mineralization, making meaningful measurements virtually impossible with present knowledge. The weighted average rate was calculated as follows:

$$r = .5 (N_{14} + N_{21} - 2N_0) / 17.5$$

where r = weighted daily rate of mineralization ($\mu\text{gN/g soil/day}$), N_0 = initial inorganic concentration ($\mu\text{gN/g}$), N_{14} = inorganic N concentration at day 14 ($\mu\text{gN/g}$), N_{21} = inorganic N concentration at day 21 ($\mu\text{gN/g}$) and 17.5 is the average of 14 and 21 days. The average rate was weighted in such manner to emphasize results of the later part of the incubation period since early in the incubation, there may have been a flush of activity due to wetting an air dry soil.

Table 7 shows the calculated weighted average daily rate of mineralization. As with nitrification, the rate of mineralization is higher in the Brookston clay than in the Berrien sandy loam, and in both soils, it decreases with depth. Unlike nitrification, the amount of the source material (organic N) is difficult to quantify and is not considered in the rate calculation of mineralization (i.e. for nitrification $\text{NH}_4^+\text{-N}$ is incorporated into the rate coefficient calculation). For this reason, correlations were extended beyond just environmental factors, but also soil characterization values (Table 8). It is evident that correlations are poor to non-existent with characterizing analyses, but better with environmental factors, particularly with moisture content and moisture content x temperature.

A generalization that 1 to 3% of soil N is mineralized yearly, has often been made (see Appendix 1 p. xix). Using this approach, the rate of mineralization can be calculated for watershed soils (Table 9). Assuming that the soil actively mineralized during 150 days (late spring to early fall) then average daily rates would range from 0.06 to 0.25 $\mu\text{gN/g/day}$. Table 9 also shows the very large pool of N in the surface 15 cm of the soils and small changes in daily rate of mineralization can be a very significant inorganic N source on a yearly basis.

The results from the soil-plant residue incubation show the influence of plant residue N concentration on the net mineralization-immobilization (Table 10). In the Brookston clay, only the soybean residue treatment resulted in net mineralization of N relative to the control, but in the Berrien sandy loam soil, both rye and soybean. Corn and wheat residue treatments resulted in about the same or less extractable inorganic N at 60 days compared with initial amounts. The soybean and rye material had the highest N concentrations and the wheat and corn residue had the lowest. Addition of plant residue appears to encourage net immobilization of N when the residue N concentration is relatively low. The N in the residue would probably be mineralized after longer incubations as the C/N ratio is narrowed.

Denitrification

Table 11 shows the results of the denitrification incubation study. The decrease in $\text{NO}_3^- \text{-N}$ was taken to be denitrification; immobilization was considered to be minimal particularly in the N amended samples since NH_4^+ was added to take care of this process. Increases in control $\text{NH}_4^+\text{-N}$ indicates that immobilization is not important even where additional $\text{NH}_4^+\text{-N}$ was not added. There was a fairly definite temperature effect, with cool temperatures retarding the $\text{NO}_3^- \text{-N}$ decrease considerably. $\text{NO}_2^- \text{-N}$ was measurable on occasion and was assumed to be a transient intermediate of denitrification. The moisture contents that were used initially were shown to have a very small effect and in subsequent incubations of soils, only one moisture content (saturation) was used. The surface Brookston clay samples had higher denitrification activity as compared with the sandier samples.

First order rate coefficients for denitrification (see Stanford et al 1975, S.S.S.A.P., 39, 284-289) were calculated as follows:

$$\text{RD}_t = -\ln (N_t / N_o) / t \quad (9)$$

where RD_t = denitrification rate coefficient at a given time, t = time, N_t and $N_0 = NO_3^- + NO_2^-$ concentrations at times t and zero, respectively. The rate coefficients are shown in Table 12. In several instances, a value could not be recorded since no NO_3^- nor NO_2^- -N was present resulting in equation (9) becoming ∞ which is unrealistic. These occur only in the control incubations where original NO_3^- -N available for denitrification was low.

Correlations of denitrification rate coefficients of fertilized surface samples with the temperature of incubation were 0.51, 0.77 and 0.91 for Berrien sandy loam, Brookston clay and N-5 soils, respectively. Similar correlations with the water content were very low (i.e. 0.07 to 0.30). This suggests that temperature has a definite influence on the rate coefficient, but the water content does not. The low correlation with water content was expected, since the water content would be adequate for microbial activity and functions more as an agent to encourage anaerobic conditions. Correlations of rate coefficient with %C, %N and glucose-C were low (less than 0.24). Glucose-C has been related to denitrification rates in other studies (Burford, J.R. and J.M. Bremner, 1975. *Soil Biol. Biochem.* 7, 389-394; Stanford, G., R.A. Vander Pol and S. Dzienia, 1975. *Soil Sci. Soc. Am. Proc.* 39, 284-289) and this difference may be due to the range of samples used in the study and/or including a wide range of temperature conditions in the rate coefficient calculation making environmental variables the overriding factors.

Nitrogen Fixation

Asymbiotic N fixation is one possible mechanism for N gain to the soil and it was not known how important the process is. The process requires anaerobic conditions, therefore, the water saturated conditions of denitrification incubations were employed to explore a maximum potential rate. It is known that excess inorganic N often retards the process, hence the incubation study for N fixation was extended such that denitrification could "strip away" any NO_3^- -N and subsequently encourage N fixation. Table 13 shows that representative soils and depths of these soils have very low asymbiotic N fixation potential. N fixation rate is shown as g ethylene produced /g soil/day and to convert this to N_2 fixation, the theoretical C_2H_2/N_2 conversion is around 3 (Hardy, R.W.F., R.C. Burns and R.D. Holsten, 1973. *Soil Biol. Biochem.* 5, 47-81), therefore, N_2 fixation appears to be rather minor input of N. Denitrification effectively reduced the NO_3^- -N content, but this reduction did not appear to increase the rate of N fixation. Time of original field sampling (eg. 2-1 vs. 2-1-b and 5-1 vs. 5-1-b) did not significantly affect the results.

DATA ANALYSIS AND INTERPRETATION

Although in the Berrien sandy loam samples, there is potential for NH_4^+ -N leaching because of a relatively high proportion of soluble NH_4^+ -N, the relatively low rate of nitrification would limit movement to the NO_3^- -N form. Fixation of added NH_4^+ -N occurred only in subsurface samples of the Brookston clay, this process would not be particularly important in N leaching considerations. Nitrification was relatively high in most surface samples studied, which would result in NH_4^+ -N being present in negligible quantities in these soils except for short periods after large NH_4^+ -N additions.

The very large pool of organic N contained in the surface horizon of the soils (see Table 9) is a potential source for N leaching to the ground water. Mineralization is the process that releases the N, which is quickly nitrified to NO_3^- -N (which is very mobile). Environmental factors are particularly important in influencing the rate of this process. The average daily rates of net mineralization measured at 15°C and 75% of field capacity moisture content were 0.1043, 0.7645 and 1.3309 $\mu\text{gN/g}$ for Berrien sandy loam, Brookston clay and N-5 surface samples, respectively. If these environmental conditions on average prevailed for 150 days in the field, approximately 36, 258 and 448 kg N/ha would be released over the summer season (i.e. entire year). This calculation of N release over a year is much higher than using the 1% of the total N in the soil. Incorporation of plant residues will alter the rate of mineral N release depending to a large degree on the nature of the residue. Material relatively high %N (e.g. legume residues or green manure materials) would encourage higher amounts of mineralization. The rate of this process must be put into context of other processes (eg. plant uptake and denitrification) and probably more detail of rates of these processes considered over seasonal changing environmental conditions are needed to determine the fate of the N in the soil. A literature review of rates of process is presented (Appendix) to enable comparison of rates with other studies.

Denitrification is much harder to assess over a period of a year because of the rather special conditions required for this process to proceed. Anaerobic conditions (whether in general or in microsites), warmth and NO_3^- -N are needed. An intimate consideration of field environmental conditions are required to apply the rate coefficients of this process to a weather conditions of field season. The quick initiation of the process upon saturation and fast rate of the process suggests sudden but large bursts of N loss during and immediately after a rain, rather than a small but continuous release over the entire season.

Asymbiotic N fixation does not appear to be a quantitatively important process in potential nitrogen leaching/pollution consideration in these soils.

Mineralization and denitrification appear to be the two most important N cycle processes to consider in N leaching potential in soils. Mineralization of the large pool of organic N contained in surface soils has great potential as a NO_3^- -N source and only a small change in the rate can have a profound effect on the amount of N released over a period of a year. It is know that fallowing will accelerate the process, whereas, incorporation of low N content plant residues would encourage immobilization. Addition of N containing materials should be considered over a relatively long time period (eg. several years) because of changing

mineralization/immobilization rates as carbon is evolved. Nitrification appears to be a relatively fast process, so any additions of NH_4^+ -N fertilizers would quickly be converted to NO_3^- -N which is then vulnerable to leaching. Denitrification is one process of depleting excess NO_3^- -N but requires rather specific environmental conditions.

RELATIONSHIP OF PROJECT RESULTS TO PLUARG OBJECTIVES

This project, of itself, does not answer the questions - From what sources and from what causes are pollutants contributed to ground water? Its contribution is an input that is to be used in relation to field plot studies, in order to assist the mathematical simulation of N transport and movement. The results do show the potential importance of biological transformations in the soil in leaching NO_3^- -N to the ground water. All soils studied have a high potential to convert NH_4^+ -N into NO_3^- -N, hence, any fertilizer without a nitrification inhibitor will quickly be converted into the mobile form. The rates of net mineralization, denitrification and plant uptake relative to one another over the entire biologically active season (spring, summer, fall) are important in controlling the amount of NO_3^- -N present in surface soils that can be moved downward. The very large pool of N in the organic fraction of surface soils is potentially vulnerable to leaching after mineralization-nitrification and any management practice (eg. fallowing, type of cultivation, etc.) that increases the rate of mineralization enhances the NO_3^- -N pollution potential. Fertilizer and soil mineralized N are two sources of N for plant growth and N fertilization must consider the amount of N that will be made available from soil organic N due to mineralization. Mineralizable N is generally higher in fine-textured, high organic matter soils, hence fertilization should be lower. Uptake of N by plants appears to change over the season (i.e. high initially, then lower, then increasing as flowering and seed set progress) but N mineralization could remain constant, hence the match of rate of these processes are important in potential NO_3^- -N loss. Incorporation of N containing organic amendments (straw residue, manure, sewage sludge) must be viewed in the long term (several years) since initially they may increase the soil org. N pool, and later be released by mineralization. Management practices should be viewed in the long term using these amendments. Denitrification is potentially a source of NO_3^- -N loss but specific environmental conditions are required. This process can be beneficial to strip away excess NO_3^- -N, but must also be considered in terms of the economics to plant production.

The remaining questions i.e. contribution to seasonal loadings and transmission to boundary waters, can not be addressed without knowledge of other projects, e.g. plot experiments and ground water studies.

SUPPLEMENTARY INFORMATION

If farm cultivation practices contribute to N pollution, one must be careful to isolate the exact cause. One must evaluate whether cultivation or fertilizer application is the culprit. Cultivation may accelerate N mineralization of the large soil N pool. Organic amendments must be considered in the long term, since immobilization may be important in the first year, but mineralization may be important in the next few years. A soil test, based on objective measurements, would be very beneficial to restrict NO_3^- -N leaching by encouraging efficient

fertilizer use. More must be known about basic processes of N transformations and transport under Ontario environmental conditions to enable application of an objective, measurement based N soil test.

An area that may require further consideration is whether or not cultivation has depleted the organic matter content of the soils in the watershed areas. If the organic matter has been depleted, then a considerable amount of N will have been released over the years. If this N had not been taken up by plants and if it was not denitrified, then a large amount of N could have been moved into the ground water over the cultivation period. This N would have to be quantified, to enable separation of cultivation from fertilization contributing NO_3^- -N to ground water.

Table 1. Description of surface *soil* samples from watershed 5 (IJC - PLUARG Task C - Canada)

Code	Soil Name	Land Use	Drainage	Slope.	Location	
					UTM	Sheet
N-5/1	Crombie or Maplewood	Pasture	Poor	1%	008 / 817	40P/2e
N-5/2	Honeywood or Guelph	Corn	Good	32	031 / 755	40P/2e
N-5/3	Embros	Corn (with grassy weeds)	Imperfect	1%	028 / 805	40P/2e
N-5/4	Travistock	Corn	Imperfect	1%	014 / 773	
N-5/5	Unnamed	Oats	Good	2 %	002 / 797	

Table 2 Some characterization analyses of soil samples from watersheds 1, 5 and 13 (IJC- PLUARG Task C - Canada)

Watershed	Sample	pH	%N	%C	C/N	% Moisture at		Glucose Equivalent (org.C/g)	Fixed NH ₄ -N (µg/g)
						Field Capacity	Saturation		
13	1-1	5.8	0.09	1.09	12.1	7.6	35	141	37
	1-2	5.8	0.08	0.96	11.9	8.6	36	119	30
	2-1-b	5.2	0.11	1.27	11.5	9.8	36	172	30
	2-2-b	5.5	0.11	1.29	12.9	9.7	39	163	28
	2-3	5.6	0.10	1.23	12.3	9.4	37	149	46
	2-4	6.1	0.06	0.76	12.7	7.5	37	43	32
	2-5	6.3	0.02	0.30	15.0	4.8	33	23	36
	2-5	6.2	0.03	0.24	8.0	4.2	33	27	29
	2-7	6.0	0.02	0.19	9.5	4.1	33	10	24
	2-1	5.6	0.12	1.87	15.4	11.5	36	149	18
	2-2	5.5	0.12	1.62	15.5	11.4	36	138	31
	3-1	7.7	0.07	0.94	13.4	9.6	35	116	79
	1	3-2	7.6	0.07	0.94	13.4	10.2	36	85
4-1		6.8	0.32	3.34	10.4	32.0	74	450	329
4-2		6.8	0.30	2.99	10.0	32.2	72	307	340
5-1-b		7.0	0.28	2.91	10.4	30.8	73	272	352
5-2-b		7.1	0.30	3.21	10.7	32.3	77	260	274
5-3		7.2	0.17	1.90	11.2	29.4	74	131	282
5-4		7.6	0.08	0.85	10.6	28.0	70	50	333
5-5		7.7	0.08	0.61	7.6	27.8	67	28	306
5-6		7.6	0.06	0.57	9.5	27.2	66	26	343
5-7		7.8	0.07	0.53	7.6	27.9	64	9	357
5-1		7.0	0.28	2.90	10.4	29.2	66	269	294
5-2		7.1	0.25	2.86	11.4	29.0	67	254	280
5		6-1	7.8	0.21	2.05	9.8	30.4	72	187
	6-2	7.7	0.20	1.94	9.7	31.5	69	187	342
	N-5/1	7.1	0.37	3.83	10.4	32.0	70	712	108
	N-5/2	7.3	0.24	2.50	10.4	27.0	63	338	99
	N-5/3	7.0	0.30	3.06	10.2	30.0	67	338	73
N-5/4	6.5	0.33	3.41	10.3	31.8	69	422	159	
N-5/5	5.5	0.13	1.50	11.5	23.4	55	215	61	

Meaning of computer abbreviations for Tables 3, 5, 7, 11 and 12

PL = plot number

DP = depth number (see p. 2 - Sample Code Designation)

SOL = abbreviated description of soil type: S = sand, C = clay

TMP = temperature in °C

WC = water content in % oven dry weight (30% and 75% of field capacity, (see p. 6)

TRT = treatment of N (see description of Incubation Systems for details pp 3-4)

AVER = average

SOL NH₄ = dilute CaCl₂ extractable NH₄-N

EX NH₄ = exchangeable NH₄-N (see p. 3)

TOTAL N = total extractable inorganic N

RN-14 = nitrification rate coefficient to day 14

RN-21 = nitrification rate coefficient to day 21

RN-AVE = nitrification rate coefficient averaged for 0-14 and 14-21 day periods (see p. 6)

MIN RATE = mineralization rate

RD-2 = denitrification rate coefficient to day 2

RD-5 = denitrification rate coefficient to day 5

RD-AVE = denitrification rate coefficients averaged

Table 3 Inorganic N analyses ($\mu\text{gN/g/soil}$) during incubation experiments examining nitrification and mineralization of watershed soils under combinations of temperature and moisture contents.

PL	DP	SOL	THP	WC	TRT	DAY	NO3		NO2		SOL NH4		Ex NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
1	1	S	4	2.67	0	0	4.6	14.3	0.0	0.0	0.6	1.3	0.8	1.8	17.4
1	1	S	4	2.87	0	5	1.4	20.0	0.0	0.0	0.1	0.8	0.8	2.3	23.0
1	1	S	4	2.87	0	14	0.6	18.8	0.0	0.0	0.1	0.5	1.0	1.6	20.9
1	1	S	4	2.87	0	21	0.1	19.6	0.0	0.0	1.1	0.8	0.9	1.1	21.5
1	1	S	4	2.87	80	0	1.5	23.8	0.0	0.0	1.1	56.4	4.0	23.6	103.8
1	1	S	4	2.87	80	5	2.3	22.3	0.0	0.0	0.5	61.0	3.4	26.8	110.4
1	1	S	4	2.87	80	14	0.2	22.5	0.0	0.0	2.1	61.8	0.4	24.5	108.8
1	1	S	4	2.87	80	21	0.0	23.8	0.0	0.0	2.1	58.6	0.6	26.8	109.2
1	1	S	4	7.19	0	0	0.8	15.5	0.0	0.0	0.4	0.6	0.7	2.8	18.8
1	1	S	4	7.19	0	5	0.5	21.9	0.0	0.0	0.1	0.6	0.0	2.5	25.1
1	1	S	4	7.19	0	14	10.5	16.4	0.0	0.0	1.1	0.8	1.2	1.6	18.9
1	1	S	4	7.19	0	21	1.6	20.4	0.0	0.0	1.6	2.5	0.8	2.0	24.9
1	1	S	4	7.19	80	0	3.3	27.9	0.0	0.0	3.4	58.6	0.6	25.0	111.5
1	1	S	4	7.19	80	5	0.9	19.8	0.0	0.0	1.1	68.1	0.6	25.3	111.3
1	1	S	4	7.19	80	14	0.5	28.4	0.0	0.0	0.6	58.4	0.6	25.2	111.9
1	1	S	4	7.19	80	21	0.9	25.9	0.0	0.0	0.5	59.4	1.6	25.0	110.4
1	1	S	15	2.87	0	0	4.6	14.3	0.0	0.0	0.6	1.3	0.8	1.8	17.4
1	1	S	15	2.87	0	5	0.2	21.6	0.0	0.0	0.6	0.8	1.8	2.6	25.0
1	1	S	15	2.87	0	14	0.1	17.8	0.0	0.0	0.7	0.8	0.7	1.1	19.7
1	1	S	15	2.87	0	21	1.7	19.3	0.0	0.0	0.0	0.3	0.1	2.1	21.8
1	1	S	15	2.87	80	0	1.5	23.8	0.0	0.0	1.1	56.4	4.0	23.6	103.8
1	1	S	15	2.87	80	5	1.9	23.9	0.0	0.0	5.7	59.9	1.3	26.3	110.1
1	1	S	15	2.87	80	14	0.8	24.4	0.0	0.0	1.5	57.3	0.7	26.3	108.0
1	1	S	15	2.67	80	21	0.1	23.3	0.0	0.0	3.7	54.6	1.2	26.3	104.2
1	1	S	15	7.19	0	0	0.8	15.5	0.0	0.0	0.4	0.6	0.7	2.8	18.8
1	1	S	15	7.19	0	5	0.8	24.8	0.0	0.0	1.2	1.3	0.2	2.0	28.1
1	1	S	15	7.19	0	14	13.1	16.6	7.0	3.5	0.3	0.5	0.5	1.3	21.8
1	1	S	15	7.19	0	21	4.1	14.8	0.0	0.0	0.9	1.6	0.1	2.4	18.8
1	1	S	15	7.19	80	0	3.3	27.9	0.0	0.0	3.4	58.6	0.6	25.0	111.5
1	1	S	15	7.19	80	5	4.3	26.3	0.0	0.0	5.9	62.1	1.5	26.4	114.8
1	1	S	15	7.19	80	14	3.0	29.6	0.0	0.0	0.0	57.1	1.3	25.8	112.6
1	1	S	15	7.19	80	21	5.2	31.9	0.0	0.0	6.2	49.1	1.6	25.2	106.2
1	1	S	30	2.87	0	0	4.6	14.3	0.0	0.0	0.6	1.3	0.6	1.8	17.4
1	1	S	30	2.87	0	5	3.2	17.1	0.0	0.0	0.1	0.8	0.6	2.0	19.9
1	1	S	30	2.87	0	14	0.2	15.2	0.0	0.0	0.1	0.3	0.1	2.3	17.8
1	1	S	30	2.87	0	21	2.5	10.8	0.0	0.0	0.1	0.4	0.1	2.1	13.3
1	1	S	30	2.87	80	0	1.5	23.8	0.0	0.0	1.1	56.4	4.0	23.6	103.8
1	1	S	30	2.87	80	5	3.1	26.6	0.0	0.0	5.3	54.3	5.2	28.5	109.5
1	1	S	30	2.87	80	14	0.0	23.8	0.0	0.0	0.0	50.9	0.9	27.3	102.0
1	1	S	30	2.87	80	21	0.2	24.4	0.0	0.0	3.1	45.9	2.4	25.9	96.2
1	1	S	30	7.19	0	0	0.8	15.5	0.0	0.0	0.4	0.6	0.7	2.8	18.8
1	1	S	30	7.19	0	5	1.0	24.6	0.0	0.0	0.3	1.0	0.5	2.6	28.2
1	1	S	30	7.19	0	14	1.9	24.8	4.0	2.0	0.0	0.3	0.7	2.0	29.1
1	1	S	30	7.19	0	21	1.9	18.8	0.0	0.0	0.0	0.3	0.1	3.3	22.3
1	1	S	30	7.19	80	0	3.3	27.9	0.0	0.0	3.4	58.6	0.6	25.0	111.5
1	1	S	30	7.19	80	5	1.0	36.2	0.0	0.0	2.3	54.3	0.5	24.3	114.8
1	1	S	30	7.19	80	14	6.6	51.7	0.0	0.0	4.8	37.1	1.0	22.0	110.8
1	1	S	30	7.19	80	21	0.9	66.8	0.0	0.0	2.0	19.8	0.6	15.4	101.9
1	2	S	4	2.60	0	0	4.9	25.6	0.0	0.0	3.5	11.1	0.0	0.0	36.9
1	2	S	4	2.60	0	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1	2	S	4	2.60	0	14	1.6	18.9	0.0	0.0	0.5	10.5	0.0	0.0	29.4
1	2	S	4	2.60	0	21	1.1	21.3	0.0	0.0	0.2	6.0	0.0	0.0	27.3
1	2	S	4	2.60	80	0	0.4	25.3	0.0	0.0	4.0	88.4	0.0	0.0	113.7
1	2	S	4	2.60	80	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1	2	S	4	2.60	80	14	2.0	34.8	0.0	0.0	0.3	84.1	0.0	0.0	118.9
1	2	S	4	2.60	80	21	8.2	37.7	0.0	0.0	0.6	75.5	0.0	0.0	113.2
1	2	S	4	6.50	0	0	1.7	24.6	0.0	0.0	0.4	6.2	0.0	0.0	31.0
1	2	S	4	6.50	0	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1	2	S	4	6.50	0	14	0.3	17.9	0.0	0.0	0.9	13.9	0.0	0.0	31.9
1	2	S	4	6.50	0	21	0.1	21.4	0.0	0.0	0.2	10.5	0.0	0.0	31.9
1	2	S	4	6.50	80	0	7.8	22.0	0.0	0.0	10.2	94.1	0.0	0.0	116.1
1	2	S	4	6.50	80	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1	2	S	4	6.50	80	14	6.0	34.0	0.0	0.0	5.2	89.1	0.0	0.0	123.1
1	2	S	4	6.50	80	21	4.7	33.5	0.0	0.0	3.1	83.8	0.0	0.0	117.3

Table 3																
PL	DP	SOL	TMP	MC	TRT	DAY	NO3		NO2		SOL NH4		EX NH4		TOTAL N	
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER		
1	2	S	15	2.60	0	0	4.9	25.8	5.0	0.0	3.5	11.1	0.0	0.0	36.9	
1	2	S	15	2.60	0	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1	2	S	15	2.60	0	14	0.1	18.4	0.0	0.0	0.0	9.0	0.0	0.0	27.4	
1	2	S	15	2.60	0	21	0.6	20.9	0.0	0.0	0.2	6.0	0.0	0.0	26.9	
1	2	S	15	2.60	80	0	0.4	25.3	0.0	0.0	4.0	88.4	0.0	0.0	113.7	
1	2	S	15	2.60	80	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1	2	S	15	2.60	80	14	3.2	29.7	0.0	0.0	0.6	80.9	0.0	0.0	110.6	
1	2	S	15	2.60	80	21	1.0	29.3	0.0	0.0	1.6	77.5	0.0	0.0	106.8	
1	2	S	15	6.50	0	0	1.7	24.8	0.0	0.0	0.4	6.2	0.0	0.0	31.0	
1	2	S	15	6.50	0	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1	2	S	15	6.50	0	14	6.2	15.3	0.0	0.0	0.9	15.7	0.0	0.0	30.9	
1	2	S	15	6.50	0	21	1.7	19.9	0.0	0.0	0.4	12.9	0.0	0.0	32.8	
1	2	S	15	6.50	80	0	7.8	22.0	0.0	0.0	10.2	94.1	0.0	0.0	116.1	
1	2	S	15	6.50	80	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1	2	S	15	6.50	80	14	3.4	33.8	0.0	0.0	1.1	85.3	0.0	0.0	118.8	
1	2	S	15	6.50	80	21	6.6	38.1	0.0	0.0	7.4	91.0	0.0	0.0	129.1	
1	2	S	30	2.60	0	0	4.9	25.8	0.0	0.0	3.5	11.1	0.0	0.0	36.9	
1	2	S	30	2.60	0	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1	2	S	30	2.60	0	14	2.0	19.2	0.0	0.0	0.2	11.1	0.0	0.0	30.3	
1	2	S	30	2.60	0	21	0.9	20.6	0.0	0.0	0.4	7.2	0.0	0.0	27.8	
1	2	S	30	2.60	80	0	0.4	25.3	0.0	0.0	4.0	88.4	0.0	0.0	113.7	
1	2	S	30	2.60	80	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1	2	S	30	2.60	80	14	5.1	29.8	0.0	0.0	1.7	83.1	0.0	0.0	112.9	
1	2	S	30	2.60	80	21	7.6	31.4	0.0	0.0	2.3	81.4	0.0	0.0	112.8	
1	2	S	30	6.50	0	0	1.7	24.8	0.0	0.0	0.4	6.2	0.0	0.0	31.0	
1	2	S	30	6.50	0	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1	2	S	30	6.50	0	14	0.3	16.6	0.0	0.0	0.3	22.5	0.0	0.0	39.2	
1	2	S	30	6.80	0	21	2.2	21.3	0.0	0.0	0.1	20.3	0.6	0.0	41.6	
1	2	S	30	6.50	80	0	7.8	22.0	0.0	0.0	10.2	94.1	0.0	0.0	116.1	
1	2	S	30	6.50	80	5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1	2	S	30	6.50	80	14	4.2	35.3	0.0	0.0	1.0	99.5	0.0	0.0	134.8	
1	2	S	30	6.50	80	21	1.1	38.6	0.0	0.0	1.1	96.8	0.0	0.0	135.3	
2	1	S	4	3.45	0	0	1.1	7.6	0.0	0.0	0.1	0.1	0.6	2.9	10.7	
2	1	S	4	3.45	0	5	1.4	10.9	0.0	0.0	0.1	1.9	0.7	3.1	16.0	
2	1	S	4	3.45	0	14	0.6	9.8	0.0	0.0	0.5	1.3	0.2	2.6	13.6	
2	1	S	4	3.45	0	21	0.3	10.0	0.0	0.0	0.6	1.9	0.4	2.4	14.3	
2	1	S	4	3.45	80	0	6.2	15.2	0.0	0.0	1.6	55.5	0.6	25.0	96.3	
2	1	S	4	3.45	80	5	3.2	17.7	0.0	0.0	4.2	58.9	0.2	27.4	104.0	
2	1	S	4	3.45	80	14	6.4	15.2	0.0	0.0	0.0	56.8	0.9	25.6	97.6	
2	1	S	4	3.45	80	21	5.8	22.9	0.0	0.0	9.0	63.4	1.6	29.1	115.4	
2	1	S	4	8.63	0	0	1.0	11.7	0.0	0.0	0.1	0.6	0.2	3.1	15.4	
2	1	S	4	8.63	0	5	0.2	10.9	0.0	0.0	0.3	0.3	0.2	3.2	14.3	
2	1	S	4	8.63	0	14	0.0	12.5	0.0	0.0	0.1	0.8	0.3	2.5	15.9	
2	1	S	4	8.63	0	21	0.3	14.5	0.0	0.0	0.3	0.9	0.1	1.3	16.3	
2	1	S	4	8.63	80	0	3.4	12.2	0.0	0.0	3.8	56.2	4.2	27.9	96.3	
2	1	S	4	8.63	80	5	1.2	18.2	0.0	0.0	2.2	58.6	0.3	27.3	104.0	
2	1	S	4	8.63	80	14	0.2	22.5	0.0	0.0	0.0	57.5	0.5	25.9	105.9	
2	1	S	4	8.63	80	21	7.2	20.8	0.0	0.0	17.1	65.6	5.3	2d.6	115.0	
2	1	S	15	3.45	0	0	1.1	7.6	0.0	0.0	0.1	0.1	0.6	2.9	10.7	
2	1	S	15	3.45	0	5	3.1	9.3	0.0	0.0	1.4	0.8	0.3	2.5	12.6	
2	1	S	15	3.45	0	14	0.5	9.8	0.0	0.0	0.3	1.4	0.0	2.6	13.8	
2	1	S	15	3.45	0	21	0.3	9.6	0.0	0.0	0.1	1.6	0.6	2.6	13.9	
2	1	S	15	3.45	80	0	6.2	15.2	0.0	0.0	1.6	55.5	0.6	25.6	96.3	
2	1	S	15	3.45	80	5	6.1	17.1	0.0	0.0	2.6	55.0	1.3	23.8	100.9	
2	1	S	15	3.45	80	14	0.6	17.9	0.0	0.0	0.5	55.6	0.9	27.3	100.7	
2	1	S	15	3.45	80	21	3.1	17.6	0.0	0.0	0.0	54.7	1.1	26.8	99.2	
2	1	S	15	8.63	0	0	1.0	11.7	0.0	0.0	0.1	0.6	0.2	3.1	15.4	
2	1	S	15	8.63	0	5	2.8	12.5	0.0	0.0	1.2	0.7	0.1	1.4	14.6	
2	1	S	15	8.63	0	14	0.5	13.a	0.0	0.0	0.0	0.1	0.1	0.8	14.6	
2	1	S	15	8.63	0	21	0.4	14.6	0.0	0.0	0.2	0.2	0.0	0.5	15.3	
2	1	S	15	8.63	80	0	3.4	12.2	0.0	0.0	3.8	56.2	4.2	27.9	96.3	
2	1	S	15	6.63	An	5	2.5	22.9	0.0	0.0	1.6	55.1	1.1	26.4	104.5	
2	1	S	15	8.63	80	14	6.7	32.5	0.0	0.0	4.4	46.5	22.1	23.3	102.6	
2	1	S	15	8.63	80	21	9.0	42.0	0.0	0.0	2.1	37.6	0.3	19.8	99.6	
2	1	S	30	3.45	0	0	1.1	7.6	0.0	0.0	0.1	0.1	0.0	2.9	10.7	
2	1	S	30	3.45	0	5	2.6	8.8	0.0	0.0	0.9	0.7	1.0	2.3	11.7	
2	1	S	30	3.45	0	14	0.8	9.6	0.0	0.0	0.2	0.9	0.0	109	12.4	
2	1	S	30	3.45	0	21	1.5	10.1	0.0	0.0	0.6	1.4	0.0	1.5	13.0	
2	1	S	30	3.45	80	0	6.2	15.2	0.0	0.0	1.6	55.5	0.6	25.6	90.3	
2	1	S	30	3.45	80	5	0.8	19.1	0.0	0.0	2.2	50.5	0.8	29.3	98.9	
2	1	S	30	3.45	80	14	2.8	14.9	0.0	0.0	3.1	97.3	0.9	29.8	107.1	
2	1	S	30	3.45	80	21	2.1	17.3	0.0	0.0	2.6	53.9	0.4	29.7	100.8	
2	1	S	30	8.63	0	0	1.0	11.7	0.0	0.0	0.1	0.6	0.2	3.1	15.4	
2	1	S	30	8.63	0	5	2.2	15.8	0.0	0.0	1.3	0.7	0.5	114	15.9	
2	1	S	30	8.63	0	14	0.3	15.3	0.0	0.0	0.1	0.1	0.1	0.6	16.1	

2	1	S	30	8.63	0	21	0.2	15.9	0.0	0.0	0.0	0.9	0.2	1.0	17.8
2	1	S	30	8.63	80	0	3.4	12.2	0.0	0.0	3.8	56.2	4.2	27.9	96.3
2	1	S	30	8.63	80	5	0.3	35.0	0.0	0.0	1.9	47.1	0.6	20.0	100.2
2	1	S	30	6.63	80	14	5.5	52.4	0.0	0.0	1.1	29.8	1.2	10.1	100.3
2	1	S	30	8.63	80		22	47.9	0.0		0.0				

PL	OP	SOL	IMP	WC	TRT	DAY	NO3		NO2		SOL NH4		EX NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
2	2	S	4	3.42	0	0	3.4	17.1	0.0	0.0	0.1	2.1	0.1	4.3	23.5
2	2	S	4	3.42	0	5	0.8	17.2	0.0	0.0	1.8	1.1	0.1	5.1	23.3
2	2	S	4	3.42	0	14	0.6	18.3	0.0	0.0	0.1	2.4	0.3	5.3	26.1
2	2	S	4	3.42	0	21	2.3	16.8	0.0	0.0	0.0	2.9	2.0	5.0	24.7
2	2	S	4	3.42	80	0	1.3	14.9	0.0	0.0	3.2	56.8	2.6	27.2	98.9
2	2	S	4	3.42	80	5	11.2	19.2	0.0	0.0	4.2	57.3	1.2	30.5	107.0
2	2	S	4	3.42	80	14	0.9	18.6	0.0	0.0	1.1	61.6	0.2	28.3	108.4
2	2	S	4	3.42	80	21	14.6	21.4	0.0	0.0	3.2	62.0	1.4	28.8	112.2
2	2	S	4	8.55	0	0	1.3	17.4	0.0	0.0	0.6	0.6	0.1	5.1	23.2
2	2	S	4	8.55	0	5	1.3	17.4	0.0	0.0	1.3	0.8	0.1	6.1	24.3
2	2	S	4	8.55	0	14	2.0	18.3	0.0	0.0	0.1	2.8	0.2	5.1	26.2
2	2	S	4	8.55	0	21	3.1	18.6	0.0	0.0	0.3	2.9	1.0	4.9	26.5
2	2	S	4	8.55	80	0	3.3	20.0	0.0	0.0	1.1	62.8	2.3	27.5	110.4
2	2	S	4	8.55	80	5	0.4	18.4	0.0	0.0	2.6	61.0	1.0	29.7	109.1
2	2	S	4	8.55	80	14	3.7	15.9	0.0	0.0	1.6	63.1	0.2	29.1	108.1
2	2	S	4	8.55	80	21	8.4	22.2	0.0	0.0	1.1	63.4	0.1	26.3	114.0
2	2	S	15	3.42	0	0	3.4	17.1	0.0	0.0	0.1	2.1	0.1	4.3	23.5
2	2	S	15	3.42	0	5	0.7	15.3	0.0	0.0	1.5	0.9	0.2	4.7	21.0
2	2	S	15	3.42	0	14	0.2	14.9	0.0	0.0	0.1	2.6	0.6	4.4	21.9
2	2	S	15	3.42	0	21	1.0	15.4	0.0	0.0	0.7	2.9	0.7	3.5	21.9
2	2	S	15	3.42	80	0	1.3	14.9	0.0	0.0	3.2	56.8	2.6	27.2	98.9
2	2	S	15	3.42	80	5	6.8	23.7	0.0	0.0	0.1	60.6	2.3	30.0	114.3
2	2	S	15	3.42	80	14	0.5	22.9	0.0	0.0	0.5	64.1	0.7	29.6	116.6
2	2	S	15	3.42	80	21	5.4	25.4	0.0	0.0	1.1	61.1	1.3	29.6	116.1
2	2	S	15	8.55	0	0	1.3	17.4	0.0	0.0	0.6	0.6	0.1	5.1	23.2
2	2	S	15	8.55	0	5	0.2	17.4	0.0	0.0	0.5	1.9	0.1	5.1	24.4
2	2	S	15	8.55	0	14	1.4	16.9	0.0	0.0	0.7	2.1	0.5	3.6	22.6
2	2	S	15	8.55	0	21	4.3	21.9	0.0	0.0	1.1	0.9	0.8	2.1	25.0
2	2	S	15	8.55	80	0	3.3	20.0	0.0	0.0	1.1	62.8	2.3	27.5	110.4
2	2	S	15	8.55	80	5	0.4	20.6	0.0	0.0	1.1	64.1	1.3	31.0	115.9
2	2	S	15	8.55	80	14	7.4	30.2	0.0	0.0	1.0	61.3	1.9	28.8	120.3
2	2	S	15	8.55	80	21	12.9	34.6	0.0	0.0	2.2	59.7	0.5	27.8	122.1
2	2	S	30	3.42	0	0	3.4	17.1	0.0	0.0	0.1	2.1	0.1	4.3	23.5
2	2	S	30	3.42	0	5	0.1	16.6	0.0	0.0	1.0	1.8	0.2	3.9	22.2
2	2	S	30	3.42	0	14	0.0	16.0	0.0	0.0	0.0	2.4	0.2	3.2	21.6
2	2	S	30	3.42	0	21	0.1	15.8	0.0	0.0	0.6	3.3	0.4	2.9	21.9
2	2	S	30	3.42	80	0	1.3	14.9	0.0	0.0	3.2	56.8	2.6	27.2	98.9
2	2	S	30	3.42	80	5	5.3	26.0	0.0	0.0	3.3	55.1	2.8	28.6	109.8
2	2	S	30	3.42	80	14	7.4	29.2	0.0	0.0	1.1	56.8	2.4	28.8	114.7
2	2	S	30	3.42	80	21	0.0	28.4	0.0	0.0	2.0	58.5	0.0	30.0	116.9
2	2	S	30	8.55	0	0	1.3	17.4	0.0	0.0	0.6	0.6	0.1	5.1	23.2
2	2	S	30	8.55	0	5	0.4	21.4	0.0	0.0	0.3	1.0	0.3	2.4	24.9
2	2	S	30	8.55	0	14	1.5	24.6	0.0	0.0	0.3	0.3	0.0	0.7	25.6
2	2	S	30	8.55	0	21	1.0	24.8	0.0	0.0	0.2	0.6	0.7	0.8	26.2
2	2	S	30	8.55	80	0	3.3	20.0	0.0	0.0	1.1	62.8	2.3	27.5	110.4
2	2	S	30	8.55	80	5	7.7	37.6	0.0	0.0	1.7	57.8	16.4	29.0	124.4
2	2	S	30	8.55	80	14	11.8	44.3	0.0	0.0	9.8	50.9	5.2	26.8	122.0
2	2	S	30	8.55	80	21	3.2	51.8	0.0	0.0	0.6	42.2	0.2	22.7	116.7
2	1	SB	4	2.94	0	0	1.0	58.6	0.0	0.0	0.1	0.1	0.8	2.7	61.4
2	1	SB	4	2.94	0	5	0.6	56.7	0.0	0.0	0.5	0.5	0.3	3.1	60.4
2	1	SB	4	2.94	0	14	0.3	59.6	0.5	0.3	0.6	0.8	0.6	6.7	67.4
2	1	SB	4	2.94	0	21	1.4	59.2	0.0	0.0	0.1	0.1	0.8	3.6	62.9
2	1	SB	4	2.94	80	0	0.5	69.1	0.0	0.0	3.9	51.8	3.5	23.6	144.4
2	1	SB	4	2.94	80	5	2.6	69.1	0.0	0.0	0.6	53.7	0.4	26.6	149.6
2	1	SB	4	2.94	80	14	0.2	68.3	0.0	0.0	1.8	57.7	3.0	30.2	156.2
2	1	SB	4	2.94	80	21	0.7	66.8	0.0	0.0	1.0	53.9	1.1	20.8	149.5
2	1	SB	4	7.34	0	0	0.2	58.8	0.0	0.0	0.5	0.4	1.1	3.4	62.6
2	1	SB	4	7.34	0	5	1.5	61.3	0.0	0.0	0.0	0.1	0.1	3.8	65.1
2	1	SB	4	7.34	0	14	0.2	60.6	0.0	0.0	0.7	0.5	0.9	3.8	65.0
2	1	SB	4	7.34	0	21	0.8	61.8	0.0	0.0	0.1	0.1	0.9	3.3	65.3
2	1	SB	4	7.34	80	0	0.2	68.2	0.0	0.0	0.9	55.6	0.4	26.6	150.3
2	1	SB	4	7.34	80	5	0.3	69.8	0.0	0.0	1.3	56.5	0.4	27.0	153.3
2	1	SB	4	7.34	80	14	4.3	67.3	0.0	0.0	5.2	59.0	0.2	26.3	152.5
2	1	SB	4	7.34	80	21	23.2	58.0	0.0	0.0	16.3	46.9	10.9	20.7	125.7
2	1	SB	15	2.94	0	0	1.0	58.6	0.0	0.0	0.1	0.1	0.8	2.7	61.4
2	1	SB	15	2.94	0	5	1.6	57.9	0.0	0.0	0.0	0.1	1.2	3.4	61.4
2	1	SB	15	2.94	0	14	1.5	57.6	7.0	6.0	0.7	1.1	0.8	3.2	67.9
2	1	SB	15	2.94	0	21	0.3	61.4	0.0	0.0	0.0	0.1	0.4	2.7	64.2
2	1	SB	15	2.94	80	0	0.5	69.1	0.0	0.0	3.9	51.8	3.5	23.6	144.4

2	1	SB	15	2.94	80	5	0.1	68.6	0.0	0.0	3.9	54.1	2.2	26.6	149.4
2	1	SB	15	2.94	80	14	1.4	68.0	1.0	0.5	7.0	55.7	3.4	27.9	152.1
2	1	SB	15	2.94	80	21	3.0	66.9	0.0	0.0	0.5	51.9	0.6	27.2	145.9
2	1	SB	15	7.34	0	0	0.2	58.8	0.0	0.0	0.5	0.4	1.1	3.4	62.6
2	1	SB	15	7.34	0	5	3.8	58.1	0.0	0.0	1.4	0.8	0.2	2.6	61.7
2	1	SB	15	7.34	0	14	3.4	64.0	1.0	0.5	0.2	0.2	0.2	3.5	68.2
2	1	SB	15	7.34	0	21	0.8	64.2	0.0	0.0	0.0	0.1	1.3	2.9	67.2
2	1	SB	15	7.34	80	0	0.2	68.2	0.0	0.0	0.9	55.6	0.4	26.6	150.3
2	1	SB	15	7.34	80	5	0.8	68.8	0.0	0.0	0.3	55.3	0.4	26.7	150.7
2	1	SB	15	7.34	80	14	0.5	65.6	0.0	0.0	3.3	56.4	0.0	27.8	152.8
2	1	SB	15	7.34	80	21	1.6	67.0	0.0	0.0	5.6	50.4	1.6	30.3	143.7

PL	DP	SOL	IMP	WC	TRT	DAY	NO3		NO2		SOL NH4		Ex NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
2	1	SB	30	2.94	0	0	1.0	58.6	0.0	0.0	0.1	0.1	0.8	2.7	61.4
2	1	SB	30	2.94	0	5	2.2	61.1	0.0	0.0	0.0	0.1	0.5	4.1	65.2
2	1	SB	30	2.94	0	14	0.9	58.6	0.0	0.0	0.6	0.5	0.0	3.6	62.7
2	1	SB	30	2.94	0	21	1.2	61.4	0.0	0.0	0.4	0.3	1.3	4.4	60.1
2	1	SB	30	2.94	80	0	0.5	69.1	0.0	0.0	3.9	51.8	3.5	23.6	144.4
2	1	SB	30	2.94	80	5	2.9	67.1	0.0	0.0	1.7	51.9	3.0	29.3	148.4
2	t	SB	30	2.94	80	14	13.1	57.8	0.5	0.3	0.3	51.1	3.3	31.0	140.3
2	1	SB	30	2.94	80	21	11.5	74.4	0.0	0.0	0.9	51.8	0.2	28.6	154.8
2	1	SB	30	7.34	0	0	0.2	58.8	0.0	0.0	0.5	0.4	1.1	3.4	62.6
2	1	SB	30	7.34	0	5	3.1	63.8	0.0	0.0	0.0	0.2	2.2	4.4	68.3
2	1	SB	30	7.34	0	14	0.1	62.8	0.0	0.0	0.0	0.1	0.3	3.8	66.7
2	1	SB	30	7.34	0	21	0.0	63.9	0.0	0.0	0.1	0.1	1.2	4.5	68.5
2	1	SB	30	7.34	80	0	0.2	68.2	0.0	0.0	0.9	55.6	0.4	26.6	150.3
2	1	SB	30	7.34	80	5	0.6	70.1	0.0	0.0	3.5	53.4	0.2	26.5	150.0
2	1	SB	30	7.34	80	14	2.9	71.3	0.0	0.0	0.3	53.5	7.2	34.5	159.3
2	1	SB	30	7.34	80	21	2.4	69.8	0.0	0.0	0.1	51.3	0.1	29.1	150.2
2	2	SB	4	2.91	0	0	1.0	22.3	0.0	0.0	0.0	0.1	0.1	3.1	25.5
2	2	SB	4	2.91	0	5	2.7	20.5	0.0	0.0	0.0	0.2	1.4	2.2	22.9
2	2	SB	4	2.91	0	14	1.7	22.3	0.0	0.0	0.1	0.1	0.2	1.3	23.8
2	2	SB	4	2.91	0	21	1.8	22.1	0.0	0.0	0.6	0.4	0.0	1.2	23.7
2	2	SB	4	2.91	80	0	0.8	20.1	0.0	0.0	0.5	52.4	0.6	28.1	100.5
2	2	SB	4	2.91	80	5	2.5	23.6	0.0	0.0	3.6	53.9	0.2	26.7	104.1
2	2	SB	4	2.91	80	14	5.1	22.1	0.0	0.0	0.0	52.1	1.4	26.8	101.0
2	2	SB	4	2.91	80	21	2.5	24.9	0.0	0.0	2.0	52.6	0.6	26.2	103.7
2	2	SB	4	7.28	0	0	1.4	21.1	0.0	0.0	3.0	2.0	0.5	2.1	25.1
2	2	SB	4	7.28	0	5	4.0	21.6	0.0	0.0	0.9	0.5	0.2	1.7	23.8
2	2	SB	4	7.28	0	14	1.7	22.8	0.0	1.5	1.2	0.8	1.2	1.3	26.3
2	2	SB	4	7.28	0	21	1.0	24.0	0.0	0.0	0.1	0.3	0.9	1.2	25.6
2	2	SB	4	7.28	80	0	4.3	23.3	0.0	0.0	1.0	53.7	0.2	28.0	104.9
2	2	SB	4	7.28	80	5	1.9	25.1	0.0	0.0	0.3	51.3	1.1	26.8	103.1
2	2	SB	4	7.28	80	14	4.3	24.5	0.5	0.3	2.1	53.1	2.0	26.1	104.0
2	2	SB	4	7.28	80	21	7.1	26.8	0.0	0.0	0.7	50.3	0.0	28.3	105.3
2	2	SB	15	2.91	0	0	1.0	22.3	0.0	0.0	0.0	0.1	0.1	3.1	25.5
2	2	SB	15	2.91	0	5	1.6	21.2	0.0	0.0	0.4	0.3	0.7	1.8	23.2
2	2	SB	15	2.91	0	14	0.5	23.1	0.0	0.0	0.0	0.1	0.1	1.6	24.7
2	2	SB	15	2.91	0	21	0.8	23.8	0.0	0.0	0.1	0.3	0.1	1.6	25.6
2	2	SB	15	2.91	80	0	0.8	20.1	0.0	0.0	0.5	52.4	0.6	28.1	100.5
2	2	SB	15	2.91	80	5	3.2	25.8	0.0	0.0	0.2	50.9	4.6	29.4	106.1
2	2	SB	15	2.91	80	14	7.8	24.1	0.0	0.0	4.3	47.8	2.5	30.4	102.4
2	2	SB	15	2.91	80	21	6.8	20.6	0.0	0.0	0.2	50.6	0.2	28.2	99.4
2	2	SB	15	7.28	0	0	1.4	21.1	0.0	0.0	3.0	2.0	0.0	2.1	25.1
2	2	SB	15	7.28	0	5	0.3	21.9	0.0	0.0	0.4	0.3	0.3	1.3	23.6
2	2	SB	15	7.28	0	14	1.6	24.5	0.0	0.0	0.2	0.2	3.9	4.6	29.3
2	2	SB	15	7.28	0	21	0.3	26.6	0.0	0.0	0.0	0.3	0.2	1.2	28.1
2	2	SB	15	7.28	80	0	4.3	23.3	0.0	0.0	1.0	53.7	0.2	28.0	104.9
2	2	SB	15	7.28	80	5	5.1	27.9	0.0	0.0	0.1	49.9	0.6	27.6	105.5
2	2	SB	15	7.28	80	14	5.6	27.1	0.0	0.0	0.1	49.1	1.1	29.8	106.0
2	2	SB	15	7.26	80	21	4.6	35.6	0.0	0.0	3.6	45.6	1.2	24.0	105.8
2	2	SB	30	2.91	0	0	1.0	22.3	0.0	0.0	0.0	0.1	0.1	3.1	25.5
2	2	SB	30	2.91	n	5	2.1	21.1	0.0	0.0	0.1	0.3	0.1	1.6	22.8
2	2	SB	30	2.91	0	14	1.3	22.1	0.0	0.0	0.0	0.2	0.6	3.3	25.6
2	2	SB	30	2.91	0	21	1.4	22.1	0.0	0.0	0.3	0.3	0.1	1.8	24.2
2	2	SB	30	2.91	80	0	0.8	20.1	0.0	0.0	0.5	52.4	0.6	28.1	100.5
2	2	SB	30	2.91	80	5	3.2	24.5	0.0	0.0	1.9	48.4	1.1	28.3	101.2
2	2	SB	30	2.91	80	14	4.0	22.1	0.0	0.0	3.0	51.1	1.3	26.8	100.1
2	2	SB	30	2.91	80	21	3.0	25.5	0.0	0.0	1.0	49.0	0.6	26.7	101.2
2	2	SB	30	7.28	0	0	1.4	21.1	0.0	0.0	3.0	2.0	0.5	2.1	25.1
2	2	SB	30	7.28	0	5	0.8	25.6	0.0	0.0	0.1	0.1	0.1	2.3	26.0
2	2	SB	30	7.28	0	14	5.0	23.9	0.0	0.0	0.1	0.3	1.2	2.5	26.6
2	2	SB	30	7.28	0	21	1.2	25.4	0.0	0.0	0.0	0.1	0.1	109	27.4
2	2	SB	30	7.28	80	0	4.3	23.3	0.0	0.0	1.0	53.7	0.2	28.0	104.9
2	2	SB	30	7.28	80	5	0.9	43.4	0.0	0.0	0.9	47.3	19.5	36.6	127.3
2	2	SB	30	7.28	80	14	7.4	39.3	0.0	0.0	0.0	44.8	0.5	28.3	110.4
2	2	SB	30	7.28	80	21	7.9	44.4	0.0	0.0	1.1	42.8	0.7	23.1	110.3

Table 3 cont'd.

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		SOL NH4		EX NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
2	3	S	4	2.81	0	0	1.6	40.4	0.0	0.0	0.1	0.3	0.8	3.4	24.0
2	3	S	4	2.81	0	5	0.2	20.9	0.0	0.0	0.1	0.3	0.0	2.6	23.7
2	3	S	4	2.81	0	10	1.2	19.3	0.0	0.0	0.5	0.8	0.3	1.8	21.8
2	3	S	4	2.81	0	21	0.8	20.6	0.0	0.0	0.0	0.0	0.0	1.4	22.0
2	3	S	4	2.81	80	0	4.9	18.8	0.0	0.0	3.9	51.8	1.3	26.0	96.5
2	3	S	4	2.61	80	S	5.6	17.9	0.0	0.0	4.6	55.6	0.1	25.9	99.6
2	3	S	4	2.81	80	14	8.7	13.4	3.3	1.6	0.4	53.1	2.5	26.3	94.4
2	3	S	4	2.81	80	21	1.4	10.8	0.0	0.0	0.4	56.6	0.3	24.8	98.1
2	3	S	4	7.04	0	0	0.2	20.7	0.0	0.0	0.6	0.6	1.9	3.1	24.3
2	3	S	4	7.04	0	5	2.7	20.4	0.0	0.0	1.3	1.4	1.9	3.1	24.9
2	3	S	4	7.04	0	14	0.1	19.8	0.0	0.0	0.2	0.9	0.0	2.4	23.1
2	3	S	4	7.04	0	21	0.3	21.1	0.0	0.0	0.1	0.6	0.1	1.1	22.9
2	3	S	4	7.04	80	0	2.7	17.9	0.0	0.0	4.5	58.1	1.3	26.0	102.1
2	3	S	4	7.04	80	5	1.6	20.4	0.0	0.0	0.2	55.6	2.3	26.6	102.6
2	3	S	4	7.04	80	14	3.1	19.8	0.0	0.0	1.6	52.8	0.9	24.6	97.2
2	3	S	4	7.04	80	21	5.7	24.4	0.0	0.0	2.5	52.1	0.1	22.8	99.3
2	3	S	15	2.81	0	0	1.6	20.4	0.0	0.0	0.1	0.3	0.6	3.4	24.0
2	3	S	15	2.81	0	5	0.5	20.9	0.0	0.0	0.1	0.1	0.3	2.1	23.1
2	3	S	15	2.61	0	14	0.3	19.3	0.0	0.0	0.5	0.4	0.1	1.6	21.3
2	3	S	15	2.81	0	21	1.4	20.6	0.0	0.0	0.0	0.1	0.6	2.3	23.0
2	3	S	15	2.81	80	0	4.9	18.8	0.0	0.0	3.9	51.8	1.3	26.0	96.5
2	3	S	15	2.81	80	5	1.6	18.4	0.0	0.0	1.9	53.6	7.9	27.8	99.8
2	3	S	15	2.81	80	14	3.1	23.1	0.0	0.0	2.4	49.1	0.6	25.2	97.4
2	3	S	15	2.81	80	21	3.7	19.4	0.0	0.0	2.4	52.5	0.6	25.4	97.3
2	3	S	15	7.04	0	0	0.2	20.7	0.0	0.0	0.8	0.6	1.9	3.1	24.3
2	3	S	15	7.04	0	5	1.2	20.9	0.0	0.0	0.3	0.3	0.7	1.8	23.0
2	3	S	15	7.04	0	10	4.2	19.7	0.0	0.0	0.4	0.3	0.6	1.7	21.7
2	3	S	15	7.04	0	21	0.6	23.5	0.0	0.0	0.1	0.3	0.2	1.9	25.7
2	3	S	15	7.04	80	0	2.7	17.9	0.0	0.0	0.5	58.1	1.3	26.0	102.1
2	3	S	15	7.04	80	5	4.1	23.1	0.0	0.0	1.9	50.1	0.1	24.3	97.5
2	3	S	15	7.04	80	14	3.4	29.4	13.5	6.8	1.6	43.3	0.1	23.4	102.9
2	3	S	15	7.04	80	21	7.9	28.1	0.0	0.0	1.0	45.1	2.7	22.4	95.6
2	3	S	30	2.81	0	0	1.6	20.4	0.0	0.0	0.1	0.3	0.6	3.4	24.0
2	3	S	30	2.81	0	5	0.8	19.4	0.0	0.0	1.2	0.8	0.5	2.3	22.4
2	3	S	30	2.81	0	14	3.3	18.3	0.5	0.3	0.0	0.2	0.2	2.1	20.9
2	3	S	30	2.81	0	21	0.3	21.1	0.0	0.0	0.0	0.2	0.4	2.1	23.4
2	3	S	30	2.81	80	0	4.9	18.8	0.0	0.0	3.9	51.8	1.3	26.0	96.5
2	3	S	30	2.81	80	5	6.5	21.4	0.0	0.0	5.4	49.3	1.1	26.8	97.5
2	3	S	30	2.81	80	14	1.8	24.0	0.5	0.0	1.7	49.3	0.1	31.6	105.1
2	3	S	30	2.81	80	21	10.6	19.0	0.0	0.0	10.0	44.2	9.1	27.3	90.5
2	3	S	30	7.04	0	0	0.2	20.7	0.0	0.0	0.6	0.6	1.9	3.1	24.3
2	3	S	30	7.04	0	5	3.5	21.6	0.0	0.0	0.2	0.2	0.7	2.1	24.1
2	3	S	30	7.04	0	14	0.9	18.6	0.0	0.0	0.0	0.1	1.0	3.0	21.6
2	3	S	30	7.04	0	21	0.4	26.4	0.0	0.0	0.7	0.4	0.1	1.8	28.6
2	3	S	30	7.04	80	0	2.7	17.9	0.0	0.0	4.5	58.1	1.3	26.0	102.1
2	3	S	30	7.04	80	5	2.4	30.1	0.0	0.0	3.1	43.8	0.3	22.8	96.6
2	3	S	30	7.04	80	14	21.4	46.8	0.0	0.0	6.0	33.1	1.7	21.8	101.8
2	3	S	30	7.04	80	21	5.1	56.4	0.0	0.0	0.7	25.9	0.8	17.4	99.8
2	4	S	4	2.26	0	0	6.8	6.2	0.0	0.0	0.0	0.8	0.7	1.3	8.3
2	4	S	4	2.26	0	5	0.4	10.2	0.0	0.0	0.0	0.0	0.1	0.9	11.1
2	4	S	4	2.26	0	14	0.2	9.3	0.0	0.0	0.0	0.0	1.4	3.4	12.7
2	4	S	4	2.26	0	21	1.5	9.4	0.0	0.0	0.0	0.0	0.0	2.9	12.3
2	4	S	4	2.26	80	0	6.4	12.7	0.0	0.0	0.6	62.8	1.4	22.1	97.6
2	4	S	4	2.26	80	5	0.8	11.7	0.0	0.0	2.3	60.3	1.6	23.0	94.9
2	4	S	4	2.26	80	14	1.0	5.6	0.0	0.0	0.0	60.8	4.4	25.5	91.9
2	4	S	4	2.26	80	21	0.9	11.8	0.0	0.0	1.1	59.6	1.8	24.0	95.5
2	4	S	4	5.59	0	0	0.1	10.3	0.0	0.0	0.4	0.2	0.2	1.2	11.7
2	4	S	4	5.59	0	5	1.3	10.5	0.0	0.0	0.3	0.0	0.1	0.9	11.5
2	4	S	4	5.59	0	14	3.9	8.5	0.0	0.0	0.0	0.0	0.1	2.1	10.6
2	4	S	4	5.59	0	21	0.4	10.7	0.0	0.0	0.0	0.0	0.3	2.0	12.7
2	4	S	4	5.59	80	0	1.5	12.4	0.0	0.0	1.7	61.6	0.8	22.3	96.4
2	4	S	4	5.59	80	5	0.2	7.3	0.0	0.0	0.6	65.6	0.1	21.6	94.4
2	4	S	4	5.59	80	14	3.2	12.1	0.0	0.0	1.2	58.5	0.5	22.6	93.1
2	4	S	4	5.59	80	21	0.9	8.9	0.0	0.0	1.7	60.4	0.4	22.2	91.6
2	4	S	15	2.26	0	0	6.8	6.2	0.0	0.0	0.0	0.8	0.7	1.3	8.3
2	4	S	15	2.26	0	5	0.2	9.2	0.0	0.0	0.0	0.0	0.3	1.4	10.6
2	4	S	15	2.26	0	10	0.9	9.8	0.0	0.0	0.1	0.1	0.2	1.8	11.7
2	4	S	15	2.26	0	21	0.7	10.1	0.0	0.0	0.0	0.0	0.1	1.6	11.8
2	4	S	15	2.26	80	0	6.4	12.7	0.0	0.0	0.6	62.8	1.4	22.1	97.6
2	4	S	15	2.26	80	5	2.8	7.1	0.0	0.0	1.7	61.6	0.6	22.1	90.8
2	4	S	15	2.26	80	14	1.9	10.8	0.0	0.0	2.9	61.6	4.5	24.4	96.8
2	4	S	15	2.26	80	21	5.8	11.4	0.0	0.0	1.7	59.9	0.4	23.2	94.5
2	4	S	15	5.59	0	0	0.1	10.3	0.0	0.0	0.4	0.2	0.2	1.2	11.7
2	4	S	15	5.59	0	5	1.4	10.8	0.0	0.0	0.0	0.0	0.6	1.3	12.1
2	4	S	15	5.59	0	14	2.9	10.3	0.0	0.0	0.0	0.0	0.1	1.1	11.4

2	4	S	15	5.59	0	21	0.7	10.9	0.0	0.0	0.1	0.1	0.0	1.0	12.0
2	4	S	15	5.59	80	0	1.5	12.4	0.0	0.0	1.7	61.6	0.8	22.3	96.4
2	4	S	15	5.59	80	5	12.5	10.4	0.0	0.0	1.7	64.4	0.6	21.0	95.9
2	4	S	15	5.59	80	14	1.3	12.3	0.0	0.0	1.7	59.9	0.2	21.7	94.0
2	4	S	15	5.59	80	21	7.7	15.6	0.0	0.0	2.3	54.8		21.7	93.5

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		SOL NH4		EX NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
2	4	S	30	2.26	0	0	6.8	6.2	0.0	0.0	0.0	0.8	0.7	1.3	8.3
2	4	S	30	2.26	0	5	0.2	9.5	0.0	0.0	0.0	0.0	0.0	1.0	10.5
2	4	S	30	2.26	0	14	0.2	9.7	0.0	0.0	0.6	0.3	0.1	1.1	11.1
2	4	S	30	2.26	0	21	1.0	9.5	0.0	0.0	0.1	0.1	0.0	1.3	10.9
2	4	S	30	2.26	80	0	6.4	12.7	0.0	0.0	0.6	62.8	1.4	22.1	97.6
2	4	S	30	2.26	80	5	0.6	11.7	0.0	0.0	3.4	59.1	0.1	22.8	93.6
2	4	S	30	2.26	80	14	0.1	12.4	0.0	0.0	0.6	56.6	0.3	24.0	93.1
2	4	S	30	2.26	80	21	7.7	13.0	0.0	0.0	2.8	54.3	1.1	23.3	90.6
2	4	S	30	5.59	0	0	0.1	10.3	0.0	0.0	0.4	0.2	0.2	1.2	11.7
2	4	S	30	5.59	0	5	0.4	10.4	0.0	0.0	0.0	0.0	0.0	0.6	11.0
2	4	S	30	5.59	0	14	1.0	8.9	0.0	0.0	0.1	0.1	0.0	0.9	9.8
2	4	S	30	5.59	0	21	1.8	9.3	0.0	0.0	0.0	0.0	0.5	0.8	10.1
2	4	S	30	5.59	80	0	1.5	12.4	0.0	0.0	1.7	61.6	0.8	22.3	96.4
2	4	S	30	5.59	80	5	1.2	12.0	0.0	0.0	2.3	61.3	0.	21.8	95.1
2	4	S	30	5.59	80	14	10.5	20.6	0.0	0.0	7.0	52.3	1.7	20.0	93.0
2	4	S	30	5.59	80	21	1.4	19.3	0.0	0.0	3.2	51.7	1.8	22.3	93.3
2	5	S	4	1.44	0	0	0.1	2.3	0.0	0.0	0.0	1.0	0.6	1.0	4.3
2	5	S	4	1.44	0	5	0.0	2.6	0.0	0.0	0.2	0.4	0.1	1.8	4.8
2	5	S	4	1.44	0	14	1.1	1.3	0.0	0.0	0.0	0.2	0.0	0.7	2.1
2	5	S	4	1.44	0	21	0.1	2.6	0.0	0.0	0.1	0.3	0.1	0.8	3.6
2	5	S	4	1.44	80	0	0.3	6.9	0.0	0.0	2.9	67.1	1.2	16.5	90.6
2	5	S	4	1.44	80	5	5.9	3.3	0.0	0.0	2.3	66.8	0.1	16.9	87.1
2	5	S	4	1.44	80	14	4.4	6.6	0.0	0.0	1.1	62.3	2.2	18.6	87.5
2	5	S	4	1.44	80	21	1.3	9.3	0.0	0.0	2.6	63.8	0.6	17.0	90.0
2	5	S	4	3.61	0	0	0.0	2.5	0.0	0.0	0.2	0.2	0.1	0.8	3.5
2	5	S	4	3.61	0	5	0.1	2.6	0.0	0.0	0.1	0.8	0.5	1.4	4.8
2	5	S	4	3.61	0	14	0.3	2.8	0.0	0.0	0.1	0.1	2.0	1.7	4.6
2	5	S	4	3.61	0	21	0.1	2.6	0.0	0.0	0.4	0.3	0.1	0.8	3.7
2	5	S	4	3.61	80	0	7.4	5.6	0.0	0.0	5.0	66.4	0.5	16.4	88.4
2	5	S	4	3.61	80	5	0.4	8.1	0.0	0.0	2.8	64.7	0.6	17.1	89.9
2	5	S	4	3.61	80	14	0.6	5.5	0.0	0.0	1.2	61.6	0.1	17.1	84.1
2	5	S	4	3.61	80	21	3.2	7.9	0.0	0.0	1.1	62.8	0.1	17.1	87.8
2	5	S	15	1.44	0	0	0.1	2.3	0.0	0.0	0.0	1.0	0.6	1.0	4.3
2	5	S	15	1.44	0	5	0.2	2.2	0.0	0.0	0.1	0.4	0.4	1.5	4.1
2	5	S	15	1.44	0	14	0.1	2.3	0.0	0.0	0.3	0.3	0.3	0.8	3.4
2	5	S	15	1.44	0	22	0.4	2.6	0.0	0.0	0.0	0.5	0.0	0.8	3.9
2	5	S	15	1.44	80	0	0.3	6.9	0.0	0.0	2.9	67.1	1.2	16.5	90.6
2	5	S	15	1.44	80	5	10.0	7.5	0.0	0.0	3.4	65.2	0.3	17.8	90.4
2	5	S	15	1.44	80	14	1.9	3.1	0.0	0.0	1.1	65.8	0.3	16.9	85.8
2	5	S	15	1.44	80	21	3.5	8.9	0.0	0.0	1.1	64.1	0.9	18.1	91.1
2	5	S	15	3.61	0	0	0.0	2.5	0.0	0.0	0.2	0.2	0.1	0.8	3.5
2	5	S	15	3.61	0	5	0.1	2.3	0.0	0.0	0.2	0.2	1.3	1.8	4.4
2	5	S	15	3.61	0	14	1.8	0.9	0.0	0.0	0.0	0.0	0.4	1.0	1.9
2	5	S	15	3.61	0	21	0.2	2.6	0.0	0.0	0.1	0.1	0.2	0.7	3.3
2	5	S	15	3.61	80	0	7.4	5.6	0.0	0.0	5.0	66.4	0.5	16.4	88.4
2	5	S	15	3.61	80	5	2.0	7.8	0.0	0.0	0.0	62.7	0.7	17.3	87.7
2	5	S	15	3.61	80	14	5.5	2.9	0.0	0.0	4.0	61.3	0.8	17.8	82.0
2	5	S	15	3.61	80	21	1.5	4.3	0.0	0.0	4.5	65.6	0.3	17.3	87.1
2	5	S	30	1.44	0	0	0.1	2.3	0.0	0.0	0.0	1.0	0.6	1.0	4.3
2	5	S	30	1.44	0	5	0.2	2.7	0.0	0.0	0.0	0.3	0.2	0.9	3.9
2	5	S	30	1.44	0	14	0.2	1.9	0.0	0.0	0.0	0.0	2.5	2.4	4.3
2	5	S	30	1.44	0	21	0.4	2.6	0.0	0.0	0.0	0.5	0.0	0.9	4.0
2	5	S	30	1.44	80	0	0.3	6.9	0.0	0.0	2.9	67.1	1.2	16.5	90.6
2	5	S	30	1.44	80	5	1.8	4.7	0.0	0.0	2.2	65.2	0.4	17.9	87.8
2	5	S	30	1.44	80	14	3.6	2.8	0.0	0.0	0.0	57.3	0.4	18.2	78.3
2	5	S	30	1.44	80	21	0.5	5.1	0.0	0.0	0.0	57.3	0.0	19.7	82.0
2	5	S	30	3.61	0	0	0.0	2.5	0.0	0.0	0.2	0.2	0.1	0.8	3.5
2	5	S	30	3.61	0	5	0.5	2.3	0.0	0.0	0.0	0.0	0.0	0.6	2.8
2	5	S	30	3.61	0	14	2.2	1.4	0.0	0.0	0.3	0.1	0.2	0.7	2.2
2	5	S	30	3.61	0	21	0.3	1.5	0.0	0.0	0.3	0.3	0.2	0.5	2.3
2	5	S	30	3.61	80	0	7.4	5.6	0.0	0.0	5.0	66.4	0.5	16.4	88.4
2	5	S	30	3.61	80	5	1.4	2.9	0.0	0.0	1.7	66.3	0.4	17.8	87.0
2	5	S	30	3.61	80	14	7.0	4.9	0.0	0.0	3.4	61.0	3.1	20.3	86.2
2	5	S	30	3.61	80	21	5.7	6.1	0.0	0.0	4.6	59.9	2.1	19.8	85.8

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		SOL NH4		Ex NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
2	6	S	4	1.26	0	0	0.3	3.3	0.0	0.0	0.5	0.5	0.0	1.9	5.8
2	6	S	4	1.26	0	5	0.6	3.4	0.0	0.0	0.1	0.5	0.0	0.7	4.6
2	6	S	4	1.26	0	14	0.3	3.3	0.0	0.0	0.5	0.4	0.0	1.3	5.0
2	6	S	4	1.26	0	21	0.3	3.3	0.0	0.0	0.2	0.6	0.5	2.6	6.4
2	6	S	4	1.26	80	0	3.9	3.8	0.0	0.0	2.0	74.4	1.3	13.0	91.2
2	6	S	4	1.26	80	5	1.8	16.7	0.0	0.0	2.9	65.6	0.4	16.6	99.0
2	6	S	4	1.26	80	14	5.7	5.3	0.0	0.0	7.7	72.9	4.4	14.1	92.3
2	6	S	4	1.26	80	21	6.7	8.4	0.0	0.0	10.4	73.6	2.5	14.5	96.6
2	6	S	4	3.14	0	0	0.3	3.1	0.0	0.0	0.0	0.3	0.2	1.1	4.5
2	6	S	4	3.14	0	5	0.2	3.7	0.0	0.0	0.1	0.3	0.0	2.1	6.0
2	6	S	4	3.14	0	14	0.3	3.3	0.0	0.0	0.2	0.4	0.1	1.1	4.8
2	6	S	4	3.14	0	21	0.4	3.2	0.0	0.0	0.2	0.7	0.1	2.6	6.4
2	6	S	4	3.14	80	0	3.0	1.5	0.0	0.0	5.2	74.4	1.8	14.7	90.6
2	6	S	4	3.14	80	5	0.7	17.9	0.0	0.0	1.2	59.7	0.8	16.0	93.6
2	6	S	4	3.14	80	14	1.7	0.8	0.0	0.0	0.5	77.6	0.7	10.3	88.6
2	6	S	4	3.14	80	21	3.3	1.7	0.0	0.0	1.5	74.1	0.0	14.3	90.2
2	6	S	15	1.26	0	0	0.3	3.3	0.0	0.0	0.5	0.5	0.0	1.9	5.6
2	6	S	15	1.26	0	5	0.6	3.3	0.0	0.0	0.3	0.8	0.9	1.2	5.3
2	6	S	15	1.26	0	14	0.0	3.1	0.0	0.0	0.3	0.3	0.8	1.2	4.6
2	6	S	15	1.26	0	21	0.2	3.1	0.0	0.0	0.2	0.6	4.8	4.8	8.5
2	6	S	15	1.26	80	0	3.9	3.8	0.0	0.0	2.0	74.4	1.3	13.0	91.2
2	6	S	15	1.26	80	5	1.1	0.6	0.0	0.0	13.3	75.3	1.5	11.1	89.9
2	6	S	15	1.26	80	14	0.0	0.0	0.0	0.0	3.4	71.9	0.4	11.6	83.5
2	6	S	15	1.26	80	21	0.8	2.1	0.0	0.0	2.5	74.4	0.2	14.6	91.0
2	6	S	15	3.14	0	0	0.3	3.1	0.0	0.0	0.0	0.3	0.2	1.1	4.5
2	6	S	15	3.14	0	5	0.1	3.3	0.0	0.0	0.1	0.1	0.3	1.3	4.6
2	6	S	15	3.14	0	14	0.1	3.6	0.0	0.0	0.1	0.3	0.0	0.6	4.5
2	6	S	15	3.14	0	21	0.7	3.8	0.0	0.0	0.2	0.7	0.3	2.0	6.5
2	6	S	15	3.14	80	0	3.0	1.5	0.0	0.0	5.2	74.4	1.8	14.7	90.6
2	6	S	15	3.14	80	5	2.0	1.0	0.0	0.0	0.3	75.0	0.4	11.9	88.0
2	6	S	15	3.14	80	14	0.1	5.4	0.0	0.0	1.8	75.2	0.2	11.5	92.1
2	6	S	15	3.14	80	21	1.2	2.6	0.0	0.0	1.4	75.6	5.0	10.4	86.6
2	6	S	30	1.26	0	0	0.3	3.3	0.0	0.0	0.5	0.5	0.4	1.9	5.6
2	6	S	30	1.26	0	5	0.2	3.1	0.0	0.0	0.4	0.6	0.1	0.8	4.5
2	6	S	30	1.26	0	14	0.1	3.3	0.0	0.0	0.2	0.4	0.0	0.5	4.2
2	6	S	30	1.26	0	21	0.7	3.5	0.0	0.0	0.4	0.8	0.2	1.9	6.2
2	6	S	30	1.26	80	0	3.9	3.8	0.0	0.0	2.0	74.4	1.3	13.0	91.2
2	6	S	30	1.26	80	5	4.9	2.4	0.0	0.0	2.9	73.1	1.3	10.5	86.0
2	6	S	30	1.26	80	14	6.0	3.9	0.0	0.0	1.4	73.5	0.2	10.7	88.1
2	6	S	30	1.26	80	21	3.5	3.3	0.0	0.0	8.4	70.0	1.1	16.3	89.5
2	6	S	30	3.14	0	0	0.3	3.1	0.0	0.0	0.0	0.3	0.2	1.1	4.5
2	6	S	30	3.14	0	5	0.2	3.1	0.0	0.0	0.6	0.5	0.3	0.8	4.3
2	6	S	30	3.14	0	14	0.5	3.1	0.0	0.0	0.1	0.3	0.0	0.5	3.9
2	6	S	30	3.14	0	21	0.0	3.9	0.0	0.0	0.0	0.6	2.9	3.1	7.5
2	6	S	30	3.14	80	0	3.0	1.5	0.0	0.0	5.2	74.4	1.8	14.7	90.6
2	6	S	30	3.14	80	5	0.8	0.4	0.0	0.0	2.6	77.4	1.7	10.4	88.2
2	6	S	30	3.14	80	14	1.1	2.9	0.0	0.0	4.9	69.3	3.7	13.3	85.4
2	6	S	30	3.14	80	21	0.0	0.0	0.0	0.0	2.1	72.1	1.6	15.8	87.9
2	7	S	4	1.22	0	0	0.4	4.5	0.0	0.0	0.4	0.4	0.1	1.1	6.0
2	7	S	4	1.22	0	5	0.1	4.1	0.0	0.0	0.4	0.3	0.0	0.8	5.1
2	7	S	4	1.22	0	14	0.1	4.4	0.0	0.0	0.3	0.5	0.0	0.8	6.0
2	7	S	4	1.22	0	21	0.3	4.8	0.0	0.0	0.1	0.8	0.6	1.7	7.3
2	7	S	4	1.22	80	0	9.0	8.0	0.0	0.0	0.6	77.2	0.5	12.3	97.4
2	7	S	4	1.22	80	5	3.6	2.6	0.0	0.0	5.4	79.3	1.0	10.6	92.5
2	7	S	4	1.22	80	14	0.0	0.0	0.0	0.0	0.3	76.8	0.4	6.8	83.5
2	7	S	4	1.22	80	21	1.8	5.6	0.0	0.0	1.7	80.0	0.7	10.6	96.3
2	7	S	4	3.05	0	0	0.0	4.4	0.0	0.0	0.0	0.3	0.1	1.1	5.8
2	7	S	4	3.05	0	5	0.3	4.8	0.0	0.0	0.1	0.8	0.2	0.9	8.5
2	7	S	4	3.05	0	14	0.0	4.7	0.0	0.0	0.1	0.3	0.0	0.9	5.9
2	7	S	4	3.05	0	21	0.1	4.6	0.0	0.0	0.1	1.6	2.2	2.5	8.7
2	7	S	4	3.05	80	0	8.0	5.8	0.0	0.0	2.3	69.4	0.1	13.3	88.6
2	7	S	4	3.05	80	5	0.0	0.0	0.0	0.0	0.3	82.6	0.8	11.1	93.9
2	7	S	4	3.05	80	14	2.4	11.6	0.0	0.0	0.6	77.2	1.3	6.5	95.3
2	7	S	4	3.05	80	21	6.9	8.8	0.0	0.0	3.4	79.5	1.5	13.6	102.0
2	7	S	15	1.22	0	0	0.4	4.5	0.0	0.0	0.4	0.4	0.1	1.1	6.0
2	7	S	15	1.22	0	5	0.6	4.4	0.0	0.0	0.2	1.0	0.3	0.8	6.2
2	7	S	15	1.22	0	14	0.3	4.6	0.0	0.0	0.1	0.5	1.0	1.3	8.5
2	7	S	15	1.22	0	21	0.0	4.9	0.0	0.0	0.7	1.6	0.1	1.3	7.8
2	7	S	15	1.22	80	0	9.0	8.0	0.0	0.0	0.6	77.2	0.5	12.3	97.4
2	7	S	15	1.22	80	5	6.3	3.1	0.0	0.0	3.7	78.4	1.0	11.5	93.1
2	7	S	15	1.22	80	14	0.6	4.8	0.0	0.0	2.3	75.8	0.1	7.0	87.6
2	7	S	15	1.22	80	21	9.8	7.8	0.0	0.0	2.3	76.6	0.5	10.9	95.6
2	7	S	15	3.05	0	0	0.0	4.4	0.0	0.0	0.0	0.3	0.1	1.1	5.8
2	7	S	15	3.05	0	5	0.4	4.6	0.0	0.0	0.2	0.5	0.2	1.1	6.2
2	7	S	15	3.05	0	14	0.7	4.4	0.0	0.0	0.3	0.3	0.1	0.8	5.4
2	7	S	15	3.05	0	21	0.2	4.6	0.0	0.0	0.0	1.9	0.3	1.1	7.6

2	7	S	15	3.05	80	0	8.0	5.8	0.0	0.0	2.3	69.4	0.1	13.3	88.6
2	7	S	15	3.05	80	5	2.9	5.3	0.0	0.0	4.6	80.7	0.4	11.0	97.0
2	7	S	15	3.05	80	14	0.5	9.4	0.0	0.0	2.3	73.8	1.3	7.9	91.1
2	7	S	15	3.05	80	21	1.6	5.0	0.0	0.0	2.5	63.9			96.2

Table 3 cont'd.- 22 -

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		SOL	NH4	Ex	NH4	TOTAL
							RANGE	AVER	RANGE	AVER					
2	7	S	30	1.22	0	0	0.4	4.5	0.0	0.0	0.4	0.4	0.4	1.1	6.0
2	7	S	30	1.22	0	5	0.5	4.6	0.0	0.0	0.3	0.9	0.5	1.4	6.9
2	7	S	30	1.22	0	14	0.3	4.3	0.0	0.0	0.1	0.9	0.1	0.8	6.0
2	7	S	30	1.22	0	21	2.3	5.5	0.0	0.0	0.1	2.4	0.9	0.7	6.6
2	7	S	30	1.22	80	0	9.0	8.0	0.0	0.0	0.6	77.2	0.5	12.3	97.4
2	7	S	30	1.22	80	5	3.4	2.8	0.0	0.0	4.0	80.0	1.3	11.8	94.6
2	7	S	30	1.22	80	14	8.2	10.7	0.0	0.0	1.7	72.5	2.1	8.6	91.9
2	7	S	30	1.22	80	21	1.1	6.3	0.0	0.0	1.7	74.3	2.2	12.5	93.0
2	7	S	30	3.05	0	0	0.0	4.4	0.0	0.0	0.0	0.3	0.1	1.1	5.8
2	7	S	30	3.05	0	5	0.5	4.4	0.0	0.0	0.0	0.2	0.0	1.4	5.9
2	7	S	30	3.05	0	14	0.2	4.8	0.0	0.0	0.0	1.5	0.0	1.2	7.5
2	7	S	30	3.05	0	21	1.6	4.9	0.0	0.0	0.0	1.6	2.0	2.0	8.5
2	7	S	30	3.05	80	0	8.0	5.8	0.0	0.0	2.3	69.4	0.1	13.3	88.6
2	7	S	30	3.05	80	5	0.0	0.0	0.0	0.0	1.1	76.3	0.4	12.5	88.8
2	7	S	30	3.05	80	14	8.4	7.1	0.0	0.0	3.8	75.6	4.6	11.0	93.7
2	7	S	30	3.05	80	21	0.0	6.9	0.0	0.0	0.0	80.1	1.0	11.4	98.4
3	1	S	4	2.87	0	0	0.7	14.8	3.5	1.8	0.1	0.1	0.0	1.7	18.2
3	1	S	4	2.87	0	5	2.2	19.4	0.0	0.0	0.7	0.8	0.2	1.5	21.7
3	1	S	4	2.87	0	14	2.1	18.3	0.0	0.0	0.0	0.0	0.2	1.8	18.0
3	1	S	4	2.87	0	21	2.5	18.6	0.0	0.0	0.1	0.6	0.3	1.8	21.0
3	1	S	4	2.87	80	0	2.2	20.2	0.0	0.0	3.3	52.8	0.5	24.4	97.5
3	1	S	4	2.87	80	5	3.4	19.0	0.0	0.0	10.2	56.5	3.3	23.6	99.1
3	1	S	4	2.87	80	14	1.9	21.8	0.0	0.0	5.9	58.6	0.1	22.8	103.2
3	1	S	4	2.87	80	21	3.7	13.8	0.0	0.0	3.6	75.5	0.9	24.8	114.1
3	1	S	4	7.17	0	0	2.0	19.9	0.0	0.0	0.9	0.4	3.4	1.8	22.1
3	1	S	4	7.17	0	5	1.8	19.8	0.0	0.0	0.7	0.4	0.6	2.2	22.4
3	1	S	4	7.17	0	14	3.1	19.8	0.0	0.0	0.3	0.3	0.2	2.1	22.2
3	1	S	4	7.17	0	21	2.7	19.9	0.0	0.0	0.2	1.2	0.5	2.4	23.5
3	1	S	4	7.17	80	0	3.2	18.6	0.0	0.0	1.2	56.8	0.7	23.6	99.0
3	1	S	4	7.17	80	5	1.4	21.6	0.0	0.0	1.2	56.8	0.5	22.1	100.5
3	1	S	4	7.17	80	14	1.2	20.7	0.0	0.0	2.3	55.1	0.1	23.1	99.0
3	1	S	4	7.17	80	21	1.7	14.1	0.0	0.0	1.2	62.3	0.5	22.1	98.5
3	1	S	15	2.87	0	0	0.7	14.6	3.5	1.8	0.1	0.1	0.0	1.7	18.2
3	1	S	15	2.87	0	5	3.7	15.4	0.0	0.0	0.2	0.3	0.0	2.2	17.9
3	1	S	15	2.87	0	14	2.6	16.2	0.0	0.0	0.0	0.0	0.6	2.4	18.6
3	1	S	15	2.87	0	21	0.1	17.3	0.0	0.0	0.2	0.6	0.5	1.8	19.6
3	1	S	15	2.87	80	0	2.2	20.2	0.0	0.0	3.3	52.8	0.5	24.4	97.5
3	1	S	15	2.87	80	5	2.7	9.5	0.0	0.0	4.8	62.8	1.2	24.1	96.4
3	1	S	15	2.87	80	14	1.4	13.8	0.0	0.0	2.5	54.4	3.1	22.1	90.3
3	1	S	15	2.87	80	21	0.2	17.8	0.0	0.0	7.8	60.7	0.8	21.7	100.2
3	1	S	15	7.17	0	0	2.0	19.9	0.0	0.0	0.9	0.4	3.4	1.8	22.1
3	1	S	15	7.17	0	5	0.3	18.9	0.0	0.0	0.0	0.0	0.9	1.9	20.9
3	1	S	15	7.17	0	14	2.5	20.4	0.0	0.0	0.7	0.3	0.1	0.8	21.4
3	1	S	15	7.17	0	21	0.2	19.3	0.0	0.0	0.1	0.5	0.1	1.3	21.2
3	1	S	15	7.17	80	0	3.2	18.6	0.0	0.0	1.2	56.8	0.7	23.6	99.0
3	1	S	15	7.17	80	5	4.9	14.7	2.0	1.0	4.1	62.1	1.7	20.3	98.1
3	1	S	15	7.17	80	14	1.6	34.4	0.0	0.0	5.2	49.3	1.5	20.4	104.0
3	1	S	15	7.17	80	21	8.0	39.4	0.0	0.0	0.8	44.3	2.0	17.8	101.5
3	1	S	30	2.87	0	0	0.7	14.8	3.5	1.8	0.1	0.1	0.0	1.7	18.2
3	1	S	30	2.87	0	5	0.2	16.3	0.0	0.0	0.2	0.1	0.4	1.8	18.2
3	1	S	30	2.87	0	14	0.3	17.0	0.0	0.0	0.5	0.4	0.4	3.5	21.0
3	1	S	30	2.87	0	21	1.5	17.4	0.0	0.0	0.1	0.6	0.0	1.8	19.8
3	1	S	30	2.87	80	0	2.2	20.2	0.0	0.0	3.3	52.8	0.5	24.4	97.5
3	1	S	30	2.87	80	5	5.3	19.1	0.0	0.0	5.0	50.6	3.1	22.6	92.3
3	1	S	30	2.87	80	14	4.4	16.9	0.0	0.0	7.9	57.1	0.2	26.0	99.9
3	1	S	30	2.87	80	21	6.0	14.6	0.0	0.0	4.8	60.4	1.0	21.8	96.8
3	1	S	30	7.17	0	0	2.0	19.9	0.0	0.0	0.9	0.4	3.4	1.8	22.1
3	1	S	30	7.17	0	5	0.1	20.8	0.0	0.0	0.3	0.3	0.3	1.3	22.3
3	1	S	30	7.17	0	14	1.0	21.5	0.0	0.0	0.2	1.2	0.4	2.8	25.5
3	1	S	30	7.17	0	21	0.2	23.2	0.0	0.0	0.1	0.3	0.6	1.2	24.6
3	1	S	30	7.17	80	0	3.2	18.6	0.0	0.0	1.2	56.8	0.7	23.6	99.0
3	1	S	30	7.17	80	5	5.4	38.7	11.5	5.8	7.5	43.9	1.8	18.1	106.4
3	1	S	30	7.17	80	14	25.1	76.1	27.0	13.5	10.6	12.7	5.2	7.0	109.3
3	1	S	30	7.17	80	21	30.2	64.4	0.0	0.0	20.2	22.5	7.7	10.5	97.4

Table 3 cont'd.

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		SOL NH4		EX NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
3	2	S	4	3.06	0	0	3.4	13.9	0.0	0.0	1.0	0.6	0.1	2.1	16.6
3	2	S	4	3.06	0	5	0.6	12.1	0.0	0.0	0.4	0.4	0.2	2.1	14.6
3	2	S	4	3.06	0	14	2.6	11.6	0.0	0.0	0.0	0.1	0.0	1.6	13.3
3	2	S	4	3.06	0	21	1.4	14.1	0.0	0.0	0.3	0.3	0.1	2.1	16.6
3	2	S	4	3.06	80	0	4.4	23.7	0.0	0.0	6.5	57.3	0.6	23.5	104.4
3	2	S	4	3.06	80	5	3.3	21.0	0.0	0.0	4.0	55.5	0.0	22.7	99.2
3	2	S	4	3.06	80	14	4.3	19.5	0.0	0.0	1.5	53.8	0.1	23.3	96.6
3	2	S	4	3.06	80	21	0.1	26.8	0.0	0.0	4.1	54.9	1.0	23.7	105.4
3	2	S	4	7.65	0	0	2.3	14.9	0.0	0.0	0.2	0.9	0.4	2.3	18.1
3	2	S	4	7.65	0	5	1.4	13.7	0.0	0.0	0.2	0.4	0.1	1.8	15.8
3	2	S	4	7.65	0	14	0.9	14.9	0.0	0.0	0.1	0.1	0.2	2.3	17.4
3	2	S	4	7.65	0	21	0.6	17.6	0.0	0.0	0.3	0.3	0.9	1.8	19.7
3	2	S	4	7.65	80	0	2.2	23.6	0.0	0.0	1.5	58.9	0.2	24.7	107.1
3	2	S	4	7.65	80	5	9.9	21.3	0.0	0.0	0.5	56.4	0.6	22.7	100.3
3	2	S	4	7.65	80	14	0.1	27.1	0.5	0.8	1.0	50.6	0.6	22.7	105.1
3	2	S	4	7.65	80	21	3.2	29.5	1.5	0.8	1.5	55.4	0.0	23.4	109.0
3	2	S	15	3.06	0	0	3.4	13.9	0.0	0.0	1.0	0.6	0.1	2.1	16.3
3	2	S	15	3.06	0	5	0.8	12.9	0.0	0.0	1.3	1.5	0.6	1.9	16.3
3	2	S	15	3.06	0	14	2.1	13.9	0.0	0.0	1.2	1.0	0.4	1.8	16.7
3	2	S	15	3.06	0	21	1.3	15.0	0.0	0.0	1.6	0.9	0.1	1.8	17.7
3	2	S	15	3.06	80	0	4.4	23.7	0.0	0.0	6.5	57.3	0.6	23.5	104.4
3	2	S	15	3.06	80	5	11.3	25.9	0.0	0.0	5.5	56.8	0.0	24.2	106.9
3	2	S	15	3.06	80	14	6.2	28.8	0.0	0.0	7.5	57.3	2.6	25.5	111.5
3	2	S	15	3.06	80	21	2.0	23.9	0.0	0.0	1.5	53.8	1.6	23.9	101.5
3	2	S	15	7.65	0	0	2.3	14.9	0.0	0.0	0.2	0.9	0.4	2.3	18.1
3	2	S	15	7.65	0	5	2.3	14.9	0.0	0.0	0.8	0.5	0.4	2.2	17.6
3	2	S	15	7.65	0	14	1.9	21.1	0.0	0.0	0.0	0.1	0.3	0.9	22.2
3	2	S	15	7.65	0	21	3.2	21.8	0.0	0.0	0.0	0.1	0.0	0.7	22.4
3	2	S	15	7.65	80	0	2.2	23.6	0.0	0.0	1.5	58.9	0.2	24.7	107.1
3	2	S	15	7.65	60	5	4.4	28.6	0.0	0.0	1.5	55.9	0.5	22.6	107.1
3	2	S	15	7.65	80	14	3.6	54.9	0.0	0.0	3.9	33.6	4.8	16.4	104.9
3	2	S	15	7.65	80	21	18.4	71.3	0.0	0.0	19.0	24.5	7.6	12.3	108.1
3	2	S	30	3.06	0	0	3.4	13.9	0.0	0.0	1.0	0.6	0.1	2.1	16.6
3	2	S	30	3.06	0	5	0.6	13.2	0.0	0.0	0.0	0.1	0.1	1.6	14.9
3	2	S	30	3.06	0	14	2.2	16.4	0.0	0.0	0.4	0.7	1.8	2.4	19.5
3	2	S	30	3.06	0	21	0.9	15.9	0.0	0.0	0.1	0.6	0.6	2.1	18.7
3	2	S	30	3.06	80	0	4.4	23.7	0.0	0.0	6.5	57.3	0.6	23.5	104.4
3	2	S	30	3.06	80	5	2.6	19.9	0.0	0.0	1.0	49.5	1.3	23.5	92.9
3	2	S	30	3.06	80	14	3.6	19.4	0.0	0.0	4.5	57.3	1.4	24.4	101.0
3	2	S	30	3.06	80	21	5.6	20.7	0.0	0.0	3.0	52.0	4.6	22.7	95.4
3	2	S	30	7.65	0	0	2.3	14.9	0.0	0.0	0.2	0.9	0.4	2.3	18.1
3	2	S	30	7.65	0	5	0.0	20.8	0.0	0.0	0.0	0.8	0.0	1.5	23.1
3	2	S	30	7.65	0	14	0.1	22.6	0.0	0.0	0.3	0.3	0.3	1.0	23.8
3	2	S	30	7.65	0	21	2.4	26.7	0.0	0.0	0.0	0.1	0.0	0.8	27.6
3	2	S	30	7.65	80	0	2.2	23.6	0.0	0.0	1.5	58.9	0.2	24.7	107.1
3	2	S	30	7.65	80	5	3.8	49.1	16.0	12.5	6.0	36.4	1.0	15.7	113.7
3	2	S	30	7.65	80	14	25.2	70.2	0.0	0.0	18.5	24.3	8.3	11.3	105.8
3	2	S	30	7.65	80	21	19.2	72.2	0.0	0.0	0.8	21.6	0.8	11.9	105.7
4	1	C	4	9.60	0	0	1.8	17.0	0.0	0.0	0.0	0.5	2.4	5.3	22.8
4	1	C	4	9.60	0	5	0.2	17.3	0.0	0.0	0.4	0.4	0.3	9.0	26.7
4	1	C	4	9.60	0	14	2.1	17.4	0.0	0.0	0.0	0.1	0.4	7.1	24.6
4	1	C	4	9.60	0	21	0.0	16.5	0.0	0.0	0.0	0.2	0.1	7.1	23.7
4	1	C	4	9.60	80	0	1.2	22.1	0.0	0.0	0.5	23.9	0.5	47.8	93.5
4	1	C	4	9.60	60	5	1.0	20.3	0.0	0.0	3.6	29.1	6.1	53.1	102.5
4	1	C	4	9.60	80	14	0.4	21.0	0.0	0.0	0.4	25.3	0.9	42.6	88.9
4	1	C	4	9.60	80	21	1.4	17.0	0.0	0.0	2.8	23.3	5.0	39.8	80.1
4	1	C	4	24.00	0	0	1.5	19.4	0.0	0.0	0.0	0.2	0.0	8.5	28.1
4	1	C	4	24.00	0	5	0.8	27.6	0.0	0.0	1.0	0.9	0.6	6.7	35.2
4	1	C	4	24.00	0	14	0.8	35.9	0.0	0.0	1.0	0.7	0.7	2.1	38.7
4	1	C	4	24.00	0	21	1.1	43.8	0.0	0.0	0.0	0.1	0.2	1.1	45.0
4	1	C	4	24.00	80	0	0.7	24.6	0.0	0.0	2.1	30.4	1.9	55.3	110.4
4	1	C	4	24.00	60	5	3.3	34.8	0.0	0.0	6.5	25.4	7.4	46.0	108.2
4	1	C	4	24.00	80	14	2.8	45.3	0.0	0.0	6.0	23.3	6.0	39.9	108.5
4	1	C	4	24.00	80	21	5.4	54.8	0.0	0.0	14.5	25.3	14.8	41.3	121.3

PL	DP	SOL	TMP	WC	TRT	DAY	NOS		NO2		SOL NH4		EX NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
4	1	C	15	9.60	0	0	1.8	17.0	0.0	0.0	0.6	0.5	2.4	5.3	22.8
4	1	C	15	9.60	0	5	1.4	15.9	0.0	0.0	0.2	0.3	1.7	8.5	24.7
4	1	C	15	9.60	0	14	0.9	18.8	0.0	0.0	0.0	0.3	5.8	4.4	23.5
4	1	C	15	9.60	0	21	1.0	18.0	0.0	0.0	0.0	0.2	0.9	3.4	21.8
4	1	C	15	9.60	80	0	1.2	22.1	0.0	0.0	0.5	23.9	0.5	47.6	93.5
4	1	C	15	9.60	80	5	1.2	21.0	0.0	0.0	18.1	32.8	14.6	47.2	100.9
4	1	C	15	9.60	80	14	2.6	20.9	0.0	0.0	2.8	26.2	2.2	39.2	86.3
4	1	C	15	9.60	80	21	1.3	20.8	0.0	0.0	6.5	21.4	7.0	38.2	80.4
4	1	C	15	24.00	0	0	1.5	19.4	0.0	0.0	0.0	0.2	0.0	8.5	28.1
4	1	C	15	24.00	0	5	4.7	36.3	0.0	0.0	0.5	0.8	0.4	1.5	38.5
4	1	C	15	24.00	0	14	0.5	40.6	0.0	0.0	0.0	0.2	0.2	0.8	41.5
4	1	C	15	24.00	0	21	8.8	44.8	0.0	0.0	0.1	0.3	0.4	0.3	45.3
4	1	C	15	24.00	80	0	0.7	24.6	0.0	0.0	2.1	30.4	1.9	55.3	110.4
4	1	C	15	24.00	80	5	0.5	51.1	0.0	0.0	2.0	22.7	0.8	38.9	112.6
4	1	C	15	24.00	80	14	8.8	83.7	0.0	0.0	5.5	8.6	6.1	19.6	111.9
4	1	C	15	24.00	80	21	4.0	106.0	0.0	0.0	2.4	2.3	10.2	10.9	119.2
4	1	C	30	9.60	0	0	1.8	17.0	0.0	0.0	0.6	0.5	2.4	5.3	22.8
4	1	C	30	9.60	0	5	1.2	18.6	0.0	0.0	0.0	0.2	0.1	5.1	23.9
4	1	C	30	9.60	0	14	0.7	15.8	0.0	0.0	0.0	0.2	0.1	2.9	18.9
4	1	C	30	9.60	0	21	1.6	15.7	0.0	0.0	1.4	1.0	0.6	2.4	19.1
4	1	C	30	9.60	80	0	1.2	22.1	0.0	0.0	0.5	23.9	0.5	47.6	93.5
4	1	C	30	9.60	80	5	0.1	21.4	0.0	0.0	11.9	28.1	12.8	41.3	90.9
4	1	C	30	9.60	80	14	2.0	21.8	0.0	0.0	5.1	19.9	3.4	40.3	82.1
4	1	C	30	9.60	80	21	2.5	20.1	0.0	0.0	2.0	19.0	3.7	37.9	77.0
4	1	C	30	24.00	0	0	1.5	19.4	0.0	0.0	0.0	0.2	0.0	8.5	28.1
4	1	C	30	24.00	0	5	1.7	45.9	0.0	0.0	0.1	0.3	0.1	0.8	47.0
4	1	C	30	24.00	0	14	2.4	50.5	0.0	0.0	0.1	0.1	0.1	0.3	51.0
4	1	C	30	24.00	0	21	8.1	57.8	0.0	0.0	0.1	0.8	0.1	0.6	59.4
4	1	C	30	24.00	80	0	0.7	24.6	0.0	0.0	2.1	30.4	1.9	55.3	110.4
4	1	C	30	24.00	80	5	1.0	101.5	0.0	0.0	3.5	3.9	6.8	16.0	121.3
4	1	C	30	24.00	80	14	13.0	132.5	0.0	0.0	0.0	0.1	3.2	2.2	134.8
4	1	C	30	24.00	80	21	3.0	131.5	0.0	0.0	0.1	0.8	0.2	0.8	133.0
4	2	C	4	9.66	0	0	0.6	4.2	0.0	0.0	0.0	0.1	0.7	4.1	8.4
4	2	C	4	9.66	0	5	1.5	4.4	0.0	0.0	0.8	0.6	0.3	4.9	10.0
4	2	C	4	9.66	n	14	0.9	4.6	0.0	0.0	0.4	0.3	0.3	5.8	10.7
4	2	C	4	9.66	0	21	0.1	4.3	0.0	0.0	0.2	0.5	0.5	5.1	10.0
4	2	C	4	9.66	80	0	1.7	5.8	0.0	0.0	6.1	20.8	0.4	55.1	87.6
4	2	C	4	9.66	80	5	1.6	5.0	0.0	0.0	3.1	24.6	1.3	54.3	84.0
4	2	C	4	9.66	80	14	1.7	5.3	0.0	0.0	6.5	25.9	9.7	51.3	82.4
4	2	C	4	9.66	80	21	3.1	4.1	0.0	0.0	7.4	24.5	5.5	51.4	80.0
4	2	C	4	24.20	0	0	0.3	6.0	0.0	0.0	0.3	0.3	1.4	5.9	12.3
4	2	C	4	24.20	0	5	5.9	6.9	0.0	0.0	0.1	0.1	0.4	8.3	15.3
4	2	C	4	24.20	0	14	0.3	19.1	0.0	0.0	0.1	0.1	0.4	6.1	25.2
4	2	C	4	24.20	0	21	0.3	21.8	0.0	0.0	0.0	0.1	0.5	3.6	25.6
4	2	C	4	24.20	80	0	0.9	7.3	0.0	0.0	0.0	23.5	2.2	48.3	79.0
4	2	C	4	24.20	80	5	2.7	14.9	0.0	0.0	2.8	26.4	0.2	57.9	99.3
4	2	C	4	24.20	80	14	2.0	20.6	0.0	0.0	0.8	25.0	0.2	56.2	101.8
4	2	C	4	24.20	80	21	0.9	26.8	0.0	0.0	0.0	22.6	2.4	50.1	99.5
4	2	C	15	9.66	0	0	0.6	4.2	0.0	0.0	0.0	0.1	0.7	4.1	8.4
4	2	C	15	9.66	0	5	0.8	4.2	0.0	0.0	0.2	0.7	0.3	4.9	9.8
4	2	C	15	9.66	0	14	0.2	4.2	0.0	0.0	0.2	1.7	0.3	4.3	10.1
4	2	C	15	9.66	0	21	0.7	4.3	0.0	0.0	0.1	1.3	0.4	2.5	8.1
4	2	C	15	9.66	80	0	1.7	5.8	0.0	0.0	6.1	26.8	0.4	55.1	87.6
4	2	C	15	9.66	80	5	2.1	6.3	0.0	0.0	9.7	20.4	14.2	47.7	74.4
4	2	C	15	9.66	80	14	0.7	4.3	0.0	0.0	4.8	19.7	11.6	41.5	65.4
4	2	C	15	9.66	80	21	1.1	7.1	0.0	0.0	13.8	30.0	15.1	53.4	90.6
4	2	C	15	24.20	0	0	0.3	6.0	0.0	0.0	0.3	0.3	1.4	5.9	12.3
4	2	C	15	24.20	0	5	1.1	19.8	0.0	0.0	0.0	0.1	0.2	3.6	23.5
4	2	C	15	24.20	0	14	5.5	28.1	0.0	0.0	0.1	0.1	0.2	0.9	29.0
4	2	C	15	24.20	0	21	0.0	30.8	0.0	0.0	0.0	0.1	0.1	0.6	31.5
4	2	C	15	24.20	80	0	0.9	7.3	0.0	0.0	0.0	23.5	2.6	48.5	79.3
4	2	C	15	24.20	80	5	5.5	25.9	0.0	0.0	0.2	20.4	0.9	48.8	95.1
4	2	C	15	24.20	80	14	2.0	57.6	0.0	0.0	2.1	13.4	2.8	29.2	100.3
4	2	C	15	24.20	80	21	1.5	87.4	0.0	0.0	0.4	3.6	1.4	11.9	102.9
4	2	C	30	9.66	0	0	0.6	4.2	0.0	0.0	0.0	0.1	0.7	4.1	8.4
4	2	C	30	9.66	0	5	0.6	4.4	0.0	0.0	0.2	0.4	0.0	2.5	7.3
4	2	C	30	9.66	0	14	1.1	2.4	0.0	0.0	0.3	1.5	0.1	2.4	6.4
4	2	C	30	9.66	0	21	0.4	1.9	0.0	0.0	1.1	2.6	0.3	1.0	5.5
4	2	C	30	9.66	80	0	1.7	5.8	0.0	0.0	6.1	26.8	0.4	55.1	87.6
4	2	C	30	9.66	80	5	1.9	4.6	0.0	0.0	0.9	18.8	6.1	38.8	62.3
4	2	C	30	9.66	80	14	1.7	5.8	0.0	0.0	0.9	18.9	0.1	38.3	63.1
4	2	C	30	9.66	80	21	1.5	6.9	0.0	0.0	1.6	18.6	9.1	37.3	62.8
4	2	C	30	24.20	0	0	0.3	6.0	0.0	0.0	0.3	0.3	1.4	5.9	12.3
4	2	C	30	24.20	0	5	1.7	30.4	0.0	0.0	0.5	0.4	0.2	0.6	31.4
4	2	C	30	24.20	0	14	0.5	42.8	0.0	0.0	0.0	0.1	0.1	0.8	43.7
4	2	C	30	24.20	0	21	4.5	49.6	0.0	0.0	0.0	0.1	0.1	1.8	51.4
4	2	C	30	24.20	80	0	0.9	7.3	0.0	0.0	0.0	23.5	2.5	46.5	79.3

4	2	C	30	24.20	80	5	2.3	58.8	0.0	0.0	2.3	14.8	6.1	31.1	104.6
4	2	C	30	24.20	80	14	9.9	103.1	0.0	0.0	0.0	0.0	1.1	1.6	104.7
4	2	C	30	24.20	80	21	7.0	113.5	0.0	0.0	0.0	0.0	0.1	1.3	114.8

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		SOL NH4		EX NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
5	1	C	4	8.76	0	0	1.3	20.3	0.0	0.0	0.0	0.0	0.3	5.3	25.7
5	1	C	4	8.76	0	5	3.3	22.3	0.0	0.0	0.0	0.1	0.0	5.9	26.2
5	1	C	4	8.76	0	14	2.2	21.3	0.0	0.0	0.1	0.1	0.1	6.8	28.2
5	1	C	4	8.76	0	21	2.1	22.8	0.0	0.0	0.1	0.1	1.0	7.0	30.0
5	1	C	4	8.76	80	0	5.2	23.4	0.0	0.0	4.9	25.4	5.0	47.8	96.6
5	1	C	4	8.76	80	5	7.7	26.4	0.0	0.0	9.9	22.0	14.3	42.6	91.8
5	1	C	4	8.76	80	14	3.1	25.3	0.0	0.0	1.2	29.7	1.8	50.9	105.9
5	1	C	4	8.76	80	21	2.3	26.8	0.0	0.0	0.0	23.0	0.9	42.8	92.6
5	1	C	4	21.90	0	0	2.8	23.0	0.0	0.0	0.1	0.1	0.4	5.4	28.5
5	1	C	4	21.90	0	5	3.8	26.2	0.0	0.0	0.4	0.3	0.3	6.6	33.1
5	1	C	4	21.90	0	14	8.9	32.6	0.0	0.0	0.0	0.1	4.7	5.5	36.3
5	1	C	4	21.90	0	21	0.9	36.3	0.0	0.0	0.1	0.1	2.0	6.8	43.2
5	1	C	4	21.90	60	0	4.6	28.2	0.0	0.5	7.7	27.5	9.1	50.4	106.7
5	1	C	4	21.90	80	5	4.2	36.4	0.0	0.0	4.6	27.3	3.5	50.8	114.4
5	1	C	4	21.90	80	14	1.5	37.4	0.0	0.0	1.6	30.3	1.3	46.3	114.1
5	1	C	4	21.90	80	21	4.1	36.8	0.0	0.0	4.7	30.1	5.0	49.9	116.9
5	1	C	15	8.76	0	0	1.3	20.3	0.0	0.0	0.0	0.0	0.3	5.3	25.7
5	1	C	15	8.76	0	5	2.4	23.5	0.0	0.0	0.2	0.7	2.2	7.4	31.6
5	1	C	15	8.76	0	14	2.7	23.0	0.0	0.0	0.0	0.1	0.1	6.3	30.0
5	1	C	15	6.76	0	21	1.1	23.6	0.0	0.0	0.0	0.2	0.3	5.8	29.7
5	1	C	15	8.76	80	0	5.2	23.4	0.0	0.0	4.9	25.4	5.0	47.8	96.6
5	1	C	15	8.76	80	5	5.8	25.6	0.0	0.0	13.3	26.8	9.5	44.6	97.0
5	1	C	15	8.76	80	14	3.3	25.9	0.0	0.0	8.4	32.7	8.1	51.1	109.8
5	1	C	15	8.76	80	21	1.9	32.4	0.0	0.0	16.4	31.7	4.5	57.7	121.8
5	1	C	15	21.90	0	0	2.6	23.0	0.0	0.0	0.1	0.1	0.9	5.4	28.5
5	1	C	15	21.90	0	5	3.1	29.8	0.0	0.0	0.1	0.1	0.3	6.5	36.5
5	1	C	15	21.90	0	14	5.5	36.4	0.0	0.0	0.0	0.1	2.1	3.1	39.7
5	1	C	15	21.90	0	21	2.7	37.3	0.0	0.0	0.0	0.1	1.7	3.4	40.9
5	1	C	15	21.90	80	0	4.6	28.2	0.0	0.5	7.7	27.5	9.1	50.4	106.7
5	1	C	15	21.90	80	5	22.2	45.6	0.0	0.0	6.3	23.0	6.3	39.9	108.6
5	1	C	15	21.90	80	14	43.2	75.2	0.0	0.0	18.0	12.6	26.3	27.4	115.2
5	1	C	15	21.90	80	21	2.0	52.1	0.0	0.0	1.4	22.4	0.3	40.8	115.2
5	1	C	30	8.76	0	0	1.3	20.3	0.0	0.0	0.0	0.0	0.3	5.3	25.7
5	1	C	30	8.76	0	5	0.8	23.7	0.0	0.0	0.0	0.1	0.3	5.8	29.5
5	1	C	30	8.76	0	14	2.6	23.9	0.0	0.0	0.7	0.8	0.3	7.3	32.0
5	1	C	30	8.76	0	21	0.6	25.2	0.0	0.0	0.2	1.4	0.2	7.2	33.6
5	1	C	30	8.76	80	0	5.2	23.4	0.0	0.0	4.9	25.4	5.0	47.8	96.6
5	1	C	30	8.76	80	5	2.7	26.3	0.0	0.0	1.6	24.2	6.1	41.8	92.4
5	1	C	30	8.76	80	14	7.6	28.0	0.0	0.0	6.3	25.9	9.4	48.7	102.6
5	1	C	30	8.76	80	21	1.5	20.9	0.0	0.0	6.1	32.6	2.0	54.0	107.6
5	1	C	30	21.90	0	0	2.8	23.0	0.0	0.0	0.1	0.1	0.9	5.4	28.5
5	1	C	30	21.90	0	5	4.7	41.4	0.0	0.0	0.4	0.3	4.5	3.4	45.1
5	1	C	30	21.00	0	14	1.6	44.0	0.0	0.0	0.0	0.1	3.2	3.7	47.8
5	1	C	30	21.90	0	21	0.9	47.3	0.0	0.0	0.1	0.3	4.2	2.8	50.4
5	1	C	30	21.90	80	0	4.6	28.2	0.0	0.5	7.7	27.5	9.1	50.4	106.7
S	1	C	30	21.90	80	5	8.4	95.8	0.0	0.0	2.3	2.8	0.7	16.4	115.0
5	1	C	30	21.90	80	14	43.0	107.5	0.0	0.0	1.2	0.7	28.6	15.8	124.0
5	1	C	30	21.90	80	21	9.0	109.5	0.0	0.0	3.4	6.8	7.7	17.8	134.1
5	2	C	4	8.70	0	0	0.5	9.8	0.0	0.0	0.0	0.1	0.6	4.3	14.2
5	2	C	4	8.70	0	5	0.8	10.1	0.0	0.0	0.6	0.6	0.5	4.6	15.2
5	2	C	4	8.70	0	14	0.2	12.0	0.0	0.0	0.0	0.1	0.2	4.1	16.2
S	2	C	4	8.70	0	21	2.2	9.2	0.0	0.0	1.2	1.0	0.1	4.8	15.0
5	2	C	4	8.70	80	0	1.8	7.2	0.0	0.0	4.0	24.9	1.3	44.3	76.4
5	2	C	4	8.70	80	5	3.8	21.5	0.0	0.0	0.3	23.5	3.3	42.0	87.1
5	2	C	4	8.70	80	14	2.0	22.9	0.0	0.0	0.5	22.3	0.4	43.8	88.9
5	2	C	4	8.70	80	21	2.1	9.1	0.0	0.0	2.3	30.3	2.7	49.0	88.4
5	2	C	4	21.80	0	0	1.1	11.1	0.0	0.0	0.1	0.3	0.1	5.1	16.5
5	2	C	4	21.80	0	5	0.4	17.0	0.0	0.0	0.1	0.3	0.1	4.3	21.6
5	2	C	4	21.80	0	14	1.4	23.0	0.0	0.0	0.0	0.1	0.0	1.4	24.5
5	2	C	4	21.80	0	21	0.9	23.1	0.0	0.0	0.1	0.1	2.5	2.6	25.6
5	2	C	4	21.80	80	0	2.3	12.4	0.5	0.3	2.6	26.8	6.8	48.5	88.0
5	2	C	4	21.80	80	5	1.7	19.9	0.5	0.3	1.9	24.8	3.3	45.3	90.2
5	2	C	4	21.80	80	14	2.3	25.8	0.5	0.3	1.4	22.9	0.9	40.3	89.1
S	2	C	4	21.80	80	21	3.7	26.0	0.0	0.0	2.8	26.4	6.0	40.4	92.8
S	2	C	15	8.70	0	0	0.5	9.8	0.0	0.0	0.0	0.1	0.6	4.3	14.2
5	2	C	15	8.70	0	5	0.3	9.8	0.0	0.0	0.0	0.9	0.3	4.4	15.1
5	2	C	15	8.70	0	14	0.2	9.3	0.0	0.0	0.0	0.1	0.1	3.6	13.0
5	2	C	15	8.70	0	21	1.2	9.4	0.0	0.0	0.3	1.0	0.2	3.4	13.8
5	2	C	15	8.70	80	0	1.8	7.2	0.0	0.0	4.0	24.9	1.3	44.3	76.4
5	2	C	15	8.70	80	5	1.4	10.9	0.0	0.0	15.9	35.9	13.4	56.8	103.6
5	2	C	15	8.70	80	14	3.2	11.0	0.0	0.0	1.4	28.9	2.2	51.8	91.7
5	2	C	15	8.70	80	21	6.4	14.4	0.0	0.0	6.5	30.9	4.6	48.1	93.3
5	2	C	15	21.80	0	0	1.1	11.1	0.0	0.0	0.1	0.3	0.1	5.1	16.5

5	2	C	15	21.80	0	5	0.4	22.5	0.0	0.0	0.0	0.1	1.6	2.2	24.8
5	2	C	15	21.80	0	14	2.3	29.6	0.0	0.0	0.0	0.1	0.2	0.7	30.4
5	2	C	15	21.80	0	21	0.6	29.5	0.0	0.0	0.0	0.1	0.1	1.3	30.9
5	2	C	15	21.80	80	0	2.3	12.4	0.5	0.3	2.6	26.8	6.3	48.5	88.0
5	2	C	15	21.80	80	5	6.1	33.9	0.0	0.0	1.7	23.3	1.3	43.0	100.2
5	2	C	15	21.80	80	14	24.8	92.6	0.0	0.0	4.0	2.6	8.1	10.8	106.0

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NOR		SOL NH4		EX NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
5	2	C	30	8.70	0	0	0.5	9.8	0.0	0.0	0.0	0.1	0.6	4.3	14.2
5	2	C	30	8.70	0	5	1.1	9.1	0.0	0.0	0.5	0.5	0.0	4.4	14.1
5	2	C	30	8.70	0	14	0.3	9.4	0.0	0.0	0.1	0.1	0.5	4.9	14.5
5	2	C	30	8.70	0	21	0.4	10.3	0.0	0.0	0.6	1.7	0.5	6.1	18.1
5	2	C	30	8.70	80	0	1.8	7.2	0.0	0.0	4.0	24.9	1.3	44.3	76.4
5	2	C	30	8.70	80	5	6.3	12.1	0.0	0.0	0.8	26.5	4.2	46.1	84.7
5	2	C	30	8.70	80	14	9.1	12.3	0.0	0.0	1.4	24.0	1.7	45.6	82.0
5	2	C	30	8.70	80	21	10.0	14.8	0.0	0.0	3.9	23.6	1.0	41.9	80.3
5	2	C	30	21.80	0	0	1.1	11.1	0.0	0.0	0.1	0.3	0.1	5.1	16.5
5	2	C	30	21.80	0	5	0.6	30.8	0.0	0.0	0.1	0.1	0.1	0.6	31.6
5	2	C	30	21.80	0	14	1.8	35.5	0.0	0.0	0.0	0.1	0.0	0.7	36.3
5	2	C	30	21.80	0	21	4.0	40.1	0.0	0.0	2.1	1.1	7.0	6.0	47.1
5	2	C	30	21.80	80	0	2.3	12.4	0.0	0.0	2.6	26.8	6.8	48.5	87.8
5	2	C	30	21.80	80	5	8.1	77.1	0.0	0.0	0.7	3.6	0.9	13.8	94.5
5	2	C	30	21.80	80	14	6.6	102.7	0.0	0.0	1.1	0.6	7.8	5.0	108.3
5	2	C	30	21.80	80	21	28.8	101.6	0.0	0.0	3.7	2.3	10.5	6.6	110.5
5	1	CB	4	9.24	0	0	0.1	17.1	0.0	0.0	0.0	0.0	0.2	6.2	23.3
5	1	CB	4	9.24	0	5	0.7	16.3	0.0	0.0	0.2	1.2	0.9	6.9	26.4
5	1	CB	4	9.24	0	14	2.1	15.3	2.5	8.8	0.0	0.5	0.1	7.4	31.9
5	1	CB	4	9.24	0	21	1.7	16.4	0.0	0.0	0.7	0.3	0.0	7.6	24.4
5	1	CB	4	9.24	80	0	1.6	15.8	0.0	0.0	2.6	25.9	7.8	50.8	92.5
5	1	CB	4	9.24	80	5	1.6	14.5	0.0	0.0	14.9	31.4	5.4	56.2	102.1
5	1	CB	4	9.24	80	14	1.6	15.9	1.0	6.5	6.1	23.9	4.8	52.4	100.7
5	1	CB	4	9.24	80	21	2.7	16.5	0.0	0.0	7.5	30.3	6.2	55.9	102.7
5	1	CB	4	23.10	0	0	0.4	20.0	0.0	0.0	0.0	0.0	0.3	5.9	25.9
5	1	CB	4	23.10	0	5	0.9	28.9	0.0	0.0	0.0	0.0	0.0	4.5	33.4
5	1	CB	4	23.10	0	14	4.0	21.7	2.0	5.0	0.0	0.0	1.9	4.3	31.0
5	1	CB	4	23.10	0	21	0.6	35.1	0.0	0.0	0.0	0.0	1.0	3.1	38.2
5	1	CB	4	23.10	80	0	4.3	20.4	0.5	0.8	0.9	27.6	0.9	50.6	99.5
5	1	CB	4	23.10	80	5	2.2	30.4	0.5	0.3	1.9	24.3	5.8	50.6	105.6
5	1	CB	4	23.10	80	14	1.3	43.8	0.0	0.0	0.5	10.9	0.9	41.9	102.7
5	1	CB	4	23.10	80	21	2.0	44.4	0.0	0.0	2.8	20.0	0.2	31.0	95.4
5	1	CB	15	9.24	0	0	0.1	17.1	0.0	0.0	0.0	0.0	0.2	6.2	23.3
5	1	CB	15	9.24	0	5	1.2	18.1	0.0	0.0	0.4	0.9	1.8	7.2	26.2
5	1	CB	15	9.24	0	14	0.8	15.8	4.5	7.8	0.3	0.3	0.4	7.5	31.3
5	1	CB	15	9.24	0	21	0.5	16.6	10.0	5.0	0.0	0.5	0.8	8.3	30.3
5	1	CB	15	9.24	80	0	1.6	15.8	0.0	0.0	2.6	25.9	7.8	50.8	92.5
5	1	CB	15	9.24	80	5	2.6	16.7	0.0	0.0	23.4	33.0	33.1	54.4	104.1
5	1	CB	15	9.24	80	14	0.5	14.1	5.5	2.8	3.1	22.3	4.5	45.3	84.4
5	1	CB	15	9.24	80	21	3.7	16.8	0.0	0.0	3.2	28.9	9.3	52.8	98.4
5	1	CB	15	23.10	0	0	0.4	20.0	0.0	0.0	0.0	0.0	0.8	5.9	25.9
5	1	CB	15	23.10	0	5	19.2	40.9	0.0	0.0	2.2	1.1	1.7	0.8	42.8
5	1	CB	15	23.10	0	14	6.5	32.3	2.0	1.0	2.3	1.3	1.3	3.3	37.9
5	1	CB	15	23.10	0	21	0.6	34.8	0.0	0.0	1.0	1.5	3.9	1.9	38.2
5	1	CB	15	23.10	80	0	4.3	20.4	0.5	0.8	0.9	27.6	0.9	50.6	99.5
5	1	CB	15	23.10	80	5	7.1	42.9	0.0	0.0	8.7	17.1	6.3	37.6	97.7
5	1	CB	15	23.10	80	14	36.8	98.6	21.0	18.5	1.9	1.8	4.3	13.3	120.3
5	1	CB	15	23.10	80	21	10.9	92.6	0.0	0.0	0.0	0.0	0.5	3.1	95.7
5	1	CB	30	9.24	0	0	0.1	17.1	0.0	0.0	0.0	0.0	0.2	6.2	23.3
5	1	CB	30	9.24	0	5	0.6	16.8	0.0	0.0	0.9	0.4	0.4	6.2	23.4
5	1	CB	30	9.24	0	14	0.0	17.0	2.5	6.3	0.1	0.3	1.1	7.9	31.4
5	1	CB	30	9.24	0	21	0.7	16.0	0.0	0.0	1.2	1.2	1.2	7.9	25.1
5	1	CB	30	9.24	80	0	1.6	15.8	0.0	0.0	2.6	25.9	7.8	50.8	92.5
5	1	CB	30	9.24	80	5	1.8	17.3	0.0	0.0	0.1	21.8	2.8	41.8	80.8
5	1	CB	30	9.24	80	14	1.9	21.1	0.0	0.0	3.8	23.8	3.5	45.1	90.0
5	1	CB	30	9.24	80	21	4.5	19.6	0.0	0.0	0.0	20.5	8.7	38.8	78.9
5	1	CB	30	23.10	0	0	0.4	20.0	0.0	0.0	0.0	0.0	0.8	5.9	25.9
5	1	CB	30	23.10	0	5	0.8	37.1	0.0	0.0	0.0	0.0	0.8	1.5	38.6
5	1	CB	30	23.10	0	14	3.0	36.2	3.0	1.5	0.0	0.0	0.1	2.1	39.7
5	1	CB	30	23.10	0	21	3.2	42.6	0.0	0.0	0.0	0.0	1.0	2.4	45.0
5	1	CB	30	23.10	80	0	4.3	20.4	0.5	0.8	0.9	27.6	0.9	50.6	99.5
5	1	CB	30	23.10	80	5	5.5	79.6	0.0	0.0	6.4	3.2	7.6	15.1	97.9
5	1	CB	30	23.10	80	14	22.0	114.0	0.0	0.0	4.9	2.7	2.0	7.8	124.5
5	1	CB	30	23.10	80	21	1.0	107.5	0.0	0.0	1.6	2.3	1.9	9.5	119.3
5	2	CB	4	9.69	0	0	1.9	16.6	0.0	0.0	0.0	0.1	0.0	5.4	22.1
5	2	CB	4	9.69	0	5	0.2	13.8	3.0	11.5	0.3	0.6	0.4	8.0	33.9
5	2	CB	4	9.69	0	14	1.8	16.8	2.0	1.0	0.1	0.3	0.0	7.4	25.5
5	2	CB	4	9.69	0	21	1.7	15.9	0.3	0.0	0.4	0.3	1.0	7.2	23.4
5	2	CB	4	9.69	80	0	2.2	20.3	0.0	0.0	11.5	27.4	15.4	48.7	96.4
5	2	CB	4	9.69	80	5	2.9	18.1	11.0	5.5	14.5	30.3	16.5	52.8	106.6

5	2	CB	4	9.69	80	14	1.0	15.5	13.5	6.8	1.2	26.3	3.1	50.8	99.3
5	2	CB	4	9.69	80	21	0.8	17.0	0.0	0.0	4.2	18.9	5.0	42.7	78.6
5	2	CB	4	24.20	0	0	1.5	21.1	0.0	0.0	0.1	0.1	0.0	7.0	28.2
5	2	CB	4	24.20	0	5	8.7	20.5	19.5	9.8	0.0	0.7	1.0	8.9	39.9
5	2	CB	4	24.20	0	14	1.2	29.1	0.0	0.0	0.2	0.1	0.1	6.1	35.3
5	2	CB	4	24.20	0	21	3.0	32.9	0.0	0.0	0.0	0.1	2.6	6.7	39.7
5	2	CB	4	24.20	80	0	3.1	24.4	0.0	0.0	0.1	28.4	0.6	50.4	103.3
5	2	CB	4	24.20	80	5	0.5	16.9	3.0	19.0	3.6	30.9	4.6	52.1	118.9
5	2	CB	4	24.20	80	14	1.9	36.3	0.0	0.0	3.5	22.6	2.6	46.9	105.9
5	2	CB	4	24.20	80	21	1.8	34.4	0.0	0.0	0.0				

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		SOL	NH4	EX NH4		TOTAL
							RANGE	AVER	RANGE	AVER			RANGE	AVER	
5	2	CB	15	9.69	0	0	1.9	16.6	0.0	0.0	0.0	0.1	0.0	5.4	22.1
5	2	CB	15	9.69	0	5	2.6	12.1	12.0	6.0	0.0	0.9	1.8	9.0	23.0
5	2	CB	15	9.69	0	14	2.4	16.1	0.0	21.5	0.2	0.4	0.1	6.8	44.8
5	2	CB	15	9.69	0	21	4.3	19.8	0.0	0.0	0.2	0.4	1.2	9.8	30.0
5	2	CB	15	9.69	80	0	2.2	20.3	0.0	0.0	11.5	27.4	15.4	48.7	96.4
5	2	CB	15	9.69	80	5	0.6	11.7	0.0	0.0	40.4	42.1	1.2	44.1	97.9
5	2	CB	15	9.69	80	14	9.4	17.5	0.0	0.0	1.9	27.6	4.4	54.4	99.5
5	2	CB	15	9.69	80	21	0.3	17.1	0.0	0.0	5.6	25.8	4.2	51.3	94.2
5	2	CB	15	24.20	0	0	1.5	21.1	0.0	0.0	0.1	0.1	0.0	7.0	28.2
5	2	CB	15	24.20	0	5	3.5	26.6	8.5	4.3	0.4	0.2	0.3	6.0	37.1
5	2	CB	15	24.20	0	14	2.9	40.1	0.0	0.0	0.1	0.3	0.2	3.2	43.5
5	2	CB	15	24.20	0	21	0.1	40.8	0.0	0.0	0.1	0.3	0.2	5.5	46.6
5	2	CB	15	24.20	80	0	3.1	24.4	0.0	0.0	0.1	26.4	0.6	50.4	103.3
5	2	CB	15	24.20	80	5	10.1	36.4	0.0	0.0	3.8	17.6	2.0	39.5	93.5
5	2	CB	15	24.20	80	14	3.7	67.3	10.5	5.3	3.7	12.6	7.5	29.4	114.6
5	2	CB	15	24.20	80	21	19.7	85.9	0.0	0.0	5.3	7.5	5.6	23.4	116.9
5	2	CB	30	9.69	0	0	1.9	16.6	0.0	0.0	0.0	0.1	0.0	5.4	22.1
5	2	CB	30	9.69	0	5	2.6	15.7	7.0	3.5	0.1	0.6	0.9	7.8	27.7
5	2	CB	30	9.69	0	14	0.5	18.6	0.0	0.0	0.0	1.1	0.1	8.8	28.5
5	2	CB	30	9.69	0	21	0.4	18.1	0.0	0.0	0.0	0.5	0.7	11.0	29.6
5	2	CB	30	9.9	80	0	2.2	20.3	0.0	0.0	11.5	27.4	15.4	48.7	96.4
5	2	CB	30	9.69	80	5	6.6	11.4	0.0	0.0	11.8	25.0	16.7	48.3	84.6
5	2	CB	30	9.69	60	14	3.1	19.9	0.0	0.0	17.9	30.8	23.7	60.9	111.7
5	2	CB	30	9.69	80	21	1.6	17.1	8.5	4.3	2.6	25.4	1.2	51.0	97.7
5	2	CB	30	24.20	0	0	1.5	21.1	0.0	0.0	0.1	0.1	0.0	7.0	28.2
5	2	CB	30	24.20	0	5	10.9	32.9	11.0	5.5	0.0	0.2	0.5	10.4	49.1
5	2	CB	30	24.20	0	14	3.3	43.9	7.0	13.0	0.1	0.3	1.3	9.5	66.7
5	2	CB	30	24.20	0	21	0.3	51.3	0.0	0.0	0.2	0.3	0.2	10.7	62.2
5	2	CB	30	24.20	80	0	3.1	24.4	0.0	0.0	0.1	28.4	0.8	50.4	103.3
5	2	CB	30	24.20	80	5	4.8	56.2	10.5	5.3	4.4	16.4	3.0	37.3	115.1
5	2	CB	30	24.20	80	14	6.6	69.1	6.0	8.5	1.4	11.5	2.1	32.4	141.5
5	2	CB	30	24.20	80	21	3.5	96.1	0.0	0.0	2.9	7.8	4.9	24.6	128.6
5	3	C	4	8.82	0	0	0.7	7.5	0.0	0.0	0.5	0.4	0.0	3.0	11.0
5	3	C	4	8.82	0	5	6.0	3.0	5.5	2.6	0.1	0.3	0.6	3.8	9.6
5	3	C	4	5.82	0	10	1.4	6.2	0.0	0.0	0.0	0.1	0.0	2.9	9.2
5	3	C	4	8.82	0	21	0.1	7.6	0.0	0.0	0.4	0.4	0.6	3.5	11.6
5	3	C	4	6.82	80	0	0.3	6.3	0.0	0.0	6.2	19.6	9.7	34.9	62.9
5	3	C	4	6.82	80	5	0.5	5.1	4.6	2.3	10.0	21.5	16.1	38.8	67.6
5	3	C	4	8.82	80	14	14.3	13.2	0.0	0.0	2.7	18.8	11.4	28.2	60.3
5	3	C	4	8.82	80	21	1.2	20.0	0.0	0.0	1.7	20.8	19.0	29.8	70.6
5	3	C	4	22.10	0	0	1.1	9.3	0.0	0.0	0.4	0.7	2.1	5.6	15.5
5	3	C	4	22.10	0	5	0.9	11.1	2.0	1.0	0.0	0.1	0.3	4.8	17.0
5	3	C	4	22.10	0	14	0.8	17.0	0.0	0.0	0.0	0.5	0.2	2.4	19.9
5	3	C	4	22.10	0	21	2.2	19.4	0.0	0.0	0.1	0.1	0.1	2.8	22.4
5	3	C	4	22.10	81	0	0.5	10.0	0.0	0.0	3.9	23.6	1.5	40.6	74.2
5	3	C	4	22.10	80	5	0.8	7.4	6.0	6.5	6.7	12.5	2.9	40.1	66.5
5	3	C	4	22.10	80	14	2.7	10.3	0.5	0.3	13.5	15.8	1.2	36.2	62.5
5	3	C	4	22.10	80	21	1.8	7.1	0.0	1.5	18.0	27.7	5.9	36.8	73.1
5	3	C	15	8.82	0	0	0.7	7.5	0.0	0.0	0.5	0.4	0.0	3.0	11.0
5	3	C	15	8.82	0	5	0.9	5.8	6.0	3.0	0.2	0.6	0.2	3.5	12.8
5	3	C	15	8.82	0	14	2.8	5.1	3.0	1.5	0.0	0.1	0.2	3.1	9.8
5	3	C	15	8.82	0	21	0.0	8.1	0.0	0.0	0.0	0.1	0.3	3.1	11.3
5	3	C	15	8.82	80	0	0.3	8.3	0.0	0.0	6.2	19.6	9.7	34.9	62.9
5	3	C	15	8.62	80	5	1.0	6.5	0.0	0.0	17.4	26.9	18.6	44.3	77.7
5	3	C	15	8.82	80	14	0.4	7.4	0.0	0.0	3.1	15.6	2.2	30.9	53.8
5	3	C	15	8.82	80	21	2.2	9.3	0.0	0.0	4.7	12.0	7.0	24.0	45.3
5	3	C	15	22.10	0	0	1.1	9.3	0.0	0.0	0.4	0.7	2.1	5.6	15.5
5	3	C	15	22.10	0	5	1.3	16.8	0.0	0.0	0.2	0.2	0.2	3.2	20.1
5	3	C	15	22.10	0	14	0.6	24.0	0.0	0.0	0.0	0.1	0.2	1.0	25.1
5	3	C	15	22.10	0	21	1.7	30.1	0.0	0.0	0.4	0.3	1.4	2.3	32.7
5	3	C	15	22.10	80	0	0.5	10.0	0.0	0.0	3.9	23.6	1.5	40.6	74.2
5	3	C	15	22.10	80	5	8.2	22.2	8.5	6.8	2.1	19.8	2.4	36.9	85.6
5	3	C	15	22.10	80	14	2.8	40.7	0.0	0.0	3.2	9.7	1.6	22.1	80.5
5	3	C	15	22.10	80	21	1.7	65.9	0.0	0.0	0.7	3.8	2.1	11.8	81.6
5	3	C	30	8.82	0	0	0.7	7.5	0.0	0.0	0.5	0.4	0.0	3.0	11.0
5	3	C	30	8.82	0	5	1.1	7.3	0.0	0.0	0.3	0.3	0.1	3.4	11.1

5	3	C	30	8.82	0	14	0.5	7.1	0.0	0.0	0.0	0.1	0.6	2.9	10.0
5	3	C	30	8.82	0	21	0.1	8.0	0.0	0.0	0.6	1.0	0.6	3.5	12.5
5	3	C	30	8.82	80	0	0.3	8.3	0.0	0.0	6.2	19.6	9.7	34.9	62.9
5	3	C	30	8.82	80	5	4.7	3.4	5.5	2.8	14.8	21.1	18.3	35.6	63.1
S	3	C	30	8.82	80	14	22.4	22.6	0.0	0.0	16.4	8.2	1.9	34.8	65.6
5	3	C	30	8.82	80	21	1.5	9.6	0.0	0.0	2.4	19.8	4.2	31.9	61.3
5	3	C	30	22.10	0	0	1.1	9.3	0.0	0.0	0.4	0.7	2.1	5.6	15.5
5	3	C	30	22.10	0	5	6.4	24.9	0.0	0.0	0.2	0.2	1.7	2.3	27.4
5	3	C	30	22.10	0	14	4.4	40.6	0.0	0.0	0.0	0.1	0.4	0.7	41.4
5	3	C	30	22.10	0	21	1.4	42.8	0.0	0.0	0.1	0.6	0.0	1.0	44.4
5	3	C	30	22.10	80	0	0.5	10.0	0.0	0.0	3.9	23.6	1.5	40.6	74.2
5	S	C	30	22.10	80	5	2.1	47.6	0.0	0.0	0.7	9.4	2.7	27.3	84.3
5	3	C	30	22.10	80	14	0.9	92.3	0.0	0.0	0.5	0.4	1.3	5.1	97.9
5	3	C	30	22.10	80	21	14.1	103.9	0.0	0.0	0.1	0.1	0.1	1.6	105.6

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	
5	4	C	4	8.40	0	0	0.8	4.6	0.0	0.0	0.3	1.5	6.1
5	4	C	4	8.40	0	14	0.4	4.7	0.0	0.0	0.1	1.4	6.1
5	4	C	4	8.40	0	21	0.2	6.8	0.0	0.0	0.1	1.8	8.5
5	4	C	4	8.40	80	0	3.4	1.7	0.0	0.0	5.9	40.6	42.3
5	4	C	4	8.40	80	14	1.7	3.5	0.0	0.0	11.9	35.8	39.4
5	4	C	4	8.40	60	21	9.6	11.1	0.0	0.0	21.1	43.1	54.2
5	4	C	4	21.00	0	0	1.7	4.6	0.0	0.0	0.0	1.6	6.3
S	4	C	4	21.00	0	14	0.1	5.9	0.0	0.0	0.0	3.6	9.6
5	4	C	4	21.00	0	21	0.3	10.8	0.0	0.0	0.7	3.9	14.7
5	4	C	4	21.00	80	0	0.6	5.5	0.0	0.0	13.5	55.1	60.6
5	4	C	4	21.00	80	14	2.8	7.5	0.0	0.0	15.7	62.0	69.5
5	4	C	4	21.00	80	21	0.9	9.8	0.0	0.0	26.5	56.6	66.4
5	4	C	15	8.40	0	0	0.8	4.6	0.0	0.0	0.3	1.5	6.1
5	4	C	15	8.40	0	14	0.6	4.6	0.0	0.0	0.1	1.4	6.0
5	4	C	15	8.40	0	21	0.4	6.3	0.0	0.0	0.4	2.2	8.5
5	4	C	15	8.40	80	0	3.4	1.7	0.0	0.0	5.7	40.8	42.4
5	4	C	15	8.40	80	14	3.5	7.6	0.0	0.0	16.7	43.6	51.3
5	4	C	15	8.40	80	21	10.0	5.0	0.0	0.0	11.5	41.9	46.9
5	4	C	15	21.00	0	0	1.7	4.6	0.0	0.0	0.0	1.6	6.3
5	4	C	15	21.00	0	14	0.5	6.9	0.0	0.0	0.0	5.5	12.4
5	4	C	15	21.00	0	21	0.7	14.4	0.0	0.0	0.0	5.9	20.3
5	4	C	15	21.00	80	0	0.6	5.5	0.0	0.0	13.5	55.1	60.6
5	4	C	15	21.00	80	14	0.3	13.1	0.0	0.0	3.5	51.6	64.7
5	4	C	15	21.00	80	21	13.7	6.8	0.0	0.0	19.0	49.9	56.7
5	4	C	30	8.40	0	0	0.8	4.6	0.0	0.0	0.3	1.5	6.1
5	4	C	30	8.40	0	14	0.0	4.0	0.0	0.0	0.0	1.6	5.6
5	4	C	30	8.40	0	21	0.2	4.3	0.0	0.0	0.2	1.7	6.0
5	4	C	30	8.40	80	0	3.4	1.7	0.0	0.0	5.7	40.6	42.4
5	4	C	30	8.40	80	14	5.6	6.0	0.0	0.0	8.3	36.8	42.8
5	4	C	30	8.40	80	21	2.8	5.3	0.0	0.0	1.0	40.5	45.8
5	4	C	30	21.00	0	0	1.7	4.6	0.0	0.0	0.0	1.6	6.3
5	4	C	30	21.00	0	14	0.6	10.6	0.0	0.0	0.2	5.5	16.1
5	4	C	30	21.00	0	21	0.3	11.5	0.0	0.0	0.8	5.9	17.4
5	4	C	30	21.00	80	0	0.6	5.5	0.0	0.0	13.5	55.1	60.6
5	4	C	30	21.00	80	14	0.5	13.6	0.0	0.0	14.0	60.4	74.0
5	4	C	30	21.00	80	21	7.4	4.1	0.0	0.0	21.9	61.1	65.1
5	5	C	4	8.30	0	0	0.0	5.0	0.0	0.0	0.0	1.1	6.1
5	5	C	4	8.30	0	14	0.3	3.5	0.0	0.0	0.3	1.5	5.1
5	5	C	4	8.30	0	21	0.0	4.4	0.0	0.0	0.4	1.8	6.2
5	5	C	4	8.30	80	0	1.0	2.9	0.0	0.0	9.0	36.2	39.1
5	5	C	4	8.30	80	14	2.7	4.4	0.0	0.0	4.3	31.0	35.5
5	5	C	4	8.30	80	21	0.1	2.4	0.0	0.0	2.4	35.1	37.5
5	5	C	4	20.90	0	0	0.4	4.3	0.0	0.0	0.0	0.9	5.2
5	5	C	4	20.90	0	14	0.2	4.6	0.0	0.0	0.2	1.9	6.5
5	5	C	4	20.90	0	21	0.8	4.3	0.0	0.0	0.1	2.3	6.5
5	5	C	4	20.90	80	0	1.2	4.3	0.0	0.0	12.8	36.6	40.9
5	5	C	4	20.90	80	14	0.9	2.8	0.0	0.0	5.0	44.5	47.3
5	5	C	4	20.90	80	21	0.3	1.8	0.0	0.0	4.2	39.4	41.2
5	5	C	15	8.30	0	0	0.0	5.0	0.0	0.0	0.0	1.1	6.1
5	5	C	15	8.30	0	14	0.4	4.1	0.0	0.0	0.0	1.4	5.5
5	5	C	15	8.30	0	21	0.5	4.3	0.0	0.0	0.2	1.5	5.8
5	5	C	15	8.30	80	0	1.0	2.9	0.0	0.0	9.0	36.2	39.1
5	5	C	15	8.30	80	14	3.0	4.9	0.0	0.0	10.7	43.3	48.2
5	5	C	15	8.30	80	21	2.4	5.1	0.0	0.0	4.9	29.6	34.7
5	5	C	15	20.90	0	0	0.4	4.3	0.0	0.0	0.0	0.9	5.2
5	5	C	15	20.90	0	14	0.4	5.0	0.0	0.0	0.4	2.9	8.0
5	5	C	15	20.90	0	21	0.3	6.0	0.0	0.0	0.4	2.9	8.9
5	5	C	15	20.90	80	0	1.2	4.3	0.0	0.0	12.8	36.6	40.9
5	5	C	15	20.90	80	14	0.8	5.8	0.0	0.0	10.2	46.0	51.8
5	5	C	15	20.90	80	21	1.6	7.0	0.0	0.0	3.9	34.6	41.6
5	5	C	30	8.30	0	0	0.0	5.0	0.0	0.0	0.0	1.1	6.1
5	5	C	30	8.30	0	14	1.1	4.6	0.0	0.0	0.0	1.6	6.2

5	5	C	30	8.30	0	21	0.3	3.9	0.0	0.0	0.2	1.7	5.6
5	5	C	30	8.30	80	0	1.0	2.9	0.0	0.0	9.0	36.2	39.1
5	5	C	30	8.30	80	14	0.1	4.9	0.0	0.0	32.3	49.3	54.2
5	5	C	30	8.30	80	21	1.5	5.9	0.0	0.0	5.8	33.2	39.1
5	5	C	30	20.90	0	0	0.4	4.3	0.0	0.0	0.0	0.9	5.2
5	5	C	30	20.90	0	14	0.2	5.1	0.0	0.0	0.7	2.9	8.0
5	5	C	30	20.90	0	21	0.7	6.3	0.0	0.0	0.1	2.4	8.7
5	5	C	30	20.90	80	0	1.2	4.3	0.0	0.0	12.8	36.6	40.9
5	5	C	30	20.90	80	14	0.7	6.1	0.0	0.0	11.5	36.9	45.0
5	5	C	30	20.90	80	21	8.4	4.3	0.0	0.0	27.7	53.1	57.4

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	
5	6	C	4	8.20	0	0	0.3	4.3	0.0	0.0	0.1	0.8	5.2
5	6	C	4	8.20	0	14	1.0	3.8	0.0	0.0	0.1	1.8	5.6
5	6	C	4	8.20	0	21	0.3	4.5	0.0	0.0	0.3	1.0	5.6
5	6	C	4	8.20	80	0	4.0	3.4	0.0	0.0	7.2	25.8	29.2
5	6	C	4	8.20	80	14	0.7	5.8	0.0	0.0	7.0	29.4	35.1
5	6	C	4	8.20	80	21	0.4	6.6	0.0	0.0	8.4	39.6	46.2
5	6	C	4	20.40	0	0	0.4	4.4	0.0	0.0	0.0	0.6	5.0
5	6	C	4	20.40	0	14	0.5	3.8	0.0	0.0	0.1	1.8	5.6
5	6	C	4	20.40	0	21	0.1	4.4	0.0	0.0	0.0	0.7	5.1
5	6	C	4	20.40	80	0	0.3	1.6	0.0	0.0	1.0	27.8	29.4
5	6	C	4	20.40	80	14	1.2	6.5	0.0	0.0	2.2	29.4	35.9
5	6	C	4	20.40	80	21	0.6	6.0	0.0	0.0	7.4	29.9	35.9
5	6	C	15	8.20	0	0	0.3	4.3	0.0	0.0	0.1	0.8	5.2
5	6	C	15	8.20	0	14	0.4	4.1	0.0	0.0	0.0	1.7	5.8
5	6	C	15	8.20	0	21	0.2	4.3	0.0	0.0	0.4	0.9	5.2
5	6	C	15	8.20	80	0	4.0	3.4	0.0	0.0	7.2	25.8	29.2
5	6	C	15	8.20	60	14	2.1	6.1	0.0	0.0	5.4	31.3	37.3
5	6	C	15	8.20	80	21	1.4	5.6	0.0	0.0	2.7	32.6	38.4
5	6	C	15	20.40	0	0	0.4	4.4	0.0	0.0	0.0	0.6	5.0
5	6	C	15	20.40	0	14	0.1	4.3	0.0	0.0	0.2	1.9	6.2
5	6	C	15	20.40	0	21	0.2	6.0	0.0	0.0	0.2	1.1	7.1
5	6	C	15	20.40	80	0	0.3	1.6	0.0	0.0	1.0	27.8	29.4
5	6	C	15	20.00	80	14	1.0	6.4	0.0	0.0	0.9	30.8	37.2
5	6	C	15	20.40	80	21	1.7	7.4	0.0	0.0	3.3	24.6	32.3
5	6	C	30	8.20	0	0	0.3	4.3	0.0	0.0	0.1	0.8	5.2
5	6	C	30	8.20	0	14	0.3	3.1	0.0	0.0	0.1	1.8	4.9
5	6	C	30	8.20	0	21	0.4	4.4	0.0	0.0	0.7	1.5	5.9
5	6	C	30	8.20	80	0	4.0	3.4	0.0	0.0	7.2	25.8	29.2
5	6	C	30	8.20	80	14	1.0	5.6	0.0	0.0	4.8	27.2	32.8
5	6	C	30	8.20	80	21	0.3	5.0	0.0	0.0	6.8	26.7	31.7
5	6	C	30	20.40	0	0	0.4	4.4	0.0	0.0	0.0	0.6	5.0
5	6	C	30	20.40	0	14	0.2	5.7	0.0	0.0	0.0	2.0	7.7
5	6	C	30	20.40	0	21	1.0	7.4	0.0	0.0	0.6	1.8	9.2
5	6	C	30	20.40	80	0	0.3	1.4	0.0	0.0	1.0	27.8	29.4
5	6	C	30	20.40	80	14	1.6	6.2	0.0	0.0	1.2	27.3	35.5
5	6	C	30	20.40	80	21	0.6	7.8	0.0	0.0	2.8	26.1	33.9
5	7	C	4	8.00	0	0	0.2	3.8	0.0	0.0	0.0	1.2	5.0
5	7	C	4	8.40	0	14	0.2	3.4	0.0	0.0	0.0	0.9	4.3
5	7	C	4	8.40	0	21	0.0	4.0	0.0	0.0	0.1	1.1	5.1
5	7	C	4	8.40	80	0	0.6	4.5	0.0	0.0	13.1	42.6	47.1
5	7	C	4	8.40	80	14	3.2	5.2	0.0	0.0	2.2	32.6	37.8
5	7	C	4	8.40	80	21	0.2	5.8	0.0	0.0	5.2	41.6	47.4
5	7	C	4	20.90	0	0	0.7	4.0	0.0	0.0	0.1	0.6	4.7
5	7	C	4	20.90	0	14	0.6	4.8	0.0	0.0	0.1	0.6	5.4
5	7	C	4	20.90	0	21	1.7	5.6	0.0	0.0	0.0	1.2	6.8
5	7	C	4	20.90	80	0	0.1	4.1	0.0	0.0	5.1	38.1	42.3
5	7	C	4	20.90	80	14	1.0	5.2	0.0	0.0	1.1	31.6	36.8
5	7	C	4	20.90	80	21	0.0	7.3	0.0	0.0	0.0	31.0	38.3
5	7	C	15	8.40	0	0	0.2	3.8	0.0	0.0	0.0	1.2	5.0
5	7	C	15	8.40	0	14	0.7	4.6	0.0	0.0	0.8	0.6	5.2
5	7	C	15	8.40	0	21	0.7	3.6	0.0	0.0	0.8	0.6	4.2
5	7	C	15	8.40	80	0	0.6	4.5	0.0	0.0	13.1	42.6	47.1
5	7	C	15	8.40	80	14	2.3	3.3	0.0	0.0	3.5	28.6	32.0
5	7	C	15	8.40	80	21	3.6	3.2	0.0	0.0	24.7	25.6	28.8
5	7	C	15	20.90	0	0	0.7	4.0	0.0	0.0	0.1	0.6	4.7
5	7	C	15	20.90	0	14	1.6	5.1	0.0	0.0	0.9	0.7	5.8
5	7	C	15	20.90	0	21	1.0	5.3	0.0	0.0	0.3	1.8	7.1
5	7	C	15	20.90	80	0	0.1	4.1	0.0	0.0	5.1	3.1	42.3
5	7	C	15	20.90	80	14	0.7	7.4	0.0	0.0	1.8	25.7	33.1
5	7	C	15	20.90	80	21	2.2	7.9	0.0	0.0	7.0	29.2	37.1
5	7	C	30	8.40	0	0	0.2	3.8	0.0	0.0	0.0	1.2	5.0
5	7	C	30	8.40	0	14	0.6	4.2	0.0	0.0	0.1	1.3	5.4
5	7	C	30	8.40	0	21	0.2	4.1	0.0	0.0	0.1	1.4	5.5
5	7	C	30	8.40	80	0	0.6	4.5	0.0	0.0	13.1	42.6	47.1
5	7	C	30	8.40	80	14	0.4	6.0	0.0	0.0	8.2	33.9	39.9
5	7	C	30	8.40	80	21	1.8	5.0	0.0	0.0	3.3	24.5	29.5

5	7	C	30	20.90	0	0	0.7	4.0	0.0	0.0	0.1	0.6	4.7
5	7	C	30	20.90	0	14	0.7	4.8	0.0	0.0	0.0	2.2	7.0
5	7	C	30	20.90	0	21	1.0	6.9	0.0	0.0	0.5	1.0	7.9
5	7	C	30	20.90	80	0	0.1	4.1	0.0	0.0	5.0	38.2	42.3
5	7	C	30	20.90	80	14	0.3	7.0	0.0	0.0	2.1	21.9	29.0
5	7	C	30	20.90	80	21	1.9	6.4	0.0	0.0	1.3	30.4	36.9

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		SOL NH4		Ex NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	RANGE	AVER	
6	1	C	4	9.12	0	0	0.4	10.4	0.0	0.0	0.0	0.3	0.1	3.6	14.3
6	1	C	4	9.12	0	5	0.2	11.4	0.0	0.0	0.7	1.3	0.9	4.1	16.7
6	1	C	4	9.12	0	14	0.6	10.9	0.0	0.0	0.0	0.2	1.1	5.6	16.7
6	1	C	4	9.12	0	21	2.1	10.8	0.0	0.0	0.9	0.4	0.7	4.3	15.5
6	1	C	4	9.12	80	0	0.2	11.4	0.0	0.0	1.2	17.2	4.9	40.1	68.7
6	1	C	4	9.12	80	5	0.4	10.2	0.0	0.0	4.3	20.8	5.3	43.5	74.5
6	1	C	4	9.12	80	14	0.4	10.7	0.5	0.3	3.5	16.3	1.1	43.6	70.8
6	1	C	4	9.12	80	21	0.7	10.8	0.0	0.0	2.2	13.9	3.9	41.1	65.9
6	1	C	4	22.80	0	0	0.1	13.4	0.5	0.3	0.6	0.5	0.3	2.5	16.7
6	1	C	4	22.80	0	5	0.3	18.4	0.5	0.8	0.1	1.4	1.3	3.1	23.8
6	1	C	4	22.80	0	14	0.3	20.6	0.0	0.0	0.0	0.0	0.5	1.8	22.4
6	1	C	4	22.80	0	21	1.9	23.3	0.0	0.0	0.1	0.1	0.1	1.8	25.1
6	1	C	4	22.80	80	0	0.2	14.6	1.0	1.5	2.0	22.1	1.0	44.6	82.8
6	1	C	4	22.80	80	5	0.2	16.8	0.0	2.0	1.3	19.3	4.1	40.3	78.5
6	1	C	4	22.60	80	14	0.5	22.8	0.0	0.0	3.7	18.8	5.0	41.8	63.4
6	1	C	4	22.80	80	21	1.0	27.2	0.0	0.5	4.2	18.1	4.2	44.6	90.4
6	1	C	15	9.12	0	0	0.4	10.4	0.0	0.0	0.0	0.3	0.1	3.6	14.3
6	1	C	15	9.12	0	5	0.6	10.8	0.0	0.0	0.6	1.2	0.1	4.9	16.9
6	1	C	15	9.12	0	14	0.6	10.5	0.0	0.0	0.1	0.1	0.0	4.8	15.4
6	1	C	15	9.12	0	21	0.2	10.9	0.0	0.0	0.1	0.1	1.7	4.6	15.9
6	1	C	15	9.12	80	0	0.2	11.4	0.0	0.0	1.2	17.2	4.9	40.1	68.7
6	1	C	15	9.12	80	5	0.4	10.3	0.0	0.0	4.0	22.6	5.2	49.9	82.8
6	1	C	15	9.12	80	14	0.0	11.8	0.0	0.0	0.0	16.6	0.0	49.7	78.1
6	1	C	15	9.12	80	21	1.0	12.4	0.0	0.0	0.5	14.3	0.5	38.9	65.6
6	1	C	15	22.80	0	0	0.1	13.4	0.5	0.3	0.6	0.5	0.3	2.5	16.7
6	1	C	15	22.80	0	5	0.5	20.3	0.0	0.0	0.4	0.7	1.1	2.9	23.9
6	1	C	15	22.80	0	14	0.1	19.4	0.0	0.0	0.0	0.0	0.0	1.2	20.6
6	1	C	15	22.80	0	21	0.1	21.6	0.0	0.0	0.2	0.1	0.4	1.5	23.1
6	1	C	15	22.80	80	0	0.2	14.6	1.0	1.5	2.0	22.1	1.0	44.6	82.8
6	1	C	15	22.80	80	5	0.9	26.1	1.5	0.8	1.2	18.1	2.7	42.8	87.6
6	1	C	15	22.80	80	14	11.0	52.1	0.0	0.0	7.3	7.3	15.6	20.6	88.0
6	1	C	15	22.80	80	21	1.4	63.4	0.0	0.0	0.4	3.5	1.9	17.8	84.7
6	1	C	30	9.12	0	0	0.4	10.4	0.0	0.0	0.0	0.3	0.1	3.6	14.3
6	1	C	30	9.12	0	5	0.0	11.3	3.0	1.5	0.2	1.3	1.0	5.7	19.8
6	1	C	30	9.12	0	14	3.1	6.1	0.0	0.0	0.7	0.5	1.6	4.1	10.7
6	1	C	30	9.12	0	21	0.1	10.8	0.0	0.0	0.2	0.3	2.0	1.7	12.8
6	1	C	30	9.12	80	0	0.2	11.4	0.0	0.0	1.2	17.2	4.9	40.1	68.7
6	1	C	30	9.12	80	5	0.8	11.7	0.0	0.0	3.7	19.5	7.8	49.0	80.2
6	1	C	30	9.12	80	14	0.6	10.5	4.5	2.3	1.2	11.4	4.5	36.4	60.6
6	1	C	30	9.12	80	21	0.5	11.8	0.0	0.0	1.3	13.8	0.8	38.6	64.2
6	1	C	30	22.80	0	0	0.1	13.4	0.5	0.3	0.6	0.5	0.3	2.5	16.7
6	1	C	30	22.80	0	5	0.9	21.3	0.0	0.0	0.5	1.6	0.4	4.4	27.4
6	1	C	30	22.80	0	14	0.0	23.7	0.0	0.0	0.0	0.1	0.1	1.1	24.9
6	1	C	30	22.80	0	21	1.4	27.6	0.0	0.0	0.1	0.1	0.4	1.6	29.3
6	1	C	30	22.80	80	0	0.2	14.6	1.0	1.5	2.0	22.1	1.0	44.6	82.8
6	1	C	30	22.80	80	5	0.1	48.6	0.0	0.0	0.1	11.8	1.2	35.6	96.1
6	1	C	30	22.80	80	14	7.5	94.6	0.0	0.0	0.0	0.0	0.1	1.3	95.8
6	1	C	30	22.80	80	21	17.7	108.1	0.0	0.0	0.0	0.2	3.2	2.5	110.6

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	
N5	1		4	9.60	0	0	0.9	37.6	0.0	0.0	0.1	5.3	43.0
N5	1		4	9.60	0	14	0.2	34.8	0.0	0.0	0.0	7.3	42.1
N5	1		4	9.60	0	21	3.2	37.1	0.0	0.0	0.3	6.6	43.7
N5	1		4	9.60	80	0	1.7	51.4	0.0	0.0	0.6	63.4	118.8
N5	1		4	9.60	80	14	3.8	48.6	0.0	0.0	3.2	66.4	115.0
N5	1		4	9.60	80	21	4.1	51.6	0.0	0.0	0.3	64.9	116.5
N5	1		4	24.00	0	0	0.4	41.8	0.0	0.0	0.1	12.8	54.6
N5	1		4	24.00	0	14	1.1	51.9	0.0	0.0	0.8	17.6	69.7
N5	1		4	24.00	0	21	2.3	64.4	0.5	0.3	0.8	13.2	77.9
N5	1		4	24.00	80	0	8.5	57.6	0.0	0.0	11.6	80.6	138.2
N5	1		4	24.00	80	14	4.2	63.8	0.0	0.0	0.6	80.4	144.2
N5	1		4	24.00	80	21	32.2	70.0	0.5	0.3	5.3	71.5	141.8
N5	1		15	9.60	0	0	0.9	37.6	0.0	0.0	0.1	5.3	43.0
N5	1		15	9.60	0	14	1.4	36.9	0.0	0.0	0.1	6.8	43.7
N5	1		15	9.60	0	21	0.2	39.5	0.0	0.0	0.5	6.6	46.0
N5	1		15	9.60	80	0	1.7	51.4	0.0	0.0	0.6	63.4	114.8
N5	1		15	9.60	80	14	25.2	65.3	0.0	0.0	36.5	84.8	150.0

N5	1	15	9.60	80	21	4.6	59.6	0.0	0.0	0.0	61.9	121.5
N5	1	15	24.00	0	0	0.4	41.6	0.0	0.0	0.1	12.8	54.6
N5	1	15	24.00	0	14	0.0	84.8	0.0	0.0	0.3	1.0	85.8
N5	1	15	24.00	0	21	0.0	88.9	0.0	0.0	0.2	0.6	89.5
N5	1	15	24.00	80	0	8.5	57.6	0.0	0.0	11.6	80.6	138.2
N5	1	15	24.00	80	14	11.0	157.5	0.0	0.0	10.1	10.3	167.8
N5	1	15	24.00	80	21	1.0	172.5	0.0	0.0	0.3	1.3	173.8

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	
N5	1		30	9.60	0	0	0.9	37.6	0.0	0.0	0.1	5.3	43.0
N5	1		30	9.60	0	14	2.5	37.1	0.0	0.0	0.4	8.7	45.8
N5	1		30	9.60	0	21	3.7	39.6	0.0	0.0	0.7	8.6	48.3
N5	1		30	9.60	80	0	1.7	51.4	0.0	0.0	0.6	63.4	114.8
N5	1		30	9.60	80	14	2.0	49.9	0.0	0.0	2.0	64.7	114.6
N5	1		30	9.60	80	21	3.6	53.8	0.0	0.0	0.3	63.5	117.3
N5	1		30	24.00	0	0	0.4	41.8	0.0	0.0	0.1	12.8	54.6
N5	1		30	24.00	0	14	1.0	101.5	0.0	0.0	0.0	1.4	102.9
N5	1		30	24.00	0	21	4.0	103.0	0.0	0.0	0.1	0.5	103.5
N5	1		30	24.00	80	0	8.5	57.6	0.0	0.0	11.6	80.6	138.2
N5	1		30	24.00	80	14	11.0	182.5	0.0	0.0	0.0	1.5	184.0
N5	1		30	24.00	80	21	7.0	199.5	0.0	0.0	0.3	1.6	201.1
N5	2		4	8.10	0	0	3.1	34.6	0.0	0.0	0.3	4.8	39.5
N5	2		4	8.10	0	14	1.8	35.4	0.0	0.0	0.1	3.8	39.2
N5	2		4	6.10	0	21	1.3	34.8	0.0	0.0	0.8	3.3	38.1
N5	2		4	8.10	80	0	1.0	41.4	2.0	1.0	8.5	82.6	125.0
N5	2		4	8.10	80	14	13.0	40.1	0.0	0.0	1.3	78.0	118.1
N5	2		4	8.10	80	21	7.9	37.8	0.0	0.0	1.3	82.8	120.7
N5	2		4	20.30	0	0	0.9	36.6	0.5	0.8	0.4	7.7	45.0
N5	2		4	20.30	0	14	1.5	47.8	0.1	2.1	0.4	5.1	54.9
N5	2		4	20.30	0	21	1.7	49.8	0.5	0.8	0.1	2.8	53.3
N5	2		4	20.30	80	0	12.4	42.7	2.0	1.0	8.6	86.3	130.0
N5	2		4	20.30	80	14	4.7	59.8	2.0	4.0	8.7	96.6	160.5
N5	2		4	20.30	80	21	1.9	57.6	0.5	4.8	1.0	75.3	137.7
N5	2		15	6.10	0	0	3.1	34.6	0.0	0.0	0.3	4.8	39.5
N5	2		15	8.10	0	14	1.2	37.7	0.0	0.0	0.1	0.6	38.3
N5	2		15	8.10	0	21	1.6	36.4	0.0	0.0	0.2	0.8	37.2
N5	2		15	8.10	80	0	1.0	41.4	2.0	1.0	8.5	82.6	125.0
N5	2		15	8.10	80	14	11.1	41.1	0.0	0.0	3.1	76.1	117.3
N5	2		15	8.10	80	21	3.3	50.8	0.0	0.0	4.1	64.8	115.6
N5	2		15	20.30	0	0	0.9	36.6	0.5	0.8	0.4	7.7	45.0
N5	2		15	20.30	0	14	0.0	57.3	0.0	0.0	0.0	0.6	57.9
N5	2		15	20.30	0	21	0.0	61.3	0.0	0.0	0.0	0.5	61.8
N5	2		15	20.30	80	0	12.4	42.7	2.0	1.0	8.6	86.3	130.0
N5	2		15	20.30	80	14	21.0	128.5	0.0	0.0	2.8	3.5	132.0
N5	2		15	20.30	80	21	22.0	129.0	0.0	0.0	0.4	2.3	131.3
N5	2		30	8.10	0	0	3.1	34.6	0.0	0.0	0.3	4.8	39.5
N5	2		30	8.10	0	14	0.6	40.3	0.0	0.0	0.0	0.5	40.8
N5	2		30	8.10	0	21	1.1	43.3	0.0	0.0	0.1	0.5	43.9
N5	2		30	8.10	80	0	1.0	41.4	2.0	1.0	8.5	82.6	125.0
N5	2		30	8.10	80	14	13.4	44.9	0.0	0.0	0.0	64.4	109.3
N5	2		30	8.10	80	21	0.0	47.6	0.0	0.0	4.4	48.6	96.2
N5	2		30	20.30	0	0	0.9	36.6	0.5	0.8	0.4	7.7	45.0
N5	2		30	20.30	0	14	0.0	70.0	0.0	0.0	0.1	0.5	70.5
N5	2		30	20.30	0	21	1.8	73.4	0.0	0.0	0.0	0.5	73.9
N5	2		30	20.30	80	0	12.4	42.7	2.0	11.0	8.6	86.3	130.0
N5	2		30	20.30	80	14	5.0	137.5	0.0	0.0	0.4	2.3	139.8
N5	2		30	20.30	80	21	3.0	139.5	0.0	0.0	0.4	2.3	141.8
N5	3		4	9.00	0	0	2.0	60.6	0.0	0.0	1.2	6.6	67.2
N5	3		4	9.00	0	14	6.1	63.9	0.0	0.0	0.3	6.3	70.3
N5	3		4	9.00	0	21	1.1	61.6	0.0	0.0	1.1	8.6	70.2
N5	3		4	9.00	80	0	12.9	67.1	0.0	0.0	6.5	89.9	157.1
N5	3		4	9.00	80	14	7.3	69.1	0.0	0.0	2.7	83.3	152.5
N5	3		4	9.00	80	21	0.8	69.6	0.0	0.0	1.7	85.8	155.4
N5	3		4	22.50	0	0	0.3	64.4	0.5	0.3	0.1	10.5	75.2
N5	3		4	22.50	0	14	2.7	79.9	0.0	0.0	0.6	10.4	90.3
N5	3		4	22.50	0	21	3.5	83.9	0.5	0.3	0.0	8.8	92.9
N5	3		4	22.50	80	0	0.8	69.9	0.5	0.8	3.5	95.9	166.6
N5	3		4	22.50	80	14	2.4	89.6	0.0	0.0	4.9	87.8	177.4
N5	3		4	22.50	80	21	4.1	88.1	0.0	0.0	24.4	95.8	183.9
N5	3		15	9.00	0	0	2.0	60.6	0.0	0.0	1.2	6.6	67.2
N5	3		15	9.00	0	14	0.0	67.3	0.0	0.0	0.8	1.9	69.2
N5	3		15	9.00	0	21	2.1	70.1	0.0	0.0	0.6	2.6	72.6
N5	3		15	9.00	80	0	12.9	67.1	0.0	0.0	6.5	89.9	157.1
N5	3		15	9.00	80	14	9.7	68.5	0.0	0.0	3.4	76.2	144.7
N5	3		15	9.00	80	21	11.9	68.1	0.0	0.0	42.3	104.6	172.9
N5	3		15	22.50	0	0	0.3	64.4	0.5	0.3	0.1	10.5	75.2

N5	3	15	22.50	0	14	1.9	95.8	0.0	0.0	0.3	0.8	96.5
N5	3	15	22.50	0	21	32.2	104.9	0.0	0.0	0.6	2.2	107.1
N5	3	15	22.50	80	0	0.8	69.9	0.5	0.8	3.5	95.9	166.6
N5	3	15	22.50	80	14	47.0	167.5	0.0	0.0	26.5	15.8	183.2
N5	3	15	22.50	80	21	6.0	185.0	0.0	0.0	3.6	3.8	188.8

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	
N5	3		30	9.00	0	0	2.0	60.6	0.0	0.0	1.2	6.6	67.2
N5	3		30	9.00	0	14	1.8	75.9	0.0	0.0	0.1	0.5	76.4
N5	3		30	9.00	0	21	6.0	77.7	0.0	0.0	0.3	2.3	80.0
N5	3		30	9.00	80	0	12.9	67.1	0.0	0.0	6.5	89.9	157.1
N5	3		30	9.00	80	14	9.9	88.6	0.0	0.0	2.7	67.6	156.3
N5	3		30	9.00	80	21	0.1	98.3	0.0	0.0	7.5	67.3	165.5
N5	3		30	22.50	0	0	0.3	64.4	0.5	0.3	0.1	10.5	75.2
N5	3		30	22.50	0	14	2.0	128.0	0.0	0.0	0.2	0.7	128.7
N5	3		30	22.50	0	21	6.0	135.0	0.0	0.0	0.8	2.4	137.4
N5	3		30	22.50	80	0	0.8	69.9	0.5	0.8	3.5	95.9	166.6
N5	3		30	22.50	80	14	2.0	185.0	0.0	0.0	0.7	2.1	187.1
N5	3		30	22.50	80	21	4.0	203.0	0.0	0.0	1.7	4.0	207.0
N5	4		4	9.50	0	0	1.9	44.1	0.0	0.0	0.2	5.9	49.9
N5	4		4	9.50	0	14	5.5	49.1	0.0	0.0	0.1	6.3	55.4
N5	4		4	9.50	0	21	0.0	50.2	0.0	0.0	0.9	8.3	58.4
N5	4		4	9.50	80	0	3.5	40.1	0.0	0.0	9.0	77.6	117.7
N5	4		4	9.50	80	14	11.2	49.9	0.0	0.0	12.8	75.4	125.3
N5	4		4	9.50	80	21	0.0	53.2	0.0	0.0	3.6	80.6	133.3
N5	4		4	23.90	0	0	0.3	46.8	0.0	0.0	0.3	8.1	54.9
N5	4		4	23.90	0	14	1.5	54.4	0.0	0.0	0.2	9.3	63.7
N5	4		4	23.90	0	21	1.6	58.6	0.0	0.0	1.0	10.5	69.1
N5	4		4	23.90	80	0	0.2	37.2	0.0	0.0	39.8	87.1	124.3
N5	4		4	23.90	80	14	5.1	58.9	0.0	0.0	7.9	61.6	120.5
N5	4		4	23.90	80	21	1.6	63.1	0.0	0.0	12.7	73.5	136.6
N5	4		15	9.50	0	0	1.9	44.1	0.0	0.0	0.2	5.9	49.9
N5	4		15	9.50	0	14	1.4	52.5	0.0	0.0	0.1	5.1	57.8
N5	4		15	9.50	0	21	3.6	52.0	0.0	0.0	0.7	7.1	59.1
N5	4		15	9.50	80	0	3.5	40.1	0.0	0.0	9.0	77.6	117.7
N5	4		15	9.50	80	14	1.0	63.5	0.0	0.0	11.8	82.8	146.3
N5	4		15	9.50	80	21	7.0	58.7	0.0	0.0	24.3	83.6	142.5
N5	4		15	23.90	0	0	0.3	46.8	0.0	0.0	0.3	8.1	54.9
N5	4		15	23.90	0	14	0.1	69.3	0.0	0.0	0.2	0.7	69.9
N5	4		15	23.90	0	21	6.1	68.1	0.0	0.0	0.6	2.2	70.3
N5	4		15	23.90	80	0	0.2	37.2	0.0	0.0	39.8	67.1	124.3
N5	4		15	23.90	80	14	17.5	108.3	0.0	0.0	16.9	40.1	148.3
N5	4		15	23.90	80	21	1.0	119.5	0.0	0.0	0.1	4.8	124.3
N5	4		30	9.50	0	0	1.9	44.1	0.0	0.0	0.2	5.9	49.9
N5	4		30	9.50	0	14	14.4	42.2	0.0	0.0	1.0	4.5	46.7
N5	4		30	9.50	0	21	1.0	42.7	0.0	0.0	0.6	7.6	50.3
N5	4		30	9.50	80	0	3.5	40.1	0.0	0.0	9.0	77.6	117.7
N5	4		30	9.50	80	14	10.3	60.3	0.0	0.0	22.7	71.3	131.3
N5	4		30	9.50	80	21	1.6	64.3	0.0	0.0	11.7	60.1	124.3
N5	4		30	23.90	0	0	0.3	46.8	0.0	0.0	0.3	8.1	54.9
N5	4		30	23.90	0	14	0.8	69.3	0.0	0.0	0.1	0.8	70.0
N5	4		30	23.90	0	21	4.1	78.1	0.0	0.0	0.5	2.4	80.5
N5	4		30	23.90	80	0	0.2	37.2	0.0	0.0	39.8	87.1	124.3
N5	4		30	23.90	80	14	9.0	132.5	0.0	0.0	10.1	7.9	140.3
N5	4		30	23.90	80	21	52.0	154.0	0.0	0.0	3.8	2.3	156.5
N5	5		4	7.00	0	0	2.9	26.3	0.0	0.0	0.0	4.5	30.3
N5	5		4	7.00	0	14	1.0	26.8	0.0	0.0	0.7	7.5	36.3
N5	5		4	7.00	0	21	0.2	28.6	0.0	0.0	0.5	9.3	37.8
N5	5		4	7.00	80	0	2.0	53.6	0.0	0.0	0.3	78.8	132.4
N5	5		4	7.00	80	14	11.2	47.5	0.0	0.0	4.1	73.1	120.5
N5	5		4	7.00	80	21	0.0	50.1	0.0	0.0	0.0	77.5	127.5
N5	5		4	17.60	0	0	0.1	30.3	0.0	0.0	0.0	6.9	37.1
N5	5		4	17.60	0	14	1.0	35.8	0.0	0.0	0.5	14.3	50.1
N5	5		4	17.60	0	21	1.5	32.6	0.0	0.0	0.4	18.3	50.9
N5	5		u	17.60	80	0	0.7	50.6	0.0	0.0	5.3	67.1	117.8
N5	5		4	17.60	80	14	5.8	51.5	0.0	0.0	3.1	77.1	128.5
N5	5		4	17.60	80	21	2.3	42.8	0.0	0.0	19.9	91.1	133.3
N5	5		15	7.00	0	0	2.9	26.3	0.0	0.0	0.0	4.5	30.3
N5	5		15	7.00	0	14	1.8	30.1	0.0	0.0	0.3	8.0	38.1
N5	5		15	7.00	0	21	0.6	26.1	0.0	0.0	0.6	9.8	35.9
N5	5		15	7.00	80	0	2.0	53.6	0.0	0.0	0.3	78.8	132.4
N5	5		15	7.00	80	14	2.3	35.8	0.0	0.0	1.6	74.6	110.4
N5	5		15	7.00	80	21	0.3	32.3	0.0	0.0	15.5	66.4	98.8

Table 3 cont'd.

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER	
N5	5		15	17.60	0	0	0.1	30.3	0.0	0.0	0.0	6.9	37.1
N5	5		15	17.60	0	14	3.3	42.3	0.0	0.0	0.8	19.8	62.1
N5	5		15	17.60	0	21	2.7	44.5	0.0	0.0	2.3	21.1	65.7
N5	5		15	17.60	80	0	0.7	50.6	0.0	0.0	5.3	67.1	117.8
N5	5		15	17.60	80	14	0.5	48.6	0.0	0.0	7.5	79.3	127.9
N5	5		15	17.60	80	21	6.0	46.3	0.0	0.0	4.7	78.1	124.4
N5	5		30	7.00	0	0	2.9	26.3	0.0	0.0	0.0	4.5	30.8
N5	5		30	7.00	0	14	6.2	31.5	0.0	0.0	4.1	15.3	46.6
N5	5		30	7.00	0	21	1.3	28.4	0.0	0.0	0.7	15.1	43.6
N5	5		30	7.00	80	0	2.0	53.6	0.0	0.0	0.3	78.8	132.6
N5	5		30	7.00	80	14	4.4	38.6	0.0	0.0	13.3	94.3	132.8
N5	5		30	7.00	80	21	25.8	49.4	0.0	0.0	5.6	87.7	137.1
N5	5		30	17.60	0	0	0.1	30.3	0.0	0.0	0.0	6.9	37.6
N5	5		30	17.60	0	14	3.5	51.1	0.0	0.0	1.1	17.3	68.5
N5	5		30	17.60	0	21	10.8	51.5	0.0	0.0	1.1	12.8	64.8
N5	5		30	17.60	80	0	0.7	50.6	0.0	0.0	5.3	67.1	117.6
N5	5		30	17.60	80	14	5.0	50.7	0.0	0.0	10.1	86.5	137.5
N5	5		30	17.60	80	21	14.9	73.6	0.0	0.0	1.4	94.5	168.1

Table 4 Recovery of added $\text{NH}_4^+\text{-N}$ ($80 \mu\text{g N/g}$) from soil profiles representing watersheds #1, 5 and 13.

Depth (cm)	% Recovery of original $\text{NH}_4^+\text{-N}$ applied						
	Sandy loam	Clay (5)	N-5/1	N-5/2	N-5/3	N-5/4	N-5/5
0-10	94	89	79	98	106	98	84
10-20	98	89					
20-30	98	68					
30-45	103	52					
45-60	102	44					
60-75	108	33					
75-90	106	49					

Table 5. Rate coefficients of nitrification of added $\text{NH}_4^+\text{-N}$ in soils from 3 watersheds

PL	DP	SOL	TMP	WC	TRT	RN-14	RN-21	RN-AVE	PL	DP	SOL	TMP	WC	TRT	RN-14	RN-21	RN-AVE
1	1	S	4	2.87	80	-0.006	-0.004	-0.005	3	1	S	4	2.87	80	-0.001	-0.009	-0.005
1	t	S	4	7.19	80	-0.001	-0.002	-0.001	3	1	S	4	7.17	80	0.002	-0.002	-0.000
1	1	S	15	2.87	80	-0.003	-0.002	-0.003	3	1	S	15	2.87	80	-0.002	-0.002	-0.002
1	1	S	15	7.19	80	-0.002	0.005	0.001	3	1	S	15	7.17	80	0.012	0.014	0.013
1	1	S	30	2.87	80	0.000	0.004	0.002	3	1	S	30	2.87	80	-0.003	-0.003	-0.003
1	1	S	30	7.19	80	0.018	0.035	0.027	3	1	S	30	7.17	80	0.132	0.040	0.086
1	2	S	4	2.60	80	0.010	0.008	0.009	3	2	S	4	3.06	80	0.000	0.002	0.001
1	2	S	4	6.50	80	0.014	0.009	0.011	3	2	S	4	7.65	80	0.004	0.002	0.003
1	2	S	15	2.60	80	0.008	0.005	0.007	3	2	S	15	3.06	80	0.001	0.001	0.001
1	2	S	15	6.50	80	0.016	0.009	0.014	3	2	S	15	7.65	80	0.031	0.036	0.033
1	2	S	30	2.60	80	0.008	0.009	0.006	3	2	S	30	3.06	80	-0.003	0.000	-0.001
1	2	S	30	6.50	80	0.015	0.009	0.012	3	2	S	30	7.65	80	0.054	0.036	0.045
2	1	S	4	3.45	80	-0.001	-0.001	-0.001	4	1	C	4	9.60	80	0.002	0.002	0.002
2	1	S	4	8.63	80	0.005	-0.002	0.001	4	1	C	4	24.00	80	0.010	0.006	0.008
2	1	S	15	3.45	80	-0.000	0.000	0.000	4	1	C	15	9.60	80	0.001	0.003	0.002
2	1	S	15	6.63	80	0.015	0.016	0.016	4	1	C	15	24.00	80	0.060	0.072	0.066
2	1	S	30	3.45	80	-0.002	-0.001	-0.001	4	1	C	30	9.60	80	0.005	0.004	0.005
2	1	S	30	8.63	80	0.041	0.024	0.052	4	1	C	30	24.00	80	0.307	0.138	0.222
2	2	S	4	3.42	80	-0.001	0.000	-0.000	4	2	C	4	9.06	80	0.003	0.002	0.002
2	2	S	4	8.55	80	-0.002	0.000	-0.001	4	2	C	4	24.20	80	-0.005	0.000	-0.002
2	2	S	15	3.42	80	0.001	0.002	0.001	4	2	C	15	9.66	80	0.010	-0.000	0.005
2	2	S	15	8.55	80	0.005	0.003	0.004	4	2	C	15	24.20	80	0.036	0.079	0.057
2	2	S	30	3.42	80	0.006	0.003	0.005	4	2	C	30	9.06	80	0.013	0.010	0.011
2	2	S	30	8.55	80	0.011	0.015	0.013	4	2	C	30	24.20	80	0.205	0.185	0.195
2	1	SB	4	2.94	80	-0.005	-0.003	-0.004									
2	1	SB	4	7.34	80	-0.002	0.000	-0.001									
2	1	SB	15	2.94	80	-0.006	-0.003	-0.004									
2	1	SB	15	7.34	80	-0.003	-0.001	-0.002									
2	1	SB	30	2.94	80	-0.008	-0.000	-0.004									
2	1	SB	30	7.34	80	-0.003	-0.000	-0.002									
2	2	SB	4	2.91	80	0.001	0.002	0.001									
2	2	SB	4	7.28	80	-0.001	0.000	-0.000									
2	2	SB	15	2.91	80	0.002	-0.000	0.001									
2	2	SB	15	7.28	80	0.002	0.005	0.003									
2	2	SB	30	2.91	80	0.002	0.003	0.003									
2	2	SB	30	7.28	80	0.011	0.010	0.011									
2	3	S	4	2.81	80	-0.002	-0.002	-0.002									
2	3	S	4	7.04	80	0.004	0.004	0.004									
2	3	S	15	2.81	80	0.004	-0.000	0.002									
2	3	S	15	7.04	80	0.017	0.007	0.012									
2	3	S	30	2.81	80	0.001	0.001	0.001									
2	3	S	30	7.04	80	0.033	0.028	0.031									
2	4	S	4	2.26	80	-0.004	-0.001	-0.002									
2	4	S	4	5.59	80	0.002	-0.001	0.001									
2	4	S	15	2.26	80	-0.003	-0.001	-0.002									
2	4	S	15	5.59	80	0.001	0.002	0.002									
2	4	S	30	2.26	80	-0.000	0.001	0.000									
2	4	S	30	5.59	80	0.010	0.005	0.007									
2	5	S	4	1.44	80	0.001	0.001	0.001									
2	5	S	4	3.61	80	0.002	0.002	0.002									
2	5	S	15	1.44	80	-0.002	0.001	-0.000									
2	5	S	15	3.61	80	0.001	-0.001	0.000									
2	5	S	30	1.44	80	0.002	0.001	0.102									
2	5	S	30	3.61	80	0.001	0.001	0.001									
2	6	S	4	1.26	80	0.001	0.001	0.001									
2	6	S	4	3.14	80	0.000	0.001	0.000									
2	6	S	15	1.26	80	-0.000	0.000	-0.000									
2	6	S	15	3.14	80	0.002	0.001	0.002									
2	6	S	30	1.26	80	0.001	0.000	0.000									
2	6	S	30	3.14	80	0.003	0.000	0.002									
2	7	S	4	1.22	80	-0.001	-0.001	-0.001									
2	7	S	4	3.05	60	0.002	-0.001	0.000									
2	7	S	15	1.22	80	0.001	0.001	0.001									
2	7	S	15	3.05	80	0.002	-0.003	-0.000									
2	7	S	30	1.22	80	0.005	0.000	0.003									
2	7	S	30	3.05	80	-0.001	-0.002	-0.001									

Table 5 cont.

PL	DP	SOL	TMP	WC	TRT	RN-14	RN-21	RN-AVE	PL	DP	SOL	TMP	WC	TRT	RN-14	RN-21	RN-AVE
5	1	C	4	8.76	80	-0.002	0.004	0.001	6	l	C	4	9.12	80	-0.001	0.001	-0.000
5	1	C	4	21.90	80	0.000	-0.002	-0.001	6	1	C	4	22.80	80	0.003	0.002	0.002
5	1	C	15	8.76	80	-0.005	-0.003	-0.004	6	1	C	15	9.12	80	-0.005	0.003	-0.001
5	1	C	15	21.90	80	0.046	0.008	0.027	6	t	C	15	22.80	80	0.045	0.051	0.048
5	1	C	30	8.76	80	0.001	-0.006	-0.002	6	1	C	30	9.12	80	0.012	0.001	0.006
5	1	C	30	21.90	80	0.119	0.064	0.092	6	1	C	30	22.80	80	*****	*****	*****
5	2	C	4	8.70	80	0.010	-0.002	0.004									
5	2	C	4	21.80	80	0.005	0.003	0.004	N5	1		4	9.60	80	-0.001	0.000	-0.000
5	2	C	15	8.70	80	-0.004	-0.001	-0.002	N5	1		4	24.00	80	0.001	-0.000	0.000
5	2	C	15	21.80	80	0.135	0.127	0.131	N5	1		15	9.60	80	-0.003	0.004	0.000
5	2	C	30	8.70	80	0.003	0.006	0.004	N5	1		15	24.00	80	0.136	0.251	0.193
5	2	C	30	21.80	80	0.198	0.119	0.158	N5	1		30	9.60	80	0.001	0.001	0.001
5	1	CB	4	9.24	80	0.002	-0.002	-0.000	N5	1		30	24.00	80	0.280	*****	*****
5	1	CB	4	23.10	80	0.019	0.012	0.018	N5	2		4	8.10	80	0.000	-0.002	-0.001
5	1	CB	15	9.24	80	0.003	-0.002	0.000	N5	2		4	20.10	80	-0.002	0.004	0.001
5	1	CB	15	23.10	80	0.176	0.106	0.141	N5	2		15	8.10	80	-0.001	0.007	0.003
5	1	CB	30	9.24	80	0.005	0.009	0.007	N5	2		15	20.30	80	0.160	0.100	0.130
5	1	CB	30	23.10	80	0.231	0.099	0.165	N5	2		30	8.10	80	0.005	0.009	0.007
5	2	CB	4	9.69	80	0.001	0.005	0.003	N5	2		30	20.30	80	0.150	0.097	0.123
5	2	CB	4	24.20	80	0.006	-0.000	0.003	N5	3		4	9.00	80	0.002	0.002	0.002
5	2	CB	15	9.69	80	-0.013	-0.001	-0.007	N5	3		4	22.50	80	0.005	-0.001	0.002
5	2	CB	15	24.20	80	0.041	0.046	0.043	N5	3		15	9.00	80	0.002	-0.007	-0.003
5	2	CB	30	9.69	80	-0.007	0.002	-0.002	N5	3		15	22.50	80	0.114	0.123	0.119
5	2	CB	30	24.20	80	0.053	0.049	0.051	N5	3		30	9.00	80	0.010	0.010	0.010
5	3	C	4	8.82	80	0.010	0.008	0.009	N5	3		30	22.50	80	0.112	0.092	0.102
5	3	C	4	22.10	60	0.001	-0.006	-0.002	N5	4		4	9.50	80	0.004	0.002	0.003
5	3	C	15	8.82	80	0.000	0.010	0.008	N5	4		4	23.90	80	0.021	0.010	0.016
5	3	C	15	22.10	80	0.042	0.056	0.049	N5	4		15	9.50	80	0.005	0.002	0.003
5	3	C	30	8.82	80	0.021	0.002	0.012	N5	4		15	23.90	80	0.058	0.097	0.078
5	3	C	30	22.10	80	0.164	*****	*****	N5	4		30	9.50	80	0.015	0.018	0.016
5	4	C	4	8.40	80	0.006	0.003	0.005	N5	4		30	23.90	80	0.177	*****	*****
5	4	C	4	21.00	80	-0.003	-0.000	-0.002	N5	5		4	7.00	80	0.000	0.000	0.000
5	4	C	15	8.40	80	0.003	0.001	0.002	N5	5		4	17.80	80	-0.004	-0.008	-0.006
5	4	C	15	21.00	80	0.009	0.000	0.005	N5	5		15	7.00	80	-0.006	-0.031	-0.004
5	4	C	30	8.40	60	0.009	0.003	0.006	N5	5		15	17.60	80	-0.007	-0.006	-0.007
5	4	C	30	21.00	80	0.000	-0.004	-0.002	N5	5		30	7.00	80	-0.011	-0.001	-0.006
5	4	C	4	48.10	80	0.009	0.001	0.005	N5	5		30	17.60	80	-0.016	-0.007	-0.012
5	5	C	4	20.90	80	-0.008	-0.003	-0.005									
5	5	C	15	8.10	80	-0.004	0.007	0.002									
5	5	C	15	20.90	80	-0.006	0.003	-0.001									
5	5	C	30	8.30	80	-0.010	0.006	-0.002									
5	5	C	30	20.90	80	0.001	-0.010	-0.005									
5	6	C	4	8.20	80	0.000	-0.009	-0.004									
5	6	C	4	20.40	80	0.007	0.002	0.005									
5	8	C	15	8.20	80	-0.002	-0.004	-0.003									
5	6	C	15	20.40	80	0.004	0.007	0.006									
5	6	C	30	8.20	80	0.004	0.001	0.003									
5	6	C	30	20.40	80	0.010	0.006	0.008									
5	7	C	4	8.40	80	0.010	0.001	0.006									
5	7	C	4	20.90	80	0.007	0.006	0.007									
5	7	C	15	8.40	80	0.011	0.010	0.010									
5	7	C	15	20.90	80	0.016	0.009	0.012									
5	7	C	30	8.40	80	0.009	0.012	0.011									
5	7	C	30	20.90	80	0.022	0.005	0.014									

Table 6. Correlations of nitrification rate coefficients with temperature and moisture in surface samples of watershed soils

Factor	Correlation coefficients			Non preassigned % reduction in total sum of squares		
	Brookston Clay	Berrien Sandy Loam	N-5	Brookston Clay	Berrien Sandy Loam	N-5
Temp. x H ₂ O	0.82	0.61	0.73	68.0	37.4	53.6
Temp.	0.45	0.36	0.25	76.9	40.5	75.4
H ₂ O	<u>0.58</u>	<u>0.40</u>	<u>0.64</u>	77.3	40.8	75.4
Multiple	0.88	0.64	0.87			

Table 7 Weighted average daily rates of mineralization of watershed soils

PL	DP	SOL	TMP	WC	TRT	MINRATE	PL	DP	SOL	TMP	WC	TRT	MINRATE	PL	DP	SOL	TMP	WC	TRT	MINRATE
1	1	8	4	2.87	0	0.2171	3	1	S	4	2.87	0	0.0771	5	1	C	4	8.76	0	0.1971
1	1	9	4	7.19	0	0.1743	3	1	S	4	7.17	0	0.0400	5	1	C	4	21.90	0	0.6943
1	1	S	15	2.87	0	0.1914	3	1	S	15	2.87	0	0.0514	5	1	c	15	8.76	0	0.2371
1	1	S	15	7.19	0	0.0886	3	1	S	15	7.17	0	-0.0429	5	1	C	15	21.90	0	0.6686
1	1	S	30	2.87	0	-0.1029	3	1	S	30	2.87	0	0.1229	5	1	C	30	8.76	0	0.4171
1	1	S	30	7.19	0	0.3943	3	1	S	30	7.17	0	0.1686	5	1	C	30	21.90	0	1.1743
1	2	S	4	2.60	0	-0.4886	3	2	S	4	3.06	0	-0.0943	5	2	C	4	8.70	0	0.0800
1	2	S	4	6.50	0	0.0486	3	2	S	4	7.65	0	0.0229	5	2	C	4	21.80	0	0.4971
1	2	S	15	2.60	0	-0.5571	3	2	S	15	3.06	0	0.0314	5	2	C	15	8.70	0	-0.0457
1	2	S	15	6.50	0	0.0514	3	2	S	15	7.65	0	0.2171	5	2	C	15	21.80	0	0.8080
1	2	S	30	2.60	0	-0.4486	3	2	S	30	3.06	0	0.1400	5	2	C	30	8.70	0	0.1200
1	2	S	30	6.50	0	0.5343	3	2	S	30	7.65	0	0.4371	5	2	C	30	21.80	0	1.4429
														5	1	CB	4	9.24	0	0.2743
2	1	S	4	3.45	0	0.1914	4	1	C	4	9.60	0	0.0800	5	1	CB	4	23.10	0	0.4971
2	1	S	4	8.63	0	0.0543	4	1	C	4	24.00	0	0.7857	5	1	CB	15	9.24	0	0.4314
2	1	S	15	3.45	0	0.1857	4	1	C	15	9.60	0	-0.0143	5	1	CB	15	23.10	0	0.6971
2	1	S	15	6.63	0	-0.0286	4	1	C	15	24.00	0	0.8800	5	1	CB	30	9.24	0	0.2857
2	1	S	30	3.45	0	0.1200	4	1	C	30	9.60	0	-0.2171	5	1	CB	30	23.10	0	0.9429
2	1	S	30	8.63	0	0.0914	4	1	C	30	24.00	0	1.5486	5	2	CB	4	9.69	0	0.1343
2	2	S	4	3.42	0	0.1086	4	2	C	4	9.66	0	0.1086	5	2	CB	4	24.20	0	0.5371
2	2	S	4	8.55	0	0.1829	4	2	C	4	24.20	0	0.7514	5	2	CB	15	9.69	0	0.8743
2	2	S	15	3.42	0	-0.0914	4	2	C	15	9.66	0	0.0400	5	2	CB	15	24.20	0	0.9686
2	2	S	15	8.55	0	0.0371	4	2	C	15	24.20	0	1.0314	5	2	CB	30	9.69	0	0.3971
2	2	S	30	3.42	0	-0.0971	4	2	C	30	9.66	0	-0.1400	5	2	CB	30	24.20	0	2.0800
2	2	S	30	8.55	0	0.1571	4	2	C	30	24.20	0	2.0200	5	3	C	4	8.82	0	-0.0257
2	1	SB	4	2.94	0	0.2143								5	3	C	4	22.10	0	0.3171
2	1	SB	4	7.34	0	0.1457								5	3	C	15	6.82	0	-0.0200
2	1	SB	15	2.94	0	0.2657	N5	1		4	9.60	0	0.0	5	3	C	15	22.10	0	0.7600
2	1	SB	15	7.34	0	0.2914	N5	1		4	24.00	0	1.0971	5	3	C	30	8.82	0	0.0229
2	1	SB	30	2.94	0	0.1714	N5	1		15	9.60	0	0.1143	5	3	C	30	22.10	0	1.5600
2	1	SB	30	7.34	0	0.2829	N5	1		15	24.00	0	1.8686	5	4	C	4	8.40	0	0.0714
2	2	SB	4	2.91	0	-0.1029	N5	1		30	9.60	0	0.2343	5	4	C	4	21.00	0	0.3371
2	2	SB	4	7.28	0	0.0457	N5	1		30	24.00	0	2.7771	5	4	C	15	8.40	0	0.0657
2	2	SB	15	2.91	0	-0.0200	N5	2		4	8.10	0	-0.0429	5	4	C	15	21.00	0	0.5800
2	2	SB	15	7.28	0	0.2000	N5	2		4	20.30	0	0.5229	5	4	C	30	8.40	0	-0.0171
2	2	SB	30	2.91	0	-0.0343	N5	2		15	8.10	0	-0.0943	5	4	C	30	21.00	0	0.6029
2	2	SB	30	7.28	0	0.1057	N5	2		15	20.30	0	0.8486	5	5	C	4	8.30	0	-0.0286
2	3	S	4	2.81	0	-0.1200	N5	2		30	8.10	0	0.1657	5	5	C	4	20.90	0	0.0771
2	3	S	4	7.04	0	-0.0743	N5	2		30	20.30	0	1.5543	5	5	C	15	8.30	0	-0.0257
2	3	S	15	2.81	0	-0.1029	N5	3		4	9.00	0	0.1714	5	5	C	15	20.90	0	0.1857
2	3	S	15	7.04	0	-0.0343	N5	3		4	22.50	0	0.9371	5	5	C	30	8.30	0	-0.0114
2	3	S	30	2.81	0	-0.1057	N5	3		15	9.00	0	0.2143	5	5	C	30	20.90	0	0.1800
2	3	S	30	7.04	0	0.0486	N5	3		15	22.50	0	1.5229	5	6	C	4	8.20	0	0.0257
2	4	S	4	2.26	0	0.2400	N5	3		30	9.00	0	0.6286	5	6	C	4	20.40	0	0.0200
2	4	S	4	5.59	0	-0.0029	N5	3		30	22.50	0	3.3057	5	6	C	15	8.20	0	0.0229
2	4	S	15	2.26	0	0.1914	N5	4		4	9.50	0	0.3971	5	6	C	15	20.40	0	0.0943
2	4	S	15	5.59	0	-0.0029	N5	4		4	23.90	0	0.6571	5	6	C	30	8.20	0	0.0171
2	4	S	30	2.26	0	0.1543	N5	4		15	9.50	0	0.4771	5	6	C	30	20.40	0	0.1971
2	4	S	30	5.59	0	-0.1000	N5	4		15	23.90	0	0.8714	5	7	C	4	8.40	0	-0.0171
2	5	S	4	1.44	0	-0.0800	N5	4		30	9.50	0	-0.0657	5	7	C	4	20.90	0	0.0857
2	5	S	4	3.61	0	0.0371	N5	4		30	23.90	0	1.1657	5	7	C	15	6.40	0	-0.0171
2	5	S	15	1.44	0	-0.0371	N5	5		4	7.00	0	0.3600	5	7	C	15	20.90	0	0.1057
2	5	S	15	5.61	0	-0.0514	N5	5		4	17.60	0	0.7600	5	7	C	30	8.40	0	0.0286
2	5	S	30	1.44	0	-0.0086	N5	5		15	7.00	0	0.3543	5	7	C	30	20.90	0	0.1629
2	5	S	30	3.61	0	-0.0714	N5	5		15	17.60	0	1.5229							
2	6	S	4	1.26	0	0.0000	N5	5		30	7.00	0	0.8200							
2	6	S	4	3.14	0	0.0629	N5	5		30	17.60	0	1.6657	6	1	C	4	9.12	0	0.1029
2	6	S	15	1.26	0	0.0457								6	1	C	4	22.80	0	0.4057
2	6	S	19	5.14	0	0.0571								6	1	C	15	9.12	0	0.0771
2	6	S	30	1.26	0	-0.0286								6	1	C	15	22.80	0	0.2971
2	6	S	30	3.14	0	0.0686								6	1	C	30	9.12	0	-0.1457
2	7	S	4	1.22	0	0.0371								6	1	C	30	22.80	0	0.5943
2	7	S	4	5.05	0	0.0857														
2	7	S	15	1.22	0	0.0629														
2	7	S	15	3.05	0	0.0400														
2	7	S	30	1.22	0	0.0743														
2	7	S	30	3.05	0	0.1257														

PL	DP	SOL	TMP	WC	TRT	RD-2	RD-5	RD-AVE	PL	DP	SOL	TMP	WC	TRT	RD-2	RD-5	RD.AVE
2	1	S	4	27.00	0	0.020	-0.020	-0.000	5	1	C	4	50.00	0	-0.054	-0.052	-0.053
2	1	S	4	27.00	80	-0.010	-0.004	-0.007	5	1	C	4	50.00	80	-0.035	-0.019	-0.027
2	1	S	4	16.00	0	-0.023	-0.009	-0.016	5	1	C	4	66.00	0	-0.057	0.011	-0.023
2	1	S	4	36.00	80	0.034	0.012	0.023	5	1	C	4	66.00	80	-0.009	-0.006	-0.007
2	1	S	4	44.00	0	-0.044	-0.017	-0.030	5	1	C	4	79.00	0	-0.049	0.014	-0.017
2	1	S	4	44.00	80	0.017	0.007	0.012	5	1	C	4	79.00	80	0.016	0.007	0.012
2	1	S	15	27.00	0	-0.050	-0.037	-0.044	5	1	C	15	50.00	0	0.175	0.551	0.363
2	1	S	15	27.00	80	0.031	0.015	0.023	5	1	C	15	50.00	80	0.058	-0.011	0.024
2	1	S	15	36.00	0	0.106	0.179	0.142	5	1	C	15	66.00	0	0.342	0.560	0.451
2	1	S	15	36.00	80	0.008	0.011	0.009	5	1	C	15	66.00	80	0.074	0.028	0.051
2	1	S	15	44.00	0	0.198	0.166	0.182	5	1	C	15	79.00	0	0.678	0.761	0.720
2	1	S	15	44.00	80	0.063	0.033	0.048	5	1	C	15	79.00	80	0.096	0.054	0.075
2	1	S	30	27.00	0	0.503	0.401	0.452	5	1	C	30	50.00	0	*****	0.054	*****
2	1	S	30	27.00	80	0.083	0.025	0.054	5	1	C	30	50.00	80	0.125	0.006	0.065
2	1	S	30	36.00	0	1.980	0.572	1.276	5	1	C	30	66.00	0	*****	0.006	*****
2	1	S	30	36.00	80	0.171	0.067	0.119	5	1	C	30	66.00	80	0.301	0.134	0.218
2	1	S	30	44.00	0	*****	0.067	*****	5	1	C	30	79.00	0	*****	0.134	*****
2	1	S	30	44.00	80	0.144	0.065	0.105	5	1	C	30	79.00	80	0.280	0.116	0.198
2	2	S	4	27.00	0	-0.027	-0.026	-0.027	5	2	C	4	50.00	0	-0.523	-0.245	-0.384
2	2	S	4	27.00	80	0.005	0.007	0.006	5	2	C	4	50.00	80	-0.009	-0.015	-0.012
2	2	S	4	36.00	0	-0.027	-0.021	-0.024	5	2	C	4	67.00	0	-0.280	-0.164	-0.222
2	2	S	4	36.00	80	0.0	-0.011	-0.006	5	2	C	4	67.00	80	0.053	0.014	0.034
2	2	S	4	44.00	0	-0.026	-0.022	-0.025	5	2	C	4	80.00	0	-0.438	-0.190	-0.314
2	2	S	4	44.00	80	-0.058	-0.019	-0.039	5	2	C	4	80.00	80	0.019	-0.002	0.009
2	2	S	15	27.00	0	-0.014	-0.031	-0.022	5	2	C	15	50.00	0	-0.765	-0.102	-0.434
2	2	S	15	27.00	80	-0.043	-0.010	-0.027	5	2	C	15	50.00	80	-0.059	-0.019	-0.039
2	2	S	15	36.00	0	0.034	0.065	0.050	5	2	C	15	67.00	0	-0.570	-0.199	-0.384
2	2	S	15	36.00	80	-0.035	-0.014	-0.025	5	2	C	15	67.00	80	0.355	0.103	0.229
2	2	S	15	44.00	0	0.029	0.093	0.061	5	2	C	15	80.00	0	-0.541	-0.191	-0.366
2	2	S	15	44.00	80	-0.074	0.005	-0.035	5	2	C	15	80.00	80	0.144	0.069	0.107
2	2	S	30	27.00	0	0.182	-0.049	0.067	5	2	C	30	50.00	0	-0.363	0.484	0.061
2	2	S	30	27.00	80	0.138	0.019	0.078	5	2	C	30	50.00	80	0.094	0.261	0.178
2	2	S	30	36.00	0	1.701	0.717	1.209	5	2	C	10	67.00	0	1.184	0.591	0.887
2	2	S	30	36.00	80	0.110	0.081	0.095	5	2	C	30	67.00	80	0.563	0.521	0.542
2	2	S	30	44.00	0	1.889	0.644	1.267	5	2	C	30	80.00	0	1.753	0.599	1.176
2	2	S	30	44.00	80	0.124	0.056	0.090	5	2	C	30	80.00	80	0.727	0.311	0.519
2	1	SB	4	36.00	0	-0.017	-0.002	-0.010	5	1	CB	4	73.00	0	-0.167	-0.119	-0.143
2	1	SB	4	36.00	80	0.009	-0.003	0.003	5	1	CB	4	73.00	80	0.012	0.010	0.011
2	1	SB	15	36.00	0	-0.013	0.013	0.000	5	1	CB	15	73.00	0	0.859	0.391	0.625
2	1	SB	15	36.00	80	-0.003	-0.009	-0.006	5	1	CB	15	73.00	80	0.124	0.156	0.140
2	1	SB	30	36.00	0	0.164	0.148	0.156	5	1	CB	30	73.00	0	0.764	0.233	0.499
2	1	SB	30	36.00	80	0.0	0.035	0.018	5	1	CB	30	73.00	80	1.220	0.527	0.874
2	2	SB	4	39.00	0	-0.090	-0.035	-0.063	5	2	CB	4	77.00	0	-0.064	-0.036	-0.050
2	2	SB	4	39.00	80	0.002	-0.009	-0.003	5	2	CB	4	77.00	80	-0.024	-0.019	-0.022
2	2	SB	15	39.00	0	-0.071	0.005	-0.033	5	2	CB	15	77.00	0	0.786	0.276	0.531
2	2	SB	15	39.00	80	-0.016	-0.008	-0.012	5	2	CB	15	77.00	80	0.122	0.182	0.152
2	2	SB	30	39.00	0	0.268	0.310	0.289	5	2	CB	30	77.00	0	0.736	0.337	0.536
2	2	SB	30	39.00	80	0.017	0.032	0.024	5	2	CB	30	77.00	80	0.887	0.387	0.637
2	3	S	4	37.00	0	-0.085	-0.051	-0.068	5	3	C	4	74.00	0	0.156	0.068	0.112
2	3	S	4	37.00	80	0.002	0.001	0.001	5	3	C	4	74.00	80	0.066	0.012	0.039
2	3	S	15	37.00	0	0.121	0.022	0.071	5	3	C	15	74.00	0	0.794	0.512	0.653
2	3	S	15	37.00	80	-0.008	0.002	-0.003	5	3	C	15	74.00	80	0.108	0.191	0.149
2	3	S	30	37.00	0	0.439	0.293	0.366	5	3	C	30	74.00	0	0.700	0.325	0.512
2	3	S	30	37.00	80	0.058	0.036	0.047	5	1	C	30	74.00	80	0.807	0.539	0.673
2	4	S	4	37.00	0	0.048	-0.044	0.002	5	4	C	4	70.00	0	0.073	0.130	0.102
2	4	S	4	37.00	80	0.023	0.005	0.014	5	4	C	4	70.00	80	-0.021	-0.065	-0.043
2	4	S	15	37.00	0	-0.110	-0.034	-0.072	5	4	C	15	70.00	0	0.226	0.202	0.214
2	4	S	15	37.00	80	0.046	0.005	0.026	5	4	C	15	70.00	80	-0.145	-0.019	-0.082
2	4	S	30	37.00	0	0.307	0.098	0.203	5	4	C	30	70.00	0	0.035	0.168	0.102
2	4	S	30	37.00	80	0.047	0.035	0.041	5	4	C	30	70.00	80	-0.023	0.055	0.014
2	5	S	4	33.00	0	-0.157	-0.213	-0.185	5	5	C	4	67.00	0	0.102	0.161	0.131
2	5	S	4	33.00	80	-0.034	-0.012	-0.023	5	5	C	4	67.00	80	0.090	0.027	0.058
2	5	S	15	33.00	0	-0.228	-0.185	-0.207	5	5	C	15	67.00	0	0.102	0.312	0.207
2	5	S	15	33.00	80	-0.062	-0.028	-0.045	5	5	C	15	67.00	80	0.009	0.003	0.006
2	5	S	30	33.00	0	-0.157	-0.189	-0.173	5	5	C	30	67.00	0	0.102	0.149	0.126
2	5	S	30	33.00	60	-0.086	-0.023	-0.054	5	5	C	30	67.00	80	-0.010	0.015	0.002
2	6	S	4	33.00	0	0.112	0.130	0.121	5	6	C	4	66.00	0	-0.321	0.045	-0.138
2	6	S	4	33.00	80	0.036	0.007	0.022	5	6	C	4	66.00	80	-0.002	-0.008	-0.005
2	6	S	15	33.00	0	0.152	0.168	0.160	5	6	C	15	66.00	0	0.215	0.160	0.188
2	6	S	15	33.00	80	0.043	0.008	0.025	5	6	C	15	66.00	80	-0.037	-0.010	-0.023
2	6	S	30	33.00	0	0.152	0.183	0.167	5	6	C	30	66.00	0	0.399	0.160	0.279
2	6	S	30	33.00	80	0.046	0.021	0.034	5	6	C	30	66.00	80	-0.101	-0.020	-0.061
2	7	S	4	33.00	0	0.074	0.036	0.055	5	7	C	4	64.00	0	-0.030	-0.045	-0.037
2	7	S	4	33.00	80	0.032	0.026	0.029	5	7	C	4	64.00	80	0.052	0.004	0.028
2	7	S	15	33.00	0	0.035	0.039	0.037	5	7	C	15	64.00	0	0.347	0.094	0.220
2	7	S	15	33.00	80	0.062	-0.021	0.020	5	7	C	15	64.00	80	0.053	0.031	0.042
2	7	S	30	33.00	0	0.050	0.026	0.038	5	7	C	30	64.00	0	0.490	0.139	0.315
2	7	S	30	33.00	80	0.018	-0.008	0.005	5	7	C	30	64.00	80	0.051	0.025	0.038

Table 8. Correlation of rate of N mineralization values with environmental and soil factors of incubated watershed surface soil samples.

Factor	Correlation Coefficient		
	Brookston Clay	Berrien Sandy Loam	N-5
%C	0.25	0.09	0.30
%N	0.24	0.12	0.30
pH	-0.22	0.00	-0.30
Fixed NH ₄ ⁺	0.24	0.03	0.30
Glucose-C	0.08	0.15	0.30
Temp.	0.30	0.16	0.37
H ₂ O	0.73	0.37	0.85
Temp. x H ₂ O	0.73	0.38	0.83
Multiple	0.90	0.58	---

Table 9. Total surface nitrogen and yearly N mineralized assuming 1% of the total N is mineralized/year

Soil	kgN/ha	N mineralized/year	
		kgN/ha	µgN/g
Berrien sandy Loam	2,466	25	9.6
Brookston clay	6,726	67	26.5
N-5/1	16,591	166	37.0
N-5/2	10,762	108	24.0
N-5/3	13,452	135	30.0
N-5/4	14,797	148	33.0
N-5/5	5,829	58	13.0

Table 10. Total inorganic N in soil-plant residue incubations

Soil	Plant Material	Day 0		Day 15		Day 30		Day 60	
		Ave	Range	Ave	Range	Ave	Range	Ave	Range
Brookston Clay	Rye	20.3	0.8	11.6	2.5	27.5	1.1	59.3	0.3
	Soybean	18.9	5.0	55.3	7.1	88.8	10.5	133.0	1.1
	Wheat	18.8	4.3	6.5	0.8	7.3	0.7	22.6	0.6
	Corn	11.4	15.5	7.4	2.5	8.3	0.4	9.4	1.3
	Control	19.0	2.0	35.8	3.8	56.6	3.4	76.2	14.5
Berrien Sandy Loam	Rye	19.6	0.4	34.8	3.1	50.2	6.4	61.2	2.2
	Soybean	20.7	2.1	35.2	1.2	73.1	9.7	99.2	0.9
	Wheat	21.7	0.7	1.9	0.1	5.4	0.2	6.7	1.1
	Corn	12.5	0.1	2.6	0.2	6.0	0.9	6.6	0.4
	Control	18.7	1.3	33.6	0.2	44.2	0.1	51.0	0.2

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N	PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER									RANGE	AVER	RANGE	AVER	RANGE	AVER	
1	1	S	4	35.00	0	0	1.0	15.6	0.0	0.0	0.8	2.8	16.4	2	1	S	4	27.00	0	0	1.2	10.4	0.0	0.0	0.0	3.0	13.4
1	1	S	4	35.00	0	2	0.7	18.3	0.0	0.0	0.0	2.6	20.8	2	1	S	4	27.00	0	2	2.0	10.0	0.0	0.0	1.1	4.6	14.6
1	1	S	4	35.00	0	5	1.2	20.9	0.0	0.0	0.6	3.9	24.8	2	1	S	4	27.00	0	5	1.0	11.5	0.0	0.0	0.7	4.3	15.8
1	1	S	4	35.00	80	0	10.4	78.1	0.0	0.0	8.2	88.0	166.1	2	1	S	4	27.00	80	0	2.0	99.0	0.0	0.0	4.0	100.0	199.0
1	1	S	4	35.00	80	2	23.2	79.3	0.0	0.0	3.6	85.4	164.7	2	1	S	4	27.00	80	2	6.0	101.0	0.0	0.0	4.0	100.0	201.0
1	1	S	4	35.00	80	5	3.4	88.1	0.0	0.0	7.4	90.3	178.4	2	1	S	4	27.00	80	5	10.0	101.0	0.0	0.0	0.0	94.0	195.6
1	1	S	15	35.00	0	0	1.0	15.6	0.0	0.0	0.8	2.8	18.4	2	1	S	4	36.00	0	0	1.0	10.5	0.0	0.0	0.0	2.0	12.5
1	1	S	15	35.00	0	2	9.5	12.5	0.5	1.8	0.9	5.3	19.6	2	1	S	4	36.00	0	2	0.0	11.0	0.0	0.0	0.4	4.2	15.2
1	1	S	15	35.00	0	5	4.7	13.5	0.0	0.0	0.4	8.3	21.8	2	1	S	4	36.00	0	5	0.0	11.0	0.0	0.0	0.0	4.4	15.4
1	1	S	15	35.00	80	0	10.4	78.1	0.0	0.0	8.2	88.0	166.1	2	1	S	4	36.00	80	0	12.0	107.0	0.0	0.0	2.0	95.0	202.0
1	1	S	15	35.00	80	2	2.0	76.5	2.5	1.3	2.8	88.9	166.6	2	1	S	4	36.00	80	2	4.0	100.0	0.0	0.0	0.0	94.0	194.0
1	1	S	15	35.00	80	5	2.0	73.7	1.5	0.8	1.3	92.4	166.9	2	1	S	4	36.00	80	5	10.0	101.0	0.0	0.0	2.0	93.0	194.0
1	1	S	30	35.00	0	0	1.0	15.6	0.0	0.0	0.8	2.8	18.4	2	1	S	4	44.00	0	0	0.0	11.0	0.0	0.0	2.0	3.0	14.0
1	1	S	30	35.00	0	2	0.6	2.5	0.0	0.0	0.4	8.6	11.1	2	1	S	4	44.00	0	2	0.0	12.0	0.0	0.0	1.8	4.9	16.9
1	1	S	30	35.00	0	5	0.5	2.1	0.0	0.0	0.6	11.3	13.3	2	1	S	4	44.00	0	5	0.0	12.0	0.0	0.0	0.0	4.0	16.0
1	1	S	30	35.00	80	0	10.4	78.1	0.0	0.0	8.2	88.0	166.6	2	1	S	4	44.00	80	0	8.0	106.0	0.0	0.0	6.0	92.0	198.0
1	1	S	30	35.00	80	2	26.0	68.5	0.0	3.0	3.7	91.8	163.3	2	1	S	4	44.00	80	2	3.0	102.5	0.0	0.0	4.0	91.0	193.5
1	1	S	30	35.00	80	5	3.3	64.6	0.0	0.0	4.0	92.9	157.5	2	1	S	4	44.00	80	5	1.0	102.5	0.0	0.0	0.0	93.0	195.5
1	2	S	4	36.00	0	0	0.5	24.6	0.0	0.0	0.4	6.0	30.6	2	1	S	15	27.00	0	0	1.2	10.4	0.0	0.0	0.0	3.0	13.4
1	2	S	4	36.00	0	2	0.1	25.3	0.0	0.0	1.1	5.8	31.1	2	1	S	15	27.00	0	2	1.0	11.5	0.0	0.0	0.4	4.9	16.4
1	2	S	4	36.00	0	5	0.3	23.5	0.0	0.0	0.1	4.6	28.2	2	1	S	15	27.00	0	5	3.0	12.5	0.0	0.0	3.0	4.5	17.0
1	2	S	4	36.00	80	0	0.0	133.0	0.0	0.0	6.3	81.4	214.4	2	1	S	15	27.00	80	0	2.0	99.0	0.0	0.0	4.0	100.0	199.0
1	2	S	4	36.00	80	2	1.0	112.5	0.0	0.0	3.8	81.7	194.2	2	1	S	15	27.00	80	2	2.0	93.0	0.0	0.0	0.0	90.0	183.0
1	2	S	4	36.00	80	5	1.0	115.5	0.0	0.0	2.0	81.4	196.9	2	1	S	15	27.00	80	5	4.0	92.0	0.0	0.0	4.0	92.0	184.0
1	2	S	15	36.00	0	0	0.5	24.6	0.0	0.0	0.4	6.0	30.6	2	1	S	15	36.00	0	0	1.0	10.5	0.0	0.0	0.0	2.0	12.5
1	2	S	15	36.00	0	2	1.3	19.3	0.5	1.8	0.6	7.5	28.5	2	1	S	15	36.00	0	2	1.3	8.0	0.0	0.5	0.0	4.8	13.3
1	2	S	19	36.00	0	5	0.9	15.7	0.0	0.0	1.0	9.9	25.6	2	1	S	15	36.00	0	5	2.7	4.3	0.0	0.0	1.3	5.8	10.2
1	2	S	15	36.00	80	0	0.0	133.0	0.0	0.0	6.3	81.4	214.4	2	1	S	15	36.00	80	0	12.0	107.0	0.0	0.0	2.0	95.0	202.0
1	2	S	15	36.00	80	2	6.0	113.0	0.0	2.0	3.9	85.6	200.6	2	1	S	15	36.00	80	2	0.0	104.0	0.5	1.3	3.0	90.5	195.7
1	2	S	15	36.00	80	5	0.0	105.0	0.0	0.5	2.8	85.4	190.9	2	1	S	15	36.00	80	5	7.0	101.5	0.0	0.0	2.0	97.0	198.5
1	2	S	30	36.00	0	0	0.5	24.6	0.0	0.0	0.4	6.0	30.6	2	1	S	15	44.00	0	0	0.0	11.0	0.0	0.0	2.0	3.0	14.9
1	2	S	30	36.00	0	2	0.0	6.6	0.0	0.0	0.4	11.1	17.7	2	1	S	15	44.00	0	2	0.0	7.1	0.5	0.3	0.0	4.9	12.3
1	2	S	30	36.00	0	5	0.3	1.8	0.0	0.0	5.1	12.5	14.4	2	1	S	15	44.00	0	5	0.0	4.8	0.0	0.0	0.5	5.6	10.4
1	2	S	30	36.00	80	0	0.0	133.0	0.0	0.0	6.3	81.4	214.4	2	1	S	15	44.00	80	0	8.0	106.0	0.0	0.0	6.0	92.0	198.0
1	2	S	30	36.00	80	2	0.2	96.0	0.0	2.5	1.6	90.2	188.7	2	1	S	15	44.00	80	2	2.0	92.0	0.0	1.5	4.0	93.0	186.5
1	2	S	30	36.00	80	5	7.7	88.8	1.5	3.3	7.9	94.9	186.9	2	1	S	15	44.00	80	5	2.0	90.0	0.0	0.0	2.0	94.0	184.0
														2	1	S	30	27.40	0	0	1.2	10.4	0.0	0.0	0.0	3.0	13.4
														2	1	S	30	27.40	0	2	1.7	3.8	0.0	0.0	1.3	9.1	13.0
														2	1	S	30	27.00	0	5	1.1	1.4	0.0	0.0	2.6	5.5	6.9
														2	1	S	30	27.00	80	0	2.0	99.0	0.0	0.0	4.0	100.0	199.0
														2	1	S	30	27.00	80	2	9.0	83.5	0.5	0.3	0.0	92.0	175.7
														2	1	S	30	27.00	80	5	9.0	87.5	0.0	0.0	8.0	94.0	181.5
														2	1	S	30	36.00	0	0	1.0	10.5	0.0	0.0	0.0	2.0	12.5
														2	1	S	30	36.00	0	2	0.4	0.2	0.0	0.0	0.9	7.4	7.6
														2	1	S	30	3640	0	5	0.5	0.6	0.0	0.0	1.6	6.1	6.7
														2	1	S	30	36.00	80	0	12.0	107.0	0.0	0.0	2.0	95.4	202.0
														2	1	S	30	36.40	80	2	2.0	75.0	0.0	1.0	2.0	93.0	169.0
														2	1	S	30	36.40	80	5	5.0	76.5	0.0	0.0	2.0	93.0	169.5
														2	1	S	30	44.00	0	0	0.0	11.0	0.0	0.0	2.0	3.0	14.0
														2	1	S	30	40.00	0	2	0.0	0.4	0.0	0.0	0.0	7.5	7.9
														2	1	S	30	44.00	0	5	0.0	0.0	0.0	0.0	0.9	7.6	7.3
														2	1	S	30	44.00	80	0	8.0	106.0	0.0	0.0	6.0	92.0	198.0
														2	1	S	30	44.00	80	2	10.0	79.0	0.0	0.5	5.0	95.5	175.0

Table 11. Inorganic N analyses ($\mu\text{gN/g}$) during incubation experiments examining denitrification under saturated moisture contents and several temperatures of watershed soils

Table 11 cont.

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N	PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER									RANGE	AVER	RANGE	AVER	RANGE	AVER	
2	2	S	4	27.00	0	0	0.0	18.0	0.0	0.0	0.0	4.3	22.3	2	1	SB	4	36.00	0	0	2.0	50.1	0.0	0.0	0.4	4.2	54.3
2	2	S	4	27.00	0	2	0.0	19.0	0.0	0.0	0.8	6.0	25.0	2	1	SB	4	36.00	0	2	1.7	51.8	0.0	0.0	0.1	3.3	55.0
2	2	S	4	27.00	0	5	1.0	20.5	0.0	0.0	0.0	5.6	26.1	2	1	SB	4	36.00	0	5	1.2	50.7	0.0	0.0	0.0	4.7	55.4
2	2	S	4	27.00	80	0	3.0	106.5	0.0	0.0	6.0	97.0	203.5	2	1	SB	4	36.00	80	0	7.0	142.5	0.0	0.0	2.2	85.8	228.3
2	2	S	4	27.00	80	2	5.0	105.5	0.0	0.0	12.0	96.0	201.5	2	1	SB	4	36.00	80	2	15.0	140.0	0.0	0.0	4.5	85.6	225.6
2	2	S	4	27.00	80	5	2.0	103.0	0.0	0.0	6.0	95.0	198.0	2	1	SB	4	36.00	80	5	8.0	145.0	0.0	0.0	3.0	84.0	229.0
2	2	S	4	36.00	0	0	0.0	18.0	0.0	0.0	0.4	4.6	22.6	2	1	SB	15	36.00	0	0	2.0	50.1	0.0	0.0	0.4	4.2	54.3
2	2	S	4	36.00	0	2	0.0	19.0	0.0	0.0	0.4	5.0	24.0	2	1	SB	15	36.00	0	2	0.0	49.9	1.0	1.5	0.2	4.3	55.7
2	2	S	4	36.00	0	5	0.0	20.0	0.0	0.0	0.4	5.9	25.9	2	1	SB	15	36.00	0	5	4.5	45.9	0.0	0.0	0.4	7.4	54.3
2	2	S	4	36.00	80	0	2.0	104.0	0.0	0.0	4.0	98.0	202.0	2	1	SB	15	36.00	80	0	7.0	142.5	0.0	0.0	2.2	85.8	228.3
2	2	S	4	36.00	80	2	2.0	104.0	0.0	0.0	2.0	95.0	199.0	2	1	SB	15	36.00	80	2	5.0	141.5	1.0	2.0	2.2	83.3	226.8
2	2	S	4	36.00	80	5	2.0	110.0	0.0	0.0	12.0	96.0	206.0	2	1	SB	15	36.00	80	5	1.0	148.5	1.5	0.8	0.9	86.3	235.5
2	2	S	4	44.00	0	0	1.0	17.5	0.0	0.0	0.0	4.4	21.9	2	1	SB	30	36.00	0	0	2.0	50.1	0.0	0.0	0.4	4.2	54.3
2	2	S	4	44.00	0	2	2.0	18.5	0.0	0.0	0.0	4.9	23.4	2	1	SB	30	36.00	0	2	2.0	35.3	0.5	0.8	0.1	6.8	42.8
2	2	S	4	44.00	0	5	1.0	19.5	0.0	0.0	1.3	7.3	26.8	2	1	SB	30	36.00	0	5	9.6	23.9	0.0	0.0	1.7	12.5	36.1
2	2	S	4	44.00	80	0	14.0	98.0	0.0	0.0	6.7	96.3	194.3	2	1	SB	30	36.00	80	0	7.0	142.5	0.0	0.0	2.2	85.8	228.3
2	2	S	4	44.00	80	2	2.0	110.0	0.0	0.0	10.0	98.0	208.0	2	1	SB	30	36.00	80	2	3.0	140.5	0.0	2.0	1.4	87.1	229.6
2	2	S	4	44.00	80	5	4.0	108.0	0.0	0.0	6.0	101.0	209.0	2	1	SB	30	36.00	80	5	3.0	118.5	1.0	1.0	6.4	87.1	206.6
2	2	S	15	27.00	0	0	0.0	18.0	0.0	0.0	0.0	4.3	22.3	2	2	SB	4	39.00	0	0	0.6	21.7	0.0	0.0	0.2	3.4	25.1
2	2	S	15	27.00	0	2	1.0	18.5	0.0	0.0	0.4	7.9	26.4	2	2	SB	4	39.00	0	2	0.0	26.0	0.0	0.0	0.2	2.4	28.4
2	2	S	15	27.00	0	5	0.0	21.0	0.0	0.0	0.5	8.8	29.8	2	2	SB	4	39.00	0	5	1.0	25.8	0.0	0.0	0.1	2.8	28.6
2	2	S	15	27.00	80	0	3.0	106.5	0.0	0.0	6.0	97.0	203.5	2	2	SB	4	39.00	80	0	11.0	111.5	0.0	0.0	3.7	83.6	185.2
2	2	S	15	27.00	80	2	2.0	116.0	0.0	0.0	0.0	96.0	212.0	2	2	SB	4	39.00	80	2	4.0	111.0	0.0	0.0	2.4	83.4	194.4
2	2	S	15	27.00	80	5	14.0	112.0	0.5	0.3	7.0	99.5	211.7	2	2	SB	4	39.00	80	5	7.0	116.5	0.0	0.0	3.0	85.6	202.3
2	2	S	15	36.00	0	0	0.0	18.0	0.0	0.0	0.4	4.6	22.6	2	2	SB	15	39.00	0	0	0.6	21.7	0.0	0.0	0.2	3.4	25.1
2	2	S	15	36.00	0	2	0.0	16.0	0.5	0.8	0.4	7.2	23.9	2	2	SB	15	39.00	0	2	0.2	24.5	1.0	0.5	4.1	5.6	30.5
2	2	S	15	36.00	0	5	0.0	13.0	0.0	0.0	0.4	9.4	22.4	2	2	SB	15	39.00	0	5	1.4	21.2	0.0	0.0	0.3	4.6	26.0
2	2	S	15	36.00	80	0	2.0	104.0	0.0	0.0	4.0	98.0	202.0	2	2	SB	15	39.00	80	0	11.0	111.5	0.0	0.0	3.7	83.8	195.2
2	2	S	15	36.00	80	2	2.0	110.0	0.0	1.5	6.0	101.0	212.5	2	2	SB	15	39.00	80	2	7.0	112.5	2.5	2.8	3.4	86.3	201.6
2	2	S	15	36.00	80	5	7.0	111.5	0.5	0.3	14.0	101.0	212.7	2	2	SB	15	39.00	80	5	3.0	115.5	0.5	0.8	4.0	86.3	202.6
2	2	S	15	44.00	0	0	1.0	17.5	0.0	0.0	0.0	4.4	21.9	2	2	SB	30	30.00	0	0	0.6	21.7	0.0	0.0	0.2	3.4	25.1
2	2	S	15	44.00	0	2	0.0	16.0	0.0	0.5	0.4	7.3	23.8	2	2	SB	30	39.00	0	2	1.6	12.7	0.1	0.0	0.2	6.0	13.7
2	2	S	15	44.00	0	5	2.0	11.0	0.0	0.0	0.2	9.9	20.9	2	2	SB	30	39.00	0	5	2.8	4.6	0.0	0.0	2.0	7.4	12.0
2	2	S	15	44.00	80	0	14.0	98.0	0.0	0.0	6.7	96.3	194.3	2	2	SB	30	39.00	80	0	11.0	111.5	0.0	0.0	3.7	83.8	195.2
2	2	S	15	44.00	80	2	1.0	112.5	0.5	1.3	2.0	92.0	205.7	2	2	SB	30	39.00	80	2	4.0	106.0	0.5	1.8	0.3	86.4	194.2
2	2	S	15	44.00	80	5	5.0	95.5	0.0	0.0	0.0	101.0	196.5	2	2	SB	30	39.00	80	5	0.8	95.1	0.0	0.0	3.0	87.5	182.6
2	2	S	30	27.00	0	0	0.0	18.0	0.0	0.0	0.0	4.3	22.3	2	3	S	4	37.00	0	0	1.6	18.5	0.0	0.0	0.5	4.1	23.5
2	2	S	30	27.00	0	2	3.0	12.5	0.0	0.0	1.0	12.5	25.0	2	3	S	4	37.00	0	2	0.7	23.1	0.0	0.0	0.0	2.3	23.4
2	2	S	30	27.00	0	5	6.0	23.0	0.0	0.0	3.0	4.5	27.5	2	3	S	4	37.00	0	5	1.6	25.2	0.0	0.0	0.3	3.1	28.3
2	2	S	30	27.00	80	0	3.0	106.5	0.0	0.0	6.0	97.0	203.5	2	3	S	4	37.00	80	0	8.4	96.8	6.0	0.0	0.6	85.3	162.1
2	2	S	30	27.00	80	2	9.0	80.5	0.5	0.3	2.0	109.0	189.7	2	3	S	4	37.00	80	2	13.0	96.5	0.0	0.0	3.1	85.9	182.4
2	2	S	30	27.00	80	5	10.0	97.0	0.0	0.0	6.0	105.0	202.0	2	3	S	4	37.00	80	5	0.6	96.4	0.0	0.0	0.4	90.8	187.2
2	2	S	30	36.00	0	0	0.0	18.0	0.0	0.0	0.4	4.6	22.6	2	3	S	15	37.00	0	0	1.6	19.5	0.0	0.0	0.5	4.1	23.5
2	2	S	30	36.00	0	2	0.5	0.6	0.0	0.0	0.0	12.0	12.6	2	3	S	15	37.00	0	2	4.5	13.8	0.0	1.5	3.4	5.6	20.9
2	2	S	30	36.00	0	5	1.0	0.5	0.0	0.0	1.0	14.5	15.0	2	3	S	15	37.00	0	5	3.0	17.8	0.0	0.0	0.8	5.1	22.5
2	2	S	30	36.00	80	0	2.0	104.0	0.0	0.0	4.0	98.0	202.0	2	3	S	15	37.00	80	0	9.4	96.8	0.0	0.0	0.6	25.3	142.1
2	2	S	30	36.00	80	2	2.0	82.0	0.0	1.5	8.0	106.0	189.5	2	3	S	15	37.00	80	2	0.7	96.1	2.5	2.3	2.7	91.5	189.9
2	2	S	30	36.00	80	5	9.0	69.5	0.0	0.0	2.0	112.0	181.5	2	3	S	15	37.00	80	5	1.3	94.9	1.0	1.0	0.7	91.8	187.7
2	2	S	30	44.00	0	0	1.0	17.5	0.0	0.0	0.0	4.4	21.9	2	3	S	30	37.00	0	0	1.6	19.5	0.0	0.0	0.5	4.1	23.5
2	2	S	30	44.00	0	2	0.0	0.4	0.0	0.0	1.0	12.5	12.9	2	3	S	30	37.00	0	2	0.1	8.1	0.0	0.0	0.3	7.8	15.9
2	2	S	30	44.00	0	5	0.6	0.7	0.0	0.0	1.0	11.5	12.2	2	3	S	30	37.00	0	5	2.7	4.5	0.0	0.0	1.2	7.1	11.6
2	2	S	30	44.00	80	0	14.0	98.0	0.0	0.0	6.7	96.3	194.3	2	3	S	30	37.00	80	0	8.4	96.8	0.0	0.0	0.6	35.3	152.6
2	2	S	30	44.00	80	2	5.0	75.5	0.0	1.0	2.0	107.0	183.5	2	3	S	30	37.00	80	2	2.3	84.4	0.5	1.8	0.8	95.6	181.8
2	2	S	30	44.00	80	5	2.0	74.0	0.0	0.0	11.0	106.5	180.5	2	3	S	30	37.00	80	5	9.9	80.9	0.0	0.0	2.7	91.7	172.9

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N	PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER									RANGE	AVER	RANGE	AVER	RANGE	AVER	
2	4	S	4	37.00	0	0	1.2	9.8	0.0	0.0	0.5	4.1	13.9	2	7	S	4	33.00	0	0	1.4	7.3	0.0	0.0	0.2	0.7	8.0
2	4	S	4	37.00	0	2	0.7	8.9	0.0	0.0	3.5	6.5	15.5	2	7	S	4	33.00	0	2	0.5	6.3	0.0	0.0	0.1	0.5	6.8
2	a	S	a	37.00	0	5	0.5	12.2	0.0	0.0	0.0	2.1	14.3	2	7	S	4	33.00	0	5	0.3	6.1	0.0	0.0	0.0	0.5	6.6
2	4	S	4	37.00	80	0	7.6	95.4	0.0	0.0	3.4	87.7	183.6	2	7	S	4	33.00	80	0	3.1	88.4	0.0	0.0	0.3	81.6	170.1
2	4	S	4	37.00	80	2	6.9	91.6	0.0	0.0	0.8	91.0	182.6	2	7	S	4	33.00	80	2	3.4	83.0	0.0	0.0	0.8	80.8	163.8
2	4	S	4	37.00	80	5	2.9	93.3	0.0	0.0	1.9	82.6	175.9	2	7	S	4	33.00	80	5	10.2	77.6	0.0	0.0	3.4	82.4	160.0
2	4	S	15	37.00	0	0	1.2	9.8	0.0	0.0	0.5	4.1	13.4	2	7	S	15	33.00	0	0	1.4	7.3	0.0	0.0	0.2	0.7	8.0
2	4	S	15	37.00	0	2	2.5	10.4	2.5	1.8	0.9	4.8	17.0	2	7	S	15	33.00	0	2	0.5	6.8	0.0	0.0	0.0	0.2	6.9
2	4	3	15	37.00	0	5	0.7	11.6	0.0	0.0	0.0	3.7	15.3	2	7	S	15	33.00	0	5	0.4	6.0	0.0	0.0	0.0	0.2	6.2
2	4	S	15	37.00	80	0	7.6	95.9	0.0	0.0	3.4	87.7	183.6	2	7	S	15	33.00	80	0	3.1	88.4	0.0	0.0	0.3	81.6	170.1
2	4	S	15	37.00	80	2	3.4	86.4	0.5	0.8	0.0	88.6	175.7	2	7	S	15	33.00	80	2	4.0	78.1	0.0	0.0	2.3	85.3	163.4
2	4	S	15	37.00	80	5	1.4	93.4	0.5	0.3	0.4	87.7	181.3	2	7	S	15	33.00	80	5	7.6	98.2	0.0	0.0	0.6	83.0	181.2
2	4	S	30	37.00	0	0	1.2	9.8	0.0	0.0	0.5	4.1	13.9	2	7	S	30	33.00	0	0	1.4	7.3	0.0	0.0	0.2	0.7	8.0
2	4	S	30	37.00	0	2	1.9	5.3	0.0	0.0	0.1	5.8	11.1	2	7	S	30	33.00	0	2	0.6	6.6	0.0	0.0	0.1	0.1	6.7
2	4	S	30	37.00	0	5	0.7	6.0	0.0	0.0	0.0	6.0	12.0	2	7	S	30	33.00	0	5	0.2	6.4	0.0	0.0	0.0	0.4	6.8
2	4	S	30	37.00	80	0	7.6	95.9	0.0	0.0	3.4	87.7	183.6	2	7	S	30	33.00	80	0	3.1	88.4	0.0	0.0	0.3	81.6	170.1
2	4	S	30	37.00	80	2	15.9	84.3	0.0	3.0	16.9	88.6	175.9	2	7	S	30	33.00	80	2	5.4	85.3	0.0	0.0	4.6	78.9	164.2
2	4	S	30	37.00	80	5	12.8	80.4	0.0	0.0	0.3	96.5	176.9	2	7	S	30	33.00	80	5	1.0	92.2	0.0	0.0	3.2	81.7	173.9
2	5	S	4	33.00	0	0	1.5	1.9	0.0	0.0	1.8	5.0	6.9	6	1	C	4	72.00	0	0	0.1	9.8	0.0	0.0	0.5	3.6	13.4
2	5	S	4	33.00	0	2	0.7	2.6	0.0	0.0	0.3	7.4	10.1	6	1	C	4	72.00	0	2	1.3	13.8	1.0	1.0	0.3	5.5	20.3
2	5	S	4	33.00	80	0	3.3	81.6	0.0	0.0	2.3	88.6	170.3	6	1	C	4	72.00	0	5	2.4	12.6	1.0	4.0	0.2	8.7	24.8
2	5	S	4	33.00	80	2	5.5	37.4	0.0	0.0	1.2	89.2	176.0	6	1	C	4	72.00	80	0	3.4	98.3	0.0	0.0	3.7	70.6	168.4
2	5	S	4	33.00	80	5	1.7	86.8	0.0	0.0	5.7	84.9	171.8	6	1	C	4	72.00	80	2	7.0	110.5	0.0	0.0	1.8	67.8	178.3
2	5	S	15	33.00	0	0	1.5	1.9	0.0	0.0	1.8	5.0	6.9	6	1	C	4	72.00	80	5	3.0	106.9	0.0	0.0	4.8	71.8	178.3
2	5	S	15	33.00	0	2	0.4	3.0	0.0	0.0	0.0	2.0	5.0	6	1	C	15	72.00	0	0	0.1	9.8	0.0	0.0	0.5	3.6	13.4
2	5	S	15	33.00	0	5	0.3	4.8	0.0	0.0	0.2	0.9	5.6	6	1	C	15	72.00	0	8	2.2	5.3	0.0	0.0	0.7	11.8	17.0
2	5	S	15	33.00	80	0	3.3	81.6	0.0	0.0	2.3	83.6	170.3	6	1	C	15	72.00	0	5	0.7	4.5	0.0	0.0	0.0	16.2	20.7
2	5	S	15	33.00	80	2	4.7	92.3	0.0	0.0	3.5	89.6	181.9	6	1	C	15	72.00	80	0	3.4	98.3	0.0	0.0	3.7	70.1	168.4
2	5	S	15	33.00	80	5	9.1	93.8	0.0	0.0	3.1	85.1	178.9	6	1	C	15	72.00	80	2	0.5	78.1	2.0	17.5	1.5	74.6	170.2
2	5	S	30	33.00	0	0	1.5	1.9	0.0	0.0	1.8	5.0	6.9	6	1	C	15	72.00	80	5	8.2	54.1	0.5	4.3	1.1	78.1	136.4
2	5	S	30	33.00	0	2	0.2	2.6	0.0	0.0	0.2	2.6	5.2	6	1	C	30	72.00	0	0	0.1	9.8	0.0	0.0	0.5	3.6	13.4
2	5	S	30	33.00	0	5	0.5	4.9	0.0	0.0	0.0	1.3	6.2	6	1	C	30	72.00	0	2	0.3	5.9	0.0	0.0	0.1	18.6	24.5
2	5	S	30	33.00	80	0	3.3	81.6	0.0	0.0	2.3	88.6	170.3	6	1	C	30	72.00	0	5	0.8	5.5	0.0	0.0	0.9	27.6	33.1
2	5	S	30	33.00	80	2	8.2	96.9	0.0	0.0	2.7	88.4	185.3	6	1	C	30	72.00	80	0	3.4	98.3	0.0	0.0	3.7	70.1	168.4
2	5	S	30	33.00	80	5	2.5	91.4	0.0	0.0	1.5	84.1	175.4	6	1	C	30	72.00	80	2	0.6	22.5	1.0	0.5	1.4	79.3	102.3
2	6	S	4	33.00	0	0	1.7	6.5	0.0	0.0	0.0	0.6	7.1	6	1	C	30	72.00	80	5	0.7	7.1	0.0	0.0	0.8	83.3	90.4
2	6	S	4	33.00	0	2	0.0	5.2	0.0	0.0	0.0	0.5	5.7	6	2	C	4	69.00	0	0	0.4	9.5	0.0	0.0	0.4	4.6	14.1
2	6	S	4	33.00	0	5	1.3	3.4	0.0	0.0	0.2	2.3	5.7	6	2	C	4	69.00	0	2	0.3	10.3	0.0	0.0	0.3	7.0	17.3
2	6	S	4	33.00	80	0	3.0	94.2	0.0	0.0	1.5	85.9	180.1	6	2	C	4	69.00	0	5	0.2	8.3	0.5	1.8	0.6	8.2	18.2
2	6	S	4	33.00	80	2	5.8	87.6	0.0	0.0	2.8	83.5	171.1	6	2	C	4	69.00	80	0	6.9	94.1	0.0	0.0	0.7	71.3	165.5
2	6	S	4	33.00	80	5	2.1	91.1	0.0	0.0	2.9	82.6	173.8	6	2	C	4	69.00	80	2	0.3	94.1	0.0	0.0	3.5	71.4	165.5
2	6	S	15	33.00	0	0	1.7	6.5	0.0	0.0	0.0	0.6	7.1	6	2	C	4	69.00	80	5	6.4	91.7	0.5	2.3	1.7	70.4	164.4
2	6	S	15	33.00	0	2	0.2	4.8	0.0	0.0	0.0	0.5	5.3	6	2	C	15	69.00	0	0	0.4	9.5	0.0	0.0	0.4	4.6	14.1
2	6	S	15	33.00	0	5	0.3	2.8	0.0	0.0	0.1	2.1	4.9	6	2	C	15	69.00	0	2	0.1	1.4	0.0	0.0	0.4	11.7	13.1
2	6	S	15	33.00	80	0	3.0	94.2	0.0	0.0	1.5	85.9	180.1	6	2	C	15	69.00	0	5	1.1	1.1	0.0	0.0	0.6	14.2	15.3
2	6	S	15	33.00	80	2	2.6	86.5	0.0	0.0	0.0	80.7	167.2	6	2	C	15	69.00	80	0	6.9	94.6	0.0	0.0	0.7	71.3	165.5
2	6	S	15	33.00	80	5	1.0	90.4	0.0	0.0	3.2	83.4	173.8	6	2	C	15	69.00	80	2	0.6	76.9	3.0	13.5	0.3	74.9	165.3
2	6	S	30	33.00	0	0	1.7	6.5	0.0	0.0	0.0	0.6	7.1	6	2	C	15	69.00	80	5	0.4	45.4	0.0	0.0	1.3	78.3	123.6
2	6	S	30	33.00	0	2	0.6	4.8	0.0	0.0	0.1	0.4	5.3	6	2	C	30	69.00	0	0	0.4	9.5	0.0	0.0	0.4	4.6	14.1
2	6	S	30	33.00	0	5	0.7	2.6	0.0	0.0	0.4	2.1	4.7	6	2	C	30	69.00	0	2	0.3	1.3	0.0	0.0	1.7	16.1	17.5
2	6	S	30	33.00	80	0	3.0	94.2	0.0	0.0	1.5	85.9	160.1	6	2	C	30	69.00	0	5	0.5	1.9	0.0	0.0	2.8	22.5	24.4
2	6	S	30	33.00	80	2	5.3	85.9	0.0	0.0	0.6	81.2	167.1	6	2	C	30	69.00	80	0	6.9	94.1	0.0	0.0	0.7	71.3	165.5
2	6	S	30	33.00	80	5	4.9	84.8	0.0	0.0	3.2	82.0	166.7	6	2	C	30	69.00	80	2	10.5	22.4	0.0	0.0	2.0	82.4	104.8
														6	2	C	30	69.00	80	5	5.1	8.6	0.0	0.0	1.4	80.7	89.3

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N	PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER									RANGE	AVER	RANGE	AVER	RANGE	AVER	
3	1	S	4	26.00	0	0	1.0	19.5	0.0	0.0	0.0	1.0	20.5	3	2	S	4	27.00	0	0	0.1	13.8	0.0	0.0	0.6	2.1	15.9
3	1	S	4	26.00	0	2	2.0	21.0	0.0	0.0	1.0	1.5	22.5	3	2	S	4	27.00	0	2	0.5	15.0	0.0	0.0	0.3	2.4	17.5
3	1	S	4	26.00	0	5	1.0	15.5	0.0	0.0	0.0	2.0	17.5	3	2	S	4	27.00	0	5	0.7	16.1	0.0	0.0	0.2	2.2	18.3
3	1	S	4	26.00	80	0	2.9	98.1	0.0	0.0	2.0	88.0	186.1	3	2	S	4	27.00	80	0	10.0	107.0	0.0	0.0	3.4	85.0	192.0
3	1	S	4	26.00	80	2	3.0	103.5	0.0	0.0	0.0	89.0	192.5	3	2	S	4	27.00	80	2	7.9	103.1	0.0	0.0	0.4	85.2	188.2
3	1	S	4	26.00	80	5	4.0	105.0	0.0	0.0	0.0	87.0	192.0	3	2	S	4	27.00	80	5	10.3	97.8	0.5	0.3	2.1	83.1	181.1
3	1	S	4	35.00	0	0	1.0	19.5	0.0	0.0	0.0	1.0	20.5	3	2	S	4	36.00	0	0	0.0	14.4	0.0	0.0	0.1	2.1	16.4
3	1	S	4	35.00	0	2	1.0	19.5	0.0	0.0	0.0	2.0	21.5	3	2	S	4	36.00	0	2	0.0	15.0	0.0	0.0	0.5	2.3	17.3
3	1	S	4	35.00	0	5	1.0	19.5	0.0	0.0	0.0	2.0	21.5	3	2	S	4	36.00	0	5	0.4	15.9	0.0	0.0	0.3	2.1	18.0
3	1	S	4	35.00	80	0	1.0	105.5	0.0	0.0	4.0	84.0	194.5	3	2	S	4	36.00	80	0	14.9	105.6	0.0	0.0	1.4	84.8	190.4
3	1	S	4	35.00	80	2	12.0	109.0	0.0	0.0	9.0	91.5	200.5	3	2	S	4	36.00	80	2	15.8	102.1	5.5	2.8	1.4	84.8	189.6
3	1	S	4	35.00	80	5	5.0	103.5	0.0	0.0	0.0	85.0	188.5	3	2	S	4	36.00	80	5	6.1	98.9	0.0	0.5	0.0	81.5	180.9
3	1	S	4	42.00	0	0	2.0	16.0	0.0	0.0	0.0	2.0	18.0	3	2	S	4	43.00	0	0	0.2	14.2	0.0	0.0	0.0	2.0	16.2
3	1	S	4	42.00	0	2	2.0	18.0	0.0	0.0	0.0	2.0	20.0	3	2	S	4	43.00	0	2	0.4	15.5	0.0	0.0	0.3	2.3	17.8
3	1	S	4	42.00	0	5	1.0	17.5	0.0	0.0	0.0	1.0	18.5	3	2	S	4	43.00	0	5	0.1	16.1	0.0	0.0	0.1	1.9	18.0
3	1	S	4	42.00	80	0	15.0	93.5	0.0	0.0	14.0	84.0	177.5	3	2	S	4	43.00	80	0	12.0	104.0	0.0	0.0	1.8	86.0	190.0
3	1	S	4	42.00	80	2	9.0	92.5	0.0	0.0	1.0	80.5	173.0	3	2	S	4	43.00	80	2	10.0	99.0	0.0	0.0	4.5	86.4	185.4
3	1	S	4	42.00	80	5	5.0	105.5	0.0	0.0	0.0	79.0	184.5	3	2	S	4	43.00	80	5	8.8	100.6	0.5	0.8	3.6	81.6	182.9
3	1	S	15	26.00	0	0	1.0	19.5	0.0	0.0	0.0	1.0	20.5	3	2	S	15	27.00	0	0	0.1	13.8	0.0	0.0	0.6	2.1	15.9
3	1	S	15	26.00	0	2	2.0	19.0	0.0	0.0	0.0	3.0	22.0	3	2	S	15	27.00	0	2	1.4	16.3	0.0	0.0	0.5	3.6	19.8
3	1	S	15	26.00	0	5	1.0	20.5	0.0	0.0	2.0	2.0	22.5	3	2	S	15	27.00	0	5	1.9	14.0	0.0	0.0	1.2	1.8	15.8
3	1	S	15	26.00	80	0	2.9	98.1	0.0	0.0	2.0	88.0	186.1	3	2	S	15	27.00	80	0	10.0	107.0	0.0	0.0	3.4	85.0	192.0
3	1	S	15	26.00	80	2	8.0	105.0	0.5	0.3	2.0	84.0	189.2	3	2	S	15	27.00	80	2	4.0	106.0	0.5	0.8	3.9	82.6	189.4
3	1	S	15	26.00	80	5	9.0	104.5	0.5	0.3	2.0	80.0	184.7	3	2	S	15	27.00	80	5	14.6	101.7	1.0	0.5	6.8	79.0	181.2
3	1	S	15	35.00	0	0	1.0	19.5	0.0	0.0	0.0	1.0	20.5	3	2	S	15	36.00	0	0	0.0	14.4	0.0	0.0	0.1	2.1	16.4
3	1	S	15	35.00	0	2	1.0	16.5	0.0	0.0	1.0	2.5	19.0	3	2	S	15	36.00	0	2	0.2	11.6	0.0	0.0	0.4	4.3	15.9
3	1	S	15	35.00	0	5	0.0	17.0	0.0	0.0	1.6	1.2	18.2	3	2	S	15	36.00	0	5	2.0	11.9	0.0	0.0	0.7	3.0	14.9
3	1	S	15	35.00	80	0	1.0	105.5	0.0	0.0	4.0	89.0	194.5	3	2	S	15	36.00	80	0	14.9	105.6	0.0	0.0	1.4	84.8	190.4
3	1	S	15	35.00	80	2	13.0	95.5	1.0	0.5	9.0	82.5	178.5	3	2	S	15	36.00	80	2	16.0	108.0	0.5	0.3	0.9	84.6	192.8
3	1	S	15	15.00	80	5	8.0	102.0	3.5	1.8	9.0	82.5	186.2	3	2	S	15	36.00	80	5	9.6	101.2	0.2	2.6	2.6	77.6	181.4
3	1	S	15	42.00	0	0	2.0	16.0	0.0	0.0	0.0	2.0	18.0	3	2	S	15	43.00	0	0	0.2	14.2	0.0	0.0	0.0	2.0	16.2
3	1	S	15	42.00	0	2	1.0	16.5	0.0	0.0	1.0	2.5	19.0	3	2	S	15	43.00	0	2	0.5	11.3	0.0	0.0	0.1	4.1	15.4
3	1	S	15	42.00	0	5	0.0	17.0	0.0	0.0	0.5	0.8	17.8	3	2	S	15	43.00	0	5	0.5	11.4	0.0	0.0	0.8	3.0	14.4
3	1	S	15	42.00	80	0	15.0	93.5	0.0	0.0	14.0	84.0	177.5	3	2	S	15	43.00	80	0	12.0	104.0	0.0	0.0	1.8	86.0	190.0
3	1	S	15	42.00	80	2	10.7	91.3	0.0	0.0	2.0	78.0	169.3	3	2	S	15	43.00	80	2	9.5	102.3	0.5	0.3	1.3	82.6	185.1
3	1	S	15	42.00	80	5	2.8	96.4	2.5	1.8	5.0	79.5	177.6	3	2	S	15	43.00	80	5	7.8	100.1	0.5	0.3	0.0	80.7	181.0
3	1	S	30	26.00	0	0	1.0	19.5	0.0	0.0	0.0	1.0	20.5	3	2	S	30	27.00	0	0	0.1	13.8	0.0	0.0	0.6	2.1	15.9
3	1	S	30	26.00	0	2	5.0	13.5	0.0	0.0	1.0	3.5	17.0	3	2	S	30	27.00	0	2	0.0	7.2	0.0	0.0	1.3	5.5	12.7
3	1	S	30	26.00	0	5	8.0	22.0	0.0	0.0	0.4	0.2	22.2	3	2	S	30	27.00	0	5	4.3	10.9	0.0	0.0	0.0	2.4	13.3
3	1	S	30	26.00	80	0	2.9	98.1	0.0	0.0	2.0	88.0	186.6	3	2	S	30	27.00	80	0	10.0	107.0	0.0	0.0	3.4	85.0	192.0
3	1	S	30	26.00	80	2	11.0	92.5	2.0	2.0	4.0	63.0	177.5	3	2	S	30	27.00	80	2	9.0	101.5	1.0	1.0	3.9	89.1	191.6
3	1	S	30	26.00	80	5	0.0	101.0	0.0	0.0	3.0	73.5	174.5	3	2	S	30	27.00	80	5	2.0	111.0	0.0	0.0	3.1	75.4	186.4
3	1	S	30	35.00	0	0	1.0	19.5	0.0	0.0	0.0	1.0	20.5	3	2	S	30	36.00	0	0	0.0	14.4	0.0	0.0	0.1	2.1	16.4
3	1	S	30	35010	0	2	5.5	9.3	0.0	0.0	2.8	3.4	12.6	3	2	S	30	36.00	0	2	0.4	2.3	0.0	0.0	0.2	6.7	9.0
3	1	S	30	35.00	0	5	3.4	11.3	0.0	0.0	0.4	0.2	11.5	3	2	S	30	36.00	0	5	1.6	4.4	0.0	0.0	1.6	3.6	8.0
3	1	S	30	35.00	80	0	1.0	105.5	0.0	0.0	4.0	89.0	194.5	3	2	S	30	36.00	80	0	14.9	105.6	0.0	0.0	1.4	84.8	190.4
3	1	S	30	35.00	80	2	11.0	96.5	1.5	0.8	0.0	85.0	182.2	3	2	S	30	36.00	80	2	4.2	101.9	2.0	11.5	0.8	87.2	190.6
3	1	S	30	35.00	80	5	12.0	95.0	0.5	0.3	4.0	85.0	180.2	3	2	S	30	36.00	80	5	27.1	109.4	11.5	6.3	0.5	58.6	174.3
3	1	S	30	42.00	0	0	2.0	16.0	0.0	0.0	0.0	2.0	16.0	3	2	S	30	43.00	0	0	0.2	14.2	0.0	0.0	0.0	2.0	16.2
3	1	S	30	42.00	0	2	0.0	7.5	0.0	0.0	0.0	4.0	11.5	3	2	S	30	43.00	0	2	1.6	1.9	0.0	0.0	1.5	6.6	8.5
3	1	S	30	42.00	0	5	0.4	9.5	0.0	0.0	1.0	3.5	13.0	3	2	S	30	43.00	0	5	2.0	3.8	0.0	0.0	1.5	2.1	5.9
3	1	S	30	42.00	80	0	15.0	93.5	0.0	0.0	14.0	64.0	177.5	5	2	S	30	43.00	80	0	12.0	104.0	0.0	0.0	1.8	86.0	190.0
3	1	S	30	42.00	80	2	4.0	93.0	1.0	1.0	0.0	84.0	178.0	3	2	S	30	43.00	80	2	3.1	96.4	0.5	2.3	0.4	81.8	180.5
3	1	S	30	42.00	80	5	6.0	100.0	2.0	5.5	11.0	71.5	177.0	3	2	S	30	43.00	80	5	0.0	114.0	13.5	6.8	15.1	63.8	184.6

Table 11 cont.

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N	PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER									RANGE	AVER	RANGE	AVER	RANGE	AVER	
4	1	C	4	55.00	0	0	0.5	19.1	0.0	0.0	0.5	5.1	24.1	4	2	C	4	54.00	0	0	0.5	5.3	0.0	0.0	1.0	3.5	8.8
4	1	C	4	55.00	0	2	6.8	18.3	0.0	3.0	0.5	9.4	30.7	4	2	C	4	54.00	0	2	0.4	7.5	0.0	0.0	0.5	7.9	15.4
4	1	C	4	55.00	0	5	20.2	22.5	2.5	1.3	6.9	6.8	30.6	4	2	C	4	54.00	0	5	2.0	14.0	0.0	0.0	0.5	7.9	21.9
4	1	C	4	55.00	80	0	14.0	114.0	0.0	0.0	0.5	81.3	195.2	4	2	C	4	54.00	80	0	2.3	85.1	0.0	0.0	2.2	82.9	168.0
4	1	C	4	55.00	80	2	1.0	121.5	0.5	0.8	6.3	86.9	209.2	4	2	C	4	54.00	80	2	22.7	79.4	0.0	0.0	15.8	79.3	157.7
4	1	C	4	55.00	80	5	36.8	107.6	5.0	2.5	11.4	84.9	195.0	4	2	C	4	54.00	80	5	22.5	90.8	1.5	1.3	0.0	79.5	171.5
4	1	C	4	74.00	0	0	0.3	19.3	0.0	0.0	0.0	5.6	24.9	4	2	C	4	72.00	0	0	0.5	4.9	0.0	0.0	0.0	4.0	8.9
4	1	C	4	74.00	0	2	0.7	15.3	0.0	4.5	0.5	9.3	29.2	4	2	C	4	72.00	0	2	1.0	3.5	0.0	2.5	0.9	8.0	14.0
4	1	C	4	74.00	0	5	0.8	12.0	0.0	1.0	0.0	9.9	22.9	4	2	C	4	72.00	0	5	0.0	2.0	0.5	0.3	0.0	11.0	13.3
4	1	C	4	74.00	80	0	12.0	111.0	0.0	0.0	3.4	86.9	197.9	4	2	C	4	72.00	80	0	7.1	81.4	0.0	0.0	2.4	86.2	167.6
4	1	C	4	74.00	80	2	17.3	104.3	0.0	5.0	1.0	86.2	195.5	4	2	C	4	72.00	80	2	16.5	81.4	0.5	3.8	2.4	79.1	164.3
4	1	C	4	74.00	80	5	16.1	98.9	1.0	4.0	1.0	90.0	192.9	4	2	C	4	72.00	80	5	9.4	77.9	0.5	5.3	2.4	83.8	166.9
4	1	C	4	88.00	0	0	1.9	19.6	0.0	0.0	0.0	5.7	25.3	4	2	C	4	87.00	0	0	0.5	5.1	0.0	0.0	0.0	4.0	9.1
4	1	C	4	88.00	0	2	0.6	15.9	0.5	4.3	1.1	9.8	29.9	4	2	C	4	87.00	0	2	3.0	1.5	0.0	2.5	0.5	8.5	12.5
4	1	C	4	88.00	0	5	1.1	4.6	0.5	1.3	1.1	12.1	18.0	4	2	C	4	87.00	0	5	10.0	6.0	0.0	0.0	7.0	7.5	13.5
4	1	C	4	88.00	80	0	12.0	115.0	0.0	0.0	0.0	85.4	200.4	4	2	C	4	87.00	80	0	7.3	86.3	0.0	0.0	2.4	81.5	167.8
4	1	C	4	88.00	80	2	3.0	107.5	0.5	4.6	0.5	89.6	201.6	4	2	C	4	87.00	80	2	0.0	85.2	0.5	3.8	4.9	77.8	166.8
4	1	C	4	88.00	80	5	9.8	98.1	1.9	4.6	4.3	89.0	191.2	4	2	C	4	87.00	80	5	5.7	79.8	2.0	7.0	2.4	81.5	168.3
4	1	C	15	55.00	0	0	0.5	19.1	0.0	0.0	0.5	5.1	24.1	4	2	C	15	54.00	0	0	0.5	5.3	0.0	0.0	1.0	3.5	8.8
4	1	C	15	55.00	0	2	23.4	14.0	0.0	0.0	5.3	11.5	25.5	4	2	C	15	54.00	0	2	1.0	1.5	0.0	0.0	2.0	16.0	17.5
4	1	C	15	55.00	0	5	1.0	27.7	0.0	0.0	0.0	0.9	28.6	4	2	C	15	54.00	0	5	16.0	12.0	0.0	0.0	6.1	8.9	20.9
4	1	C	15	55.00	80	0	14.0	114.0	0.0	0.0	0.5	81.3	195.2	4	2	C	15	54.00	80	0	2.3	85.1	0.0	0.0	2.2	82.9	168.0
4	1	C	15	55.00	80	2	38.0	100.0	0.0	0.0	12.3	84.8	184.8	4	2	C	15	54.00	80	2	13.7	74.9	0.5	1.3	2.3	87.4	163.6
4	1	C	15	55.00	80	5	19.5	91.3	0.0	0.0	4.5	79.6	170.9	4	2	C	15	54.00	80	5	13.9	100.1	0.0	0.0	6.8	71.5	171.6
4	1	C	15	74.00	0	0	0.3	19.3	0.0	0.0	0.0	5.6	24.9	4	2	C	15	72.00	0	0	0.5	4.9	0.0	0.0	0.0	4.0	8.9
4	1	C	15	74.00	0	2	0.0	2.9	0.0	0.0	0.5	15.3	18.2	4	2	C	15	72.00	0	2	1.8	1.9	0.0	0.0	0.0	15.0	16.9
4	1	C	15	74.00	0	5	0.8	2.4	0.0	0.0	0.5	13.9	16.3	4	2	C	15	72.00	0	5	2.1	1.9	0.0	0.0	8.0	16.0	17.9
4	1	C	15	74.00	80	0	12.0	111.0	0.0	0.0	3.4	86.9	197.9	4	2	C	15	72.00	80	0	7.1	81.4	0.0	0.0	2.4	86.2	167.6
4	1	C	15	74.00	80	2	6.6	70.1	0.0	0.0	1.9	95.1	165.2	4	2	C	15	72.00	80	2	2.0	41.0	2.0	3.0	2.3	88.5	132.5
4	1	C	15	74.00	80	5	9.0	49.5	0.0	0.0	4.3	91.6	141.1	4	2	C	15	72.00	80	5	4.6	11.7	0.0	0.0	4.8	96.8	108.5
4	1	C	15	88.00	0	0	1.9	19.6	0.0	0.0	0.0	5.7	25.3	4	2	C	15	87.00	0	0	0.5	5.1	0.0	0.0	0.0	4.0	9.1
4	1	C	15	88.00	0	2	0.1	2.3	0.0	0.0	1.0	15.7	18.0	4	2	C	15	87.00	0	2	0.0	3.0	0.0	0.0	0.0	14.0	17.0
4	1	C	15	88.00	0	5	1.0	1.3	0.0	0.0	0.5	14.3	15.6	4	2	C	15	87.00	0	5	1.0	2.5	0.0	0.0	1.0	16.5	19.0
4	1	C	15	88.00	80	0	12.0	115.0	0.0	0.0	0.0	85.4	200.4	4	2	C	15	87.00	80	0	7.3	86.3	0.0	0.0	2.4	81.5	167.8
4	1	C	15	88.00	80	2	10.3	74.4	0.0	0.0	3.5	95.4	169.9	4	2	C	15	87.00	80	2	0.0	14.0	0.0	3.5	0.0	114.0	131.5
4	1	C	15	88.00	80	5	3.9	62.4	0.0	0.0	3.9	88.3	150.8	4	2	C	15	87.00	80	5	3.0	22.5	0.0	0.0	2.5	88.8	111.3
4	1	C	30	55.00	0	0	0.5	19.1	0.0	0.0	0.5	5.1	24.1	4	2	C	30	54.00	0	0	0.5	5.3	0.0	0.0	1.0	3.5	8.8
4	1	C	30	55.00	0	2	1.4	3.3	0.0	0.0	4.0	19.6	22.9	4	2	C	30	54.00	0	2	0.0	5.0	0.0	0.0	4.0	31.0	36.0
4	1	C	30	55.00	0	5	0.7	1.8	0.0	0.0	2.1	3.6	5.4	4	2	C	30	54.00	0	5	1.0	3.5	0.0	0.0	10.0	15.0	18.5
4	1	C	30	55.00	80	0	14.0	114.0	0.0	0.0	0.5	81.3	195.2	4	2	C	30	54.00	80	0	2.3	85.1	0.0	0.0	2.2	82.9	168.0
4	1	C	30	55.00	80	2	45.9	28.4	0.0	0.0	9.1	97.4	125.9	4	2	C	30	54.00	80	2	36.1	32.1	0.5	0.3	0.0	93.1	125.4
4	1	C	30	55.00	80	5	12.0	39.0	0.0	0.0	17.7	76.3	115.2	4	2	C	30	54.00	80	5	4.1	7.1	0.0	0.0	13.6	84.0	91.0
4	1	C	30	74.00	0	0	0.3	19.3	0.0	0.0	0.0	5.6	24.9	4	2	C	30	72.00	0	0	0.5	4.9	0.0	0.0	0.0	4.0	8.9
4	1	C	30	74.00	0	2	0.0	0.0	0.0	0.0	0.6	22.5	22.5	4	2	C	30	72.00	0	2	0.0	0.4	0.0	0.0	0.0	31.0	31.4
4	1	C	30	74.00	0	5	1.8	0.9	0.0	0.0	6.6	29.5	30.4	4	2	C	30	72.00	0	5	0.9	0.4	0.0	0.0	9.0	39.5	39.9
4	1	C	30	74.00	80	0	12.0	111.0	0.0	0.0	3.4	86.9	197.9	4	2	C	30	72.00	80	0	7.1	81.4	0.0	0.0	2.4	86.2	167.6
4	1	C	30	74.00	80	2	6.5	12.8	0.0	0.0	21.8	87.1	99.8	4	2	C	30	72.00	80	2	5.1	4.6	0.0	0.0	2.4	98.0	102.5
4	1	C	30	74.00	80	5	4.2	7.8	0.0	0.0	11.5	95.3	103.0	4	2	C	30	72.00	80	5	7.1	3.6	0.0	0.0	0.0	87.4	90.9
4	1	C	30	88.00	0	0	1.9	19.6	0.0	0.0	0.0	5.7	25.3	4	2	C	30	87.00	0	0	0.9	5.1	0.0	0.0	0.0	4.0	9.1
4	1	C	30	88.00	0	2	1.5	1.6	0.0	0.0	0.0	26.6	28.3	4	2	C	30	87.00	0	2	0.5	0.8	0.0	0.0	2.0	31.0	31.8
4	1	C	30	88.00	0	5	1.4	1.9	0.0	0.0	5.5	29.9	31.8	4	2	C	30	87.00	0	5	0.0	0.0	0.0	0.0	1.0	32.5	32.5
4	1	C	30	88.00	80	0	12.0	115.0	0.0	0.0	0.0	85.4	200.4	4	2	C	30	87.00	80	0	7.3	86.3	0.0	0.0	2.4	81.5	167.8
4	1	C	30	88.00	80	2	0.0	16.0	0.0	0.0	0.0	100.0	116.0	4	2	C	30	87.00	80	2	3.0	3.5	0.0	0.0	5.2	102.4	105.9
4	1	C	30	88.00	80	5	5.4	5.1	0.0	0.0	8.3	92.5	97.6	4	2	C	30	87.00	80	5	5.0	2.5	0.0	0.5	7.4	88.8	91.8

Table 11 cont.

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL	PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL
							RANGE	AVER	RANGE	AVER	RANGE	AVER	N								RANGE	AVER	RANGE	AVER	RANGE	AVER	N
5	1	C	4	50.00	0	0	2.0	22.0	0.0	0.0	0.9	4.4	26.4	5	2	C	4	50.00	0	0	9.0	4.5	0.0	0.0	5.1	5.6	10.1
5	1	C	4	50.00	0	2	6.0	24.0	0.0	0.5	0.4	4.7	29.2	5	2	C	4	50.00	0	2	2.0	12.0	1.5	0.8	0.0	4.5	17.3
5	1	C	4	50.00	0	5	3.0	28.5	0.0	0.0	0.4	4.7	33.2	5	2	C	4	50.00	0	5	4.0	15.0	0.5	0.3	1.8	5.9	21.1
5	1	C	4	50.00	80	0	2.0	111.0	0.0	0.0	4.5	80.9	191.9	5	2	C	4	50.00	80	0	9.0	92.3	0.0	0.0	2.2	75.4	167.7
5	1	C	4	50.00	80	2	2.0	118.0	0.0	1.0	0.0	78.8	197.7	5	2	C	4	50.00	80	2	4.5	92.3	0.5	1.8	2.3	75.3	169.3
5	1	C	4	50.00	80	5	11.0	120.5	1.0	1.5	2.3	77.5	199.5	5	2	C	4	50.00	80	5	16.0	98.0	1.5	1.3	0.0	76.5	175.7
5	1	C	4	66.00	0	0	0.0	23.0	0.0	0.0	0.0	4.0	27.0	5	2	C	a	67.00	0	0	0.7	9.6	0.0	0.0	1.0	2.5	12.1
5	1	C	4	66.00	0	2	2.0	24.0	0.5	1.8	0.5	5.4	31.1	5	2	C	4	67.00	0	2	1.0	13.5	0.5	3.3	0.0	4.0	20.8
5	1	C	4	66.00	0	5	0.0	21.0	0.5	0.8	0.5	6.8	28.5	5	2	C	4	67.00	0	5	0.9	19.6	1.5	2.3	0.4	4.9	26.7
5	1	C	4	66.00	80	0	7.0	115.5	0.0	0.0	2.4	80.4	195.9	5	2	C	4	67.00	80	0	51.0	109.5	0.0	0.0	2.3	78.1	187.6
5	1	C	4	66.00	80	2	11.0	115.5	1.0	2.0	2.3	75.8	193.2	5	2	C	4	67.00	80	2	9.0	95.5	0.0	3.0	0.0	79.3	177.8
5	1	C	4	66.00	80	5	11.0	117.5	0.0	1.5	0.0	72.2	191.2	5	2	C	4	67.00	80	5	2.0	99.0	0.0	3.0	2.3	78.1	180.1
5	1	C	4	79.00	0	0	1.0	22.5	0.0	0.0	0.0	4.0	26.5	5	2	C	4	80.00	0	0	0.0	10.0	0.0	0.0	0.0	3.0	13.0
5	1	C	4	79.00	0	2	0.0	23.0	0.5	1.8	0.0	5.3	30.0	5	2	C	4	80.00	0	2	6.0	20.0	0.0	4.0	0.0	4.0	28.0
5	1	C	4	79.00	0	5	1.0	20.5	0.0	0.5	0.5	7.4	28.4	5	2	C	4	80.00	0	5	2.0	24.0	0.5	1.8	1.0	3.5	29.3
5	1	C	4	79.00	80	0	5.0	122.5	0.0	0.0	2.4	77.9	200.4	5	2	C	4	80.00	80	0	0.0	98.4	0.0	0.0	2.4	73.2	171.6
5	1	C	4	79.00	80	2	7.0	116.5	0.5	2.3	0.0	71.9	190.6	5	2	C	4	80.00	80	2	0.0	91.2	1.0	3.5	4.8	81.6	176.3
5	1	C	4	79.00	80	5	22.0	116.0	1.0	2.0	2.4	68.3	186.3	5	2	C	4	80.00	80	5	0.0	96.0	0.5	3.3	0.0	76.8	176.1
5	1	C	15	50.00	0	0	2.0	22.0	0.0	0.0	0.0	4.4	26.4	5	2	C	15	50.00	0	0	9.0	4.5	0.0	0.0	5.1	5.6	10.1
5	1	C	15	50.00	0	2	5.0	15.5	0.0	0.0	2.3	7.8	23.3	5	2	C	15	50.00	0	2	16.0	20.0	1.5	0.8	4.7	5.3	26.1
5	1	C	15	50.00	0	5	1.1	1.4	0.0	0.0	0.9	8.0	9.5	5	2	C	15	50.00	0	5	13.0	7.5	0.0	0.0	1.6	6.2	13.7
5	1	C	15	50.00	80	0	2.0	111.0	0.0	0.0	4.5	80.9	191.9	5	2	C	15	50.00	80	0	9.0	92.3	0.0	0.0	2.2	75.4	167.7
5	1	C	15	50.00	80	2	6.6	97.7	1.5	1.3	26.9	67.4	166.4	5	2	C	15	50.00	80	2	13.5	101.3	1.0	2.5	0.0	76.5	180.2
5	1	C	15	50.00	80	5	4.0	117.0	0.0	0.0	2.3	73.0	190.0	5	2	C	15	50.00	80	5	17.7	101.1	0.0	0.5	22.0	52.0	153.6
5	1	C	15	66.00	0	0	0.0	23.0	0.0	0.0	0.0	4.0	27.0	5	2	C	15	67.00	0	0	0.7	9.6	0.0	0.0	1.0	2.5	12.1
5	1	C	15	66.00	0	2	4.7	11.6	0.0	0.0	1.0	8.4	20.0	5	2	C	15	67.00	0	2	2.0	30.0	0.0	0.0	0.0	3.0	33.0
5	1	C	15	66.00	0	5	1.1	1.4	0.0	0.0	0.9	9.3	10.8	5	2	C	15	67.00	0	5	4.0	26.0	0.0	0.0	1.0	2.5	28.5
5	1	C	15	66.00	80	0	7.0	115.5	0.0	0.0	2.4	80.4	195.9	5	2	C	15	67.00	80	0	51.0	109.5	0.0	0.0	2.3	78.1	187.6
5	1	C	15	66.00	80	2	11.8	99.1	0.0	0.5	2.4	80.4	180.0	5	2	C	15	67.00	80	2	26.0	50.0	0.5	3.8	39.3	94.3	148.1
5	1	C	15	66.00	80	5	5.1	100.4	0.0	0.0	2.3	71.0	171.5	5	2	C	15	67.00	80	5	0.0	65.3	0.0	0.0	4.6	77.0	142.3
5	1	C	15	79.00	0	0	1.0	22.5	0.0	0.0	0.0	4.0	26.5	5	2	C	15	80.00	0	0	0.0	10.0	0.0	0.0	0.0	3.0	13.0
5	1	C	15	79.00	0	2	1.9	5.8	0.0	0.0	1.0	10.5	16.3	5	2	C	15	80.00	0	2	3.0	29.5	0.0	0.0	0.0	3.0	32.5
5	1	C	15	79.00	0	5	0.0	0.5	0.0	0.0	1.0	10.5	11.0	5	2	C	15	80.00	0	5	0.0	26.0	0.0	0.0	0.0	3.0	29.0
5	1	C	15	79.00	80	0	5.0	122.5	0.0	0.0	2.4	77.9	200.4	5	2	C	15	80.00	80	.0	0.0	98.4	0.0	0.0	2.4	73.2	171.6
5	1	C	15	79.00	80	2	0.0	101.0	0.0	0.0	2.4	77.9	178.9	5	2	C	15	80.00	80	2	4.8	72.0	3.5	1.8	2.4	78.0	151.7
5	1	C	15	79.00	0	5	0.0	93.4	0.0	0.0	4.8	71.9	165.3	5	2	C	15	80.00	80	5	4.8	69.6	0.0	0.0	4.8	74.4	144.0
5	1	C	30	50.00	0	0	2.0	22.0	0.0	0.0	0.9	4.4	26.4	S	2	C	30	50.00	0	0	9.0	4.5	0.0	0.0	5.1	5.6	10.1
5	1	C	30	50.00	0	2	4.0	2.0	0.0	0.0	3.1	11.4	13.4	5	2	C	30	50.00	0	2	3.3	9.3	0.0	0.0	1.3	9.3	18.6
5	1	C	30	50.00	.0	S	0.0	0.0	0.0	0.0	2.0	13.0	13.0	5	.2	C	30	50.00	0	5	0.9	0.4	0.0	0.0	9.0	14.5	14.9
5	1	C	30	50.00	80	0	2.0	111.0	0.0	0.0	4.5	80.9	191.9	5	2	C	30	50.00	80	0	9.0	92.3	0.0	0.0	2.2	75.4	167.7
5	1	C	30	50.00	80	2	2.3	86.5	0.0	0.0	6.8	79.8	166.3	5	2	C	30	50.00	80	2	27.0	72.0	1.0	4.5	13.5	76.6	153.1
5	1	C	30	50.00	83	5	54.1	107.9	0.0	0.0	26.9	58.4	166.4	5	2	C	30	50.00	80	5	5.0	20.5	9.0	4.5	29.3	68.6	93.6
5	1	C	30	66.00	0	0	0.0	23.0	0.0	0.0	0.0	4.0	27.0	5	2	C	30	67.00	0	0	0.7	9.6	0.0	0.0	1.0	2.5	12.1
5	1	C	30	66.00	0	2	0.0	0.0	0.0	0.0	1.0	13.5	13.5	5	2	C	30	67.00	0	2	0.1	0.9	0.0	0.0	2.0	15.0	15.9
5	1	C	30	66.00	0	5	0.0	0.0	0.0	0.0	7.0	14.5	14.5	5	2	C	30	67.00	0	5	1.0	0.5	0.0	0.0	3.0	24.5	25.0
5	1	C	30	66.00	80	0	7.0	115.5	0.0	0.0	2.4	80.4	195.9	5	2	C	30	67.00	80	0	51.0	109.5	0.0	0.0	2.3	78.1	187.6
5	1	C	30	66.00	80	2	23.3	62.9	0.5	0.3	11.7	78.0	141.2	5	2	C	30	67.00	80	2	0.0	35.0	1.0	0.5	4.6	84.0	119.5
5	1	C	30	66.00	80	5	13.9	58.3	1.5	0.8	23.3	65.3	124.2	5	2	C	30	67.00	80	5	2.3	8.1	0.0	0.0	7.0	82.8	90.9
5	1	C	30	79.00	0	0	1.0	22.5	0.0	0.0	0.0	4.0	26.5	5	2	C	30	80.00	0	0	0.0	10.0	0.0	0.0	0.0	3.0	13.0
5	1	C	30	79.00	0	2	0.0	0.0	0.0	0.0	1.0	12.5	12.5	5	2	C	30	80.00	0	2	0.5	0.3	0.0	0.0	4.0	17.0	17.3
5	1	C	30	79.00	0	5	0.0	0.0	0.0	0.0	5.0	13.5	13.5	5	2	C	30	80.00	0	5	1.0	0.5	0.0	0.0	1.0	24.5	25.0
5	1	C	30	79.00	80	0	5.0	122.5	0.0	0.0	2.4	77.9	200.4	5	2	C	30	80.00	80	0	0.0	98.4	0.0	0.0	2.4	73.2	171.6
5	1	C	30	79.00	80	2	4.8	69.5	0.0	0.5	0.0	61.5	151.5	5	2	C	30	80.00	80	2	1.0	22.5	1.0	0.5	2.4	85.2	108.2
5	1	C	30	79.00	80	5	12.0	68.3	0.5	0.3	21.6	63.5	132.1	5	2	C	30	60.00	80	5	3.0	15.5	4.5	5.3	0.0	60.0	80.8

Table 11 cont.

PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N	PL	DP	SOL	TMP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER									RANGE	AVER	RANGE	AVER	RANGE	AVER	
5	1	CB	4	73.00	0	0	0.2	10.6	0.0	0.0	0.1	8.5	19.1	5	4	C	4	70.00	0	0	0.0	4.4	0.0	0.0	0.0	1.2	5.6
5	1	CB	4	73.00	0	2	1.0	14.5	0.5	0.3	0.6	10.0	24.8	5	4	C	4	70.00	0	2	0.6	3.8	0.0	0.0	0.0	1.5	5.3
5	1	CB	4	73.00	0	5	4.6	12.7	0.0	6.5	0.8	14.3	33.5	5	4	C	4	70.00	0	5	0.1	2.3	0.0	0.0	0.5	33.8	6.0
5	1	CB	4	73.00	80	0	4.0	109.0	0.0	0.0	0.3	79.4	188.4	5	4	C	4	70.00	80	0	2.2	82.7	0.0	0.0	1.1	62.6	145.3
5	1	CB	4	73.00	80	2	1.0	106.5	0.0	0.0	0.9	81.3	187.7	5	4	C	4	70.00	80	2	8.7	86.3	0.0	0.0	0.8	59.6	145.8
5	1	CB	4	73.00	80	5	2.8	98.6	4.0	5.0	1.5	83.1	186.7	5	4	C	4	70.00	80	5	6.0	103.0	21.0	11.5	1.5	61.1	175.6
5	1	CB	15	73.00	0	0	0.2	10.6	0.0	0.0	0.1	8.5	19.1	5	4	C	15	70.00	0	0	0.0	4.4	0.0	0.0	0.0	1.2	5.6
5	1	CB	15	73.00	0	2	2.4	1.9	0.0	0.0	0.4	19.1	21.0	5	4	C	15	70.00	0	2	0.7	2.8	0.0	0.0	0.1	2.9	5.7
5	1	CB	15	73.00	0	5	0.3	1.5	0.0	0.0	0.3	25.1	26.7	5	4	C	15	70.00	0	5	0.4	1.6	0.0	0.0	0.3	7.4	9.0
5	1	CB	15	73.00	60	0	4.0	109.0	0.0	0.0	0.3	79.4	188.4	5	4	C	15	70.00	80	0	2.2	82.7	0.0	0.0	1.1	62.6	145.3
5	1	CB	15	73.00	80	2	9.2	78.6	0.0	6.5	0.6	68.8	173.9	5	4	C	15	70.00	80	2	8.7	69.1	2.0	21.5	0.0	60.0	170.6
5	t	CB	15	73.00	80	5	4.3	50.0	0.0	0.0	0.0	95.5	145.5	5	4	C	15	70.00	80	5	3.0	70.1	2.0	21.0	0.0	64.7	155.8
5	1	CB	30	73.00	0	0	0.2	10.6	0.0	0.0	0.1	8.5	19.1	5	4	C	30	70.00	0	0	0.0	4.4	0.0	0.0	0.0	1.2	5.6
5	1	CB	30	73.00	0	2	0.9	2.3	0.0	0.0	0.1	32.8	35.2	5	4	C	30	70.00	0	2	0.1	4.1	0.0	0.0	0.1	6.1	10.7
5	1	CB	30	73.00	0	5	3.2	3.3	0.0	0.0	1.1	47.1	50.4	5	4	C	30	70.00	0	5	0.1	1.9	0.0	0.0	0.0	14.0	15.9
5	1	CB	30	73.00	80	0	4.0	109.0	0.0	0.0	0.3	79.4	188.4	5	4	C	30	70.00	80	0	2.2	82.7	0.0	0.0	1.1	62.0	145.3
5	1	CB	30	73.00	80	2	3.6	9.5	0.0	0.0	3.1	101.4	110.9	5	4	C	30	70.00	80	2	10.4	54.7	4.5	32.8	1.7	62.6	150.1
5	1	CB	30	73.00	80	5	5.3	7.8	0.0	0.0	2.0	105.0	112.7	5	4	C	30	70.00	80	5	3.7	44.9	0.0	18.0	2.1	66.1	131.0
5	2	CB	4	77.00	0	0	0.2	16.3	0.0	0.0	0.0	5.4	23.7	5	5	C	4	67.00	0	0	0.2	3.8	0.0	0.0	0.1	0.9	4.7
5	2	CB	4	77.00	0	2	0.7	20.8	0.0	0.0	0.1	8.3	29.1	5	5	C	4	67.00	0	2	0.3	3.1	0.0	0.0	0.0	0.9	4.6
5	2	CB	4	77.00	0	5	0.5	16.6	0.5	5.3	0.0	11.9	33.8	5	5	C	4	67.00	0	5	0.2	1.7	0.0	0.0	0.0	2.8	4.5
5	2	CB	4	77.00	80	0	1.8	99.1	0.0	0.0	2.6	81.2	180.3	5	5	C	4	67.00	80	0	23.3	109.3	0.0	0.0	10.5	67.8	177.1
5	2	CB	4	77.00	80	2	2.0	104.0	0.0	0.0	0.6	81.5	185.5	5	5	C	4	67.00	80	2	0.4	91.3	0.0	0.0	1.5	53.1	144.4
5	2	CB	4	77.00	80	5	5.0	103.5	1.0	5.5	1.6	66.1	195.1	5	5	C	4	67.00	80	5	6.7	95.6	0.0	0.0	0.0	53.6	149.2
5	2	CB	15	77.00	0	0	0.2	18.3	0.0	0.0	0.0	5.4	23.7	5	5	C	15	67.00	0	0	0.2	3.8	0.0	0.0	0.1	0.9	4.7
5	2	CB	15	77.00	0	2	0.1	3.8	0.0	0.0	0.1	16.4	20.2	5	5	C	15	67.00	0	2	0.5	3.1	0.0	0.0	0.2	1.1	4.1
5	2	CB	15	77.00	0	5	0.7	4.6	0.0	0.0	1.0	24.4	29.0	5	5	C	15	67.00	0	5	0.1	0.8	0.0	0.0	0.3	3.8	4.6
5	2	CB	15	77.00	80	0	1.8	99.1	0.0	0.0	2.6	61.2	180.3	5	5	C	15	67.00	80	0	23.3	109.3	0.0	0.0	10.5	67.8	177.1
5	2	CB	15	77.00	80	2	7.1	66.1	1.0	11.5	0.3	89.9	167.5	5	5	C	15	67.00	80	2	0.8	96.2	0.5	11.3	2.0	51.1	158.6
5	2	CB	15	77.00	80	5	3.7	39.8	0.0	0.0	0.0	95.9	135.7	5	5	C	15	67.00	80	5	0.1	89.1	1.0	18.5	0.9	52.6	160.1
5	2	CB	30	77.00	0	0	0.2	18.3	0.0	0.0	0.0	5.4	23.7	5	5	C	30	67.00	0	0	0.2	3.8	0.0	0.0	0.1	0.9	4.7
5	2	CB	30	77.00	0	2	1.4	4.2	0.0	0.0	2.8	29.3	33.5	5	5	C	30	67.00	0	2	0.7	3.1	0.0	0.0	0.2	2.5	5.6
5	2	CB	30	77.00	0	5	2.3	3.4	0.0	0.0	1.8	38.7	42.1	5	5	C	30	67.00	0	5	0.1	1.8	0.0	0.0	0.9	5.9	7.7
5	2	CB	30	77.00	80	0	1.8	99.1	0.0	0.0	2.6	81.2	180.3	5	5	C	30	67.00	80	0	23.3	109.3	0.0	0.0	10.5	67.8	177.1
5	2	CB	30	77.00	80	2	1.5	16.8	0.0	0.0	0.3	99.5	116.3	5	5	C	30	67.00	80	2	2.0	83.8	1.5	27.8	0.9	51.9	163.5
5	2	CB	30	77.00	80	5	1.6	14.3	0.0	0.0	1.6	98.3	112.6	5	5	C	30	67.00	80	5	6.2	78.3	0.0	23.0	0.6	52.7	154.0
5	3	C	4	74.00	0	0	8.9	14.2	0.0	0.0	3.0	4.3	18.4	5	6	C	4	66.00	0	0	0.3	2.0	0.0	0.0	0.5	2.8	4.3
5	3	C	4	74.00	0	2	0.6	10.4	0.0	0.0	0.6	4.4	14.8	5	6	C	4	66.00	0	2	3.7	3.8	0.0	0.0	3.7	4.5	8.4
5	3	C	4	74.00	0	5	0.7	6.3	0.5	3.8	0.1	8.0	18.1	5	6	C	4	66.00	0	5	0.3	1.6	0.0	0.0	0.9	2.8	4.4
5	3	C	4	74.00	80	0	4.0	105.0	0.0	0.0	4.7	65.3	170.2	5	6	C	4	66.00	80	0	13.2	89.4	0.0	0.0	0.6	55.1	144.5
5	3	C	4	74.00	80	2	1.8	92.0	0.0	0.0	0.9	66.3	156.2	5	6	C	4	66.00	80	2	0.3	89.8	0.0	0.0	0.6	49.4	139.1
5	3	C	4	74.00	80	5	7.4	94.2	0.5	4.8	2.6	68.3	167.2	5	6	C	4	66.00	80	5	1.4	93.0	0.0	0.0	0.6	46.6	139.6
5	3	C	15	74.00	0	0	8.9	14.2	0.0	0.0	3.0	4.3	18.4	5	6	C	15	66.00	0	0	0.3	2.0	0.0	0.0	0.5	2.8	4.3
5	3	C	15	74.00	0	2	0.1	2.9	0.0	0.0	0.1	9.3	12.2	5	6	C	15	66.00	0	2	0.2	1.3	0.0	0.0	0.6	2.2	3.5
5	3	C	15	74.00	0	5	0.7	1.1	0.0	0.0	0.5	16.3	17.4	5	6	C	15	66.00	0	5	0.1	0.9	0.0	0.0	0.2	2.1	3.0
5	3	C	15	74.00	80	0	4.0	105.0	0.0	0.0	4.7	65.3	170.2	5	6	C	15	66.00	80	0	13.2	89.4	0.0	0.0	0.6	55.1	144.5
5	3	C	15	74.00	80	2	2.5	70.1	0.0	14.5	1.5	71.6	156.2	5	6	C	15	66.00	80	2	3.3	91.9	0.5	4.3	0.6	45.7	141.7
5	3	C	15	74.00	80	5	5.5	40.3	0.5	0.3	2.7	77.1	117.6	5	6	C	15	66.00	80	5	3.4	87.6	1.0	6.5	1.5	44.6	138.7
5	3	C	30	74.00	0	0	8.4	14.2	0.0	0.0	3.0	4.3	18.4	5	6	C	30	66.00	0	0	0.3	2.0	0.0	0.0	0.5	2.8	4.8
5	3	C	30	74.00	0	2	0.8	3.5	0.0	0.0	0.7	16.5	20.0	5	6	C	30	66.00	0	2	.0.0	0.9	0.0	0.0	0.6	2.9	3.8
5	3	C	30	74.00	0	5	1.3	2.8	0.0	0.0	0.1	28.1	30.9	5	6	C	30	66.00	0	5	0.3	0.9	0.0	0.0	0.2	3.6	4.5
5	3	C	30	74.00	80	0	4.0	105.0	0.0	0.0	4.7	65.3	170.2	5	6	C	30	66.00	80	0	13.2	89.4	0.0	0.0	0.6	55.1	144.5
5	3	C	30	74.00	80	2	1.2	17.9	0.0	3.0	1.2	77.6	98.5	5	6	C	30	66.00	80	2	0.0	97.2	2.5	12.3	1.5	45.6	155.0
5	3	C	30	74.00	80	5	2.1	7.1	0.0	0.0				5	6	C	30	66.00	80	5							

Table 11 cont

PL	OP	SOL	THP	WC	TRT	DAY	NO3		NO2		NH4		TOTAL N	PL	DP	SOL	TMP	WC	TRT	NO3		NO2		NH4		TOTAL N
							RANGE	AVER	RANGE	AVER	RANGE	AVER								DAY	RANGE	AVER	RANGE	AVER	RANGE	
5	7	C	4	64.00	0	0	0.1	1.6	0.0	0.0	0.2	2.4	3.9	N5	3	4	67.00	0	0	8.9	61.1	0.0	0.0	0.4	6.7	67.8
5	7	C	4	64.00	0	2	0.0	1.7	0.0	0.0	0.4	2.1	3.8	N5	3	4	67.00	0	2	1.7	62.8	0.5	3.3	0.7	11.8	77.8
5	7	C	4	64.00	0	5	0.2	2.0	0.0	0.0	0.6	2.3	4.3	N5	3	4	67.00	0	5	1.2	57.6	0.5	9.3	0.1	20.1	86.9
5	7	C	4	64.00	80	0	0.6	99.4	0.0	0.0	3.3	56.3	155.6	N5	3	4	67.00	80	0	3.0	137.5	0.0	0.0	1.0	79.1	216.5
5	7	C	4	64.00	80	2	10.6	89.5	0.0	0.0	2.7	50.5	140.0	N5	3	4	67.00	80	2	3.0	106.5	0.5	2.3	1.5	89.1	197.9
5	7	C	4	64.00	80	5	4.4	97.6	0.0	0.0	2.4	45.8	143.4	N5	3	4	67.00	80	5	1.0	124.5	2.0	10.0	3.3	97.8	232.3
5	7	C	15	64.00	0	0	0.1	1.6	0.0	0.0	0.2	2.4	3.9	N5	3	15	67.00	0	0	8.9	61.1	0.0	0.0	0.4	6.7	67.8
5	7	C	15	64.00	0	2	0.1	0.6	0.0	0.0	0.3	1.6	2.4	N5	3	15	67.00	0	2	0.1	7.1	0.0	2.5	0.5	26.3	35.8
5	7	C	15	64.00	0	5	1.0	1.0	0.0	0.0	3.8	3.8	4.8	N5	3	15	67.00	0	5	1.3	7.3	0.0	0.0	2.3	42.4	49.8
5	7	C	15	64.00	80	0	0.6	99.4	0.0	0.0	3.3	56.3	155.6	N5	3	15	67.00	80	0	3.0	137.5	0.0	0.0	1.0	79.1	216.6
5	7	C	15	64.00	80	2	0.5	87.1	1.5	2.3	2.8	45.6	135.0	N5	3	15	67.00	80	2	3.0	84.2	1.0	10.0	2.0	103.0	197.2
5	7	C	15	64.00	80	5	1.9	82.6	0.5	2.8	1.8	44.3	129.6	N5	3	15	67.00	80	5	3.1	27.6	0.0	0.0	1.0	120.5	148.1
5	7	C	30	64.00	0	0	0.1	1.6	0.0	0.0	0.2	2.4	3.9	N5	3	30	67.00	0	0	8.9	61.1	0.0	0.0	0.4	6.7	67.8
5	7	C	30	64.00	0	2	0.2	0.6	0.0	0.0	2.2	3.6	4.2	N5	3	30	67.00	0	2	0.1	8.3	0.0	0.0	1.7	48.9	57.2
5	7	C	30	64.00	0	5	0.1	0.8	0.0	0.0	0.5	3.1	3.9	N5	3	30	67.00	0	5	0.1	0.5	0.0	0.0	2.1	60.3	60.8
5	7	C	30	64.00	80	0	0.6	99.4	0.0	0.0	3.3	56.3	155.6	N5	3	30	67.00	80	0	3.0	137.5	0.0	0.0	1.0	79.1	216.6
5	7	C	30	64.00	80	2	0.0	81.2	0.0	8.5	0.0	49.2	138.9	N5	3	30	67.00	80	2	1.7	2.6	0.0	0.0	11.0	127.5	130.1
5	7	C	30	64.00	80	5	0.8	77.7	1.5	10.3	0.3	40.3	128.3	N5	3	30	67.00	80	5	2.7	5.9	0.0	0.0	11.0	133.5	139.4
														N5	4	4	69.00	0	0	3.4	43.8	0.0	0.0	2.3	8.4	52.3
N5	1		0	70.00	0	0	0.3	33.1	1.0	0.05	0.0	9.1	42.7	N5	4	4	69.00	0	2	0.8	45.1	0.0	2.5	0.3	9.8	57.3
N5	1		4	70.00	0	2	0.4	30.4	0.5	5.8	0.3	15.0	51.2	N5	4	4	69.00	0	5	1.7	38.3	0.0	7.5	0.0	13.4	59.2
N5	1		4	70.00	0	5	4.1	18.4	2.0	24.0	0.2	21.0	63.4	N5	4	4	69.00	80	0	5.0	138.5	0.0	0.0	0.0	84.1	222.6
N5	1		4	70.00	80	0	4.0	112.0	0.0	0.5	1.4	71.9	184.0	N5	4	4	69.00	80	2	9.0	137.5	0.0	3.5	0.6	80.7	221.7
N5	1		4	70.00	80	2	13.0	109.5	0.0	6.5	2.3	78.3	194.3	N5	4	4	69.00	80	5	2.0	132.0	0.0	7.0	2.5	86.3	225.2
N5	1		4	70.00	80	5	8.8	99.6	0.5	30.8	1.2	84.8	215.1	N5	4	15	69.00	0	0	3.4	43.8	0.0	0.0	2.3	8.4	52.3
N5	1		15	70.00	0	0	0.3	33.1	1.0	0.5	0.0	9.1	42.7	N5	4	15	69.00	0	2	0.4	6.2	0.5	0.3	0.2	17.6	24.0
N5	1		15	70.00	0	2	0.0	0.0	0.0	0.0	2.2	32.4	32.4	N5	4	15	69.00	0	5	0.1	0.6	0.0	0.0	0.1	26.8	27.5
N5	t		15	70.00	0	5	2.8	1.4	0.0	0.0	0.5	42.8	44.1	N5	4	15	69.00	80	0	5.0	138.5	0.0	0.0	0.0	84.1	222.6
N5	1		15	70.00	80	0	4.0	112.0	0.0	0.5	1.4	71.9	184.4	N5	4	15	69.00	80	2	2.0	100.0	1.0	1.5	1.8	88.1	189.6
N5	1		15	70.00	80	2	0.8	15.1	0.5	2.3	0.0	94.8	112.1	N5	4	15	69.00	80	5	0.7	55.6	0.0	0.0	5.3	100.3	156.0
N5	1		15	70.00	80	5	0.5	10.8	0.0	0.0	5.0	111.5	122.3	N5	4	30	69.00	0	0	3.4	43.8	0.0	0.0	2.3	8.4	52.3
N5	1		30	70.00	0	0	0.3	33.1	1.0	0.5	0.0	9.1	42.7	N5	4	30	69.00	0	2	1.1	1.8	0.0	0.0	0.6	32.2	33.9
N5	1		30	70.00	0	2	0.2	35.7	0.0	0.0	2.3	48.4	84.1	N5	4	30	69.00	0	5	0.2	0.1	0.0	0.0	1.6	39.0	39.1
N5	1		30	70.00	0	5	0.2	2.8	0.0	0.0	1.6	68.8	71.6	N5	4	30	69.00	80	0	5.0	138.5	0.0	0.0	0.0	84.1	222.6
N5	1		30	70.00	80	0	4.0	112.0	0.0	0.5	1.4	71.9	184.4	N5	4	30	69.00	80	2	4.5	13.9	0.0	0.0	2.0	103.0	116.9
N5	1		30	70.00	80	2	0.2	8.4	0.0	0.0	1.0	125.5	133.9	N5	4	30	69.00	80	5	2.8	8.4	0.0	0.0	3.0	113.5	121.9
N5	1		30	70.00	80	5	3.3	9.3	0.0	0.0	5.0	134.5	143.8	N5	5	4	55.00	0	0	1.0	28.6	0.0	0.0	0.4	6.1	34.7
N5	2		4	63.00	0	0	0.6	28.1	0.0	0.0	0.1	5.9	34.0	N5	5	4	55.00	0	2	1.1	29.9	0.0	0.0	0.2	6.4	36.3
N5	2		4	63.00	0	2	2.5	26.9	1.0	4.0	0.7	10.3	41.2	N5	5	4	55.00	0	5	0.3	28.0	0.0	0.0	0.9	7.1	35.2
N5	2		4	63.00	0	5	1.6	25.6	1.0	8.0	0.4	13.3	46.9	N5	5	4	55.00	80	0	1.0	119.5	0.0	0.0	3.0	85.6	205.1
N5	2		4	63.00	80	0	2.0	123.0	0.0	0.0	4.6	84.0	207.0	N5	5	4	55.00	80	2	10.0	114.0	0.0	0.0	1.2	84.7	198.7
N5	2		4	63.00	80	2	3.0	120.5	0.5	3.8	0.4	86.5	210.7	N5	5	4	55.00	80	5	0.0	107.0	0.0	0.0	1.2	84.1	191.1
N5	2		4	63.00	80	5	10.0	115.0	1.0	9.5	0.3	92.1	216.6	N5	5	15	55.00	0	0	1.0	28.6	0.0	0.0	0.4	6.1	34.7
N5	2		15	63.00	0	0	0.6	28.1	0.0	0.0	0.1	5.9	34.0	N5	5	15	55.00	0	2	3.0	21.8	1.0	6.5	2.4	10.3	38.6
N5	2		15	63.00	0	2	0.9	1.1	0.0	0.0	0.6	20.0	21.1	N5	5	15	55.00	0	5	0.4	2.3	0.0	0.0	0.4	16.7	19.0
N5	2		15	63.00	0	5	0.0	1.1	0.0	0.0	3.9	24.8	25.9	N5	5	15	55.00	80	0	1.0	119.5	0.0	0.0	3.0	85.6	205.1
N5	2		15	63.00	80	0	2.0	123.0	0.0	0.0	4.6	84.0	207.0	N5	5	15	55.00	80	2	7.0	106.5	0.5	5.8	3.0	86.2	198.4
N5	2		15	63.00	80	2	1.0	84.3	1.0	10.0	2.0	94.2	188.5	N5	5	15	55.00	80	5	5.4	90.8	0.0	0.0	4.5	92.1	182.9
N5	2		15	63.00	80	5	3.4	51.2	0.0	0.0	0.0	103.0	154.2	N5	5	30	55.00	0	0	1.0	28.6	0.0	0.0	0.4	6.1	34.7
N5	2		30	63.00	0	0	0.6	28.1	0.0	0.0	0.1	5.9	34.0	N5	5	30	55.00	0	2	1.4	3.2	0.0	0.0	0.1	20.6	23.8
N5	2		30	63.00	0	2	0.3	0.6	0.0	0.0	0.2	14.0	14.6	N5	5	30	55.00	0	5	2.5	6.6	0.0	0.0	0.4	40.3	46.9
N5	2		30	63.00	0	5	1.1	0.6	0.0	0.0	1.2	12.4	12.9	N5	5	30	55.00	80	0	1.0	119.5	0.0	0.0	3.0	85.6	205.1
N5	2		30	63.00	80	0	2.0	123.0	0.0	0.0	4.6	84.0	207.0	N5	5	30	55.00	80	2	11.5	94.3	0.5	6.3	5.5	92.8	193.2
N5	2		30	63.10	80	2	1.2	11.8	0.0	0.0	6.0	107.0	118.8	N5	5	30	55.00	80	5		42.6	0.0				

Table 12 Denitrification rate coefficients of representative soil samples at saturated or near saturated moisture content.

PL	DP	SOL	TMP	WC	TRT	RD-2	RD-5	RD-AVE	PL	DP	SOL	TMP	WC	TRT	RD-2	RD-5	RD-AVE
1	1	S	4	35.00	0	-0.080	-0.058	-0.069	4	1	C	4	55.00	0	-0.055	-0.044	-0.049
1	1	S	4	35.00	80	-0.008	-0.024	-0.016	4	1	C	4	55.00	80	-0.035	0.007	-0.014
1	1	S	15	35.00	0	0.044	0.029	0.036	4	1	C	4	74.00	0	-0.013	0.079	0.033
1	1	S	15	35.00	80	0.002	0.010	0.006	4	1	C	4	74.00	80	0.008	0.015	0.011
1	1	S	30	35.00	0	0.915	0.401	0.658	4	1	C	4	88.00	0	-0.013	0.240	0.114
1	1	S	30	35.00	80	0.044	0.038	0.041	4	1	C	4	88.00	80	0.013	0.024	0.019
1	2	S	4	36.00	0	-0.014	0.009	-0.002	4	1	C	15	55.00	0	0.155	-0.074	0.040
1	2	S	4	36.00	60	0.064	0.028	0.056	4	1	C	15	55.00	80	0.066	0.044	0.055
1	2	S	15	36.00	0	0.079	0.090	0.084	4	1	C	15	74.00	0	0.948	0.417	0.682
1	2	S	15	36.00	80	0.073	0.040	0.060	4	1	C	15	74.00	80	0.230	0.162	0.196
1	2	S	30	36.00	0	0.650	0.523	0.590	4	1	C	15	88.00	0	1.071	0.543	0.807
1	2	S	30	36.00	80	0.150	0.074	0.112	4	1	C	15	88.00	80	0.218	0.122	0.170
									4	1	C	30	55.00	0	0.878	0.472	0.675
3	1	S	4	26.00	0	-0.037	0.046	0.004	4	1	C	30	55.00	80	0.695	0.215	0.455
3	1	S	4	26.00	80	-0.027	-0.014	-0.020	4	1	C	30	74.00	0	*****	0.613	*****
3	1	S	4	35.00	0	0.0	0.0	0.0	4	1	C	30	74.00	80	1.080	0.531	0.806
3	1	S	4	35.00	80	-0.016	0.004	-0.006	4	1	C	30	88.00	0	1.253	0.467	0.860
3	1	S	4	42.00	0	-0.059	-0.018	-0.030	4	1	C	30	88.00	80	0.986	0.623	0.805
3	1	S	4	42.00	80	0.005	-0.024	-0.009	4	2	C	4	54.00	0	-0.174	-0.194	-0.184
3	1	S	15	26.00	0	0.013	-0.010	0.001	4	2	C	4	54.00	80	0.035	-0.016	0.010
3	1	S	15	26.00	80	-0.035	-0.013	-0.024	4	2	C	4	72.00	0	-0.101	0.151	0.025
3	1	S	15	35.00	0	0.084	0.027	0.055	4	2	C	4	72.00	80	-0.023	-0.004	-0.013
3	1	S	15	35.00	80	0.047	0.003	0.025	4	2	C	4	87.00	0	0.121	-0.033	0.044
3	1	S	15	42.00	0	-0.015	-0.012	-0.014	4	2	C	4	87.00	80	-0.015	-0.001	-0.008
3	1	S	15	42.00	80	0.012	-0.010	0.001	4	2	C	15	54.00	0	0.631	0.163	0.234
3	1	S	30	26.00	0	0.184	-0.024	0.080	4	2	C	15	54.00	80	0.055	-0.032	0.011
3	1	S	30	26.00	80	0.019	-0.006	0.006	4	2	C	15	72.00	0	0.474	0.189	0.332
3	1	S	30	35.00	0	0.370	0.109	0.240	4	2	C	15	72.00	80	0.308	0.388	0.348
3	1	S	30	35.00	80	0.040	0.020	0.030	4	2	C	15	87.00	0	0.265	0.143	0.204
3	1	S	30	42.00	0	0.379	0.104	0.242	4	2	C	15	87.00	80	0.798	0.269	0.533
3	1	S	30	42.00	80	-0.003	-0.024	-0.013	4	2	C	30	54.00	0	0.029	0.083	0.056
3	2	S	4	27.00	0	-0.042	-0.031	-0.036	4	2	C	30	54.00	80	0.484	0.497	0.491
3	2	S	4	27.00	80	0.019	0.017	0.018	4	2	C	30	72.00	0	1.253	0.501	0.877
3	2	S	4	36.00	0	-0.020	-0.020	-0.020	4	2	C	30	72.00	80	1.437	0.624	1.030
3	2	S	4	36.00	80	0.004	0.012	0.008	4	2	C	30	87.00	0	*****	0.624	*****
3	2	S	4	43.00	0	-0.044	-0.025	-0.034	4	2	C	30	87.00	80	1.603	0.672	1.137
3	2	S	4	43.00	80	0.025	0.005	0.015									
3	2	S	15	27.00	0	-0.083	-0.003	-0.043	N5	1		4	70.00	0	-0.036	-0.047	-0.041
3	2	S	15	27.00	80	0.001	0.009	0.005	N5	1		4	70.00	80	-0.015	-0.029	-0.022
3	2	S	15	36.00	0	0.108	0.038	0.073	N5	t		15	70.00	0	*****	0.636	
3	2	S	15	36.00	80	-0.013	0.003	-0.005	N5	1		15	70.00	80	0.936	0.469	0.702
3	2	S	15	43.00	0	0.114	0.044	0.079	N5	1		30	70.00	0	-0.030	0.497	0.233
3	2	S	15	43.00	80	0.007	0.007	0.007	N5	1		30	70.00	80	1.297	0.499	0.898
3	2	S	30	27.00	0	0.325	0.047	0.186	N5	2		4	63.00	0	-0.047	-0.036	-0.042
3	2	S	30	27.00	80	0.021	-0.007	0.007	N5	2		4	63.00	80	-0.005	-0.002	-0.004
3	2	S	30	36.00	0	0.917	0.237	0.577	N5	2		15	63.00	0	1.620	0.648	1.134
3	2	S	30	36.00	80	0.011	-0.018	-0.004	N5	2		15	63.00	80	0.133	0.175	0.154
3	2	S	30	43.00	0	1.006	0.264	0.635	N5	2		30	63.00	0	1.923	0.769	1.346
3	2	S	30	43.00	80	0.026	-0.030	-0.002	N5	2		30	63.00	80	1.172	0.544	0.858
									N5	3		4	67.00	0	-0.039	-0.018	-0.029
6	1	C	4	72.00	0	-0.206	-0.099	-0.153	N5	3		4	67.00	80	0.117	0.004	0.061
6	1	C	4	72.00	80	-0.058	-0.016	-0.037	N5	3		15	67.00	0	0.925	0.425	0.675
6	1	C	15	72.00	0	0.307	0.156	0.231	N5	3		15	67.00	80	0.189	0.321	0.255
6	1	C	15	72.00	80	0.014	0.104	0.059	N5	3		30	67.00	0	0.998	0.961	0.980
6	1	C	30	72.00	0	0.254	0.116	0.185	N5	3		30	67.00	80	1.984	0.630	1.307
6	1	C	30	72.00	80	0.726	0.526	0.626	N5	4		4	69.00	0	-0.042	-0.009	-0.025
6	2	C	4	69.00	0	-0.040	-0.010	-0.025	N5	4		4	69.00	80	-0.008	-0.001	-0.005
6	2	C	4	69.00	80	0.0	0.000	0.000	N5	4		15	69.00	0	0.962	0.858	0.910
6	2	C	15	69.00	0	0.957	0.431	0.694	N5	4		15	69.00	80	0.155	0.183	0.169
6	2	C	15	69.00	80	0.020	0.146	0.083	N5	4		30	69.00	0	1.596	1.216	1.408
6	2	C	30	69.00	0	0.994	0.322	0.658	N5	4		30	69.00	80	1.149	0.561	0.855
6	2	C	30	69.00	80	0.718	0.479	0.598	N5	5		4	55.00	0	-0.022	0.004	-0.009
									N5	5		4	55.00	80	0.024	0.022	0.023
									N5	5		15	55.00	0	0.005	0.504	0.255
									N5	5		15	55.00	80	0.032	0.055	0.043
									N5	5		30	55.00	0	1.095	0.293	0.694
									N5	5		30	55.00	80	0.087	0.206	0.146

Table 13 Potential N fixation (expressed as μg ethylene produced/g soil/day) and extractable N in representative soil profile samples.

Soil	Day 7			Day 14		
	N fixation	$\mu\text{g NO}_3\text{-N/g}$	$\mu\text{g NH}_4\text{-N/g}$	N fixation	$\mu\text{g NO}_3\text{-N/g}$	$\mu\text{g NH}_4\text{-N/g}$
2-1	0.002	0.2	15.7	0.002	0.3	15.5
2-1-b	<0.002	10.1	10.3	<0.002	8.7	14.9
2-2	0.002	3.1	21.8	<0.002	1.2	26.6
2-2-b	0.002	1.3	11.0	<0.002	1.3	10.5
2-4	<0.002	0.0	6.3	<0.002	0.6	5.0
2-6	<0.002	2.3	1.2	<0.002	3.1	1.8
5-1	0.021	1.2	35.2	0.015	2.6	24.7
5-1-b	0.039	0.1	38.9	0.031	0.9	25.4
5-2	0.014	3.6	31.3	0.012	1.0	20.7
5-2-b	0.019	0.7	36.7	0.010	1.0	23.9
5-4	0.003	0.2	13.3	0.002	0.6	14.0
5-6	0.443	0.5	2.8	0.062	0.6	6.0

APPENDIX

Rates of gaseous loss, mineralization and plant uptake of nitrogen in soil systems: A Literature Review.

Rates of gaseous loss, mineralization and plant uptake of nitrogen in soil systems: A literature review.

C. Grant Kowalenko. S.R.I.

The purpose of this review is to present the extent of knowledge of the rate of processes that are important in the nitrogen economy of soils, particularly under field conditions. Some mention will be given to factors effecting these rates, but most of the effort is directed toward presenting a range of the rates of these processes. It will be seen that very few reports deal directly with the measurement of rates, especially under field conditions and much of the information is derived from literature dealing with other aspects. An attempt is made to present a unified term for the rate, but this is not always possible. The term chosen was kg/ha/day.

GASEOUS LOSS OF N

Several mechanisms have been shown as being possible for the loss of N from the soil system. These include volatilization as NH_3 , biological denitrification, chemical decomposition of oxidized inorganic N forms and gaseous leakage during nitrification and plant growth. Only the first two mechanisms will be considered in terms of rates of loss and the later two will be discussed briefly in terms of what is known of the mechanisms.

(1) Chemical decomposition of oxidized inorganic N forms

The term chemical decomposition of oxidized inorganic N is used to distinguish this chemical mechanism from NH_3 volatilization. There has been considerable recurring interest in possible losses of nitrogen from soils and mechanisms have been discussed (Broadbent and Clark, 1965; Reuss and Smith, 1965; Allison, 1966; Woldendorp, 1968; Chao, 1967; Bremner and Nelson, 1968; Nelson and Bremner, 1969, 1970; Nommik and Thorin, 1972; Bollag *et al.*, 1973; Van Clumput and Baert, 1976). Most of the processes require fairly specific conditions before they proceed (eg) low pH and/or significant concentrations of NO_2^- . Agricultural soils are not always acidic nor have significant concentrations of nitrite been detected, however, certain agricultural practices such as band fertilization may result in lowering soil pH and formation of high nitrite concentrations in localized areas (Pang *et al.*, 1975a,b) which could result in chemical decomposition involving nitrite. Although mechanisms and conditions of chemical decomposition of oxidized forms of inorganic nitrogen have been clearly shown, the quantitative significance of this loss mechanism has not been shown in field situations. Without quantitative measurements, rates can not be estimated.

(2) Gaseous "leakage" during nitrification and plant growth

Gaseous losses of nitrogen as nitrous oxide has been shown to occur during nitrification (Yoshida and Alexander, 1970). It can be argued that conditions required to produce significant amounts of nitrous oxide are not usually present in soils, however, localized situations such as during fertilization may encourage the process. The study referred to examined only one species of *Nitrosomonas* in culture systems, and more extensive studies of different organisms and in soil situations are needed to show the quantitative significance in soil systems.

Nitrogen gases can also be lost from plants during growth. Viets (1965) has discussed some of the possible mechanisms of loss. Recently Craswell and Martin (1975a,b) and Daigger *et al.* (1976) have suggested that significant losses of N due to plants does occur in cropping systems. Other workers (eg. Rumberg and Sneva. 1970) have noted losses of N during plant maturation. but attributed these losses to shedding of pollen. anthers and filaments. Further study on this area is required to quantitatively evaluate this possible N lost from field systems. From the work of Daigger *et al.* (1976). the period during crop maturation appears to be the critical time. possibly during extensive translocation of N compounds.

(3) Volatilization as NH₃

Several reviews (Allison, 1955, 1965; Harmsen and Kolenbrander, 1965; Woldendorp, 1968) have shown the following factors influence the volatilization of nitrogen as NH₃:

- (i) pH and/or CaCO₃ content
 - high pH encourages loss. insignificant losses below pH 6.
 - formation of NH₃ results in localized pH increase.
- (ii) soil texture (C.E.C.)
 - sandy soils (low C.E.C.) result in higher loss.
- (iii) moisture content of soil
 - drying increases loss.
- (iv) temperature
 - increased losses with higher temperatures
- (v) method of fertilizer application
 - losses greater during surface application of ammonia (anhydrous) applied to dry sandy soils.
- (vi) form of fertilizer or associated cation
 - volatilization shown to decrease in the following order:
NH₃ (liquid or anhydrous) > (NH₄)₂SO₄ > NH₄NO₃ > NH₄-phosphates.

This information suggests that NH₃ volatilization is significant only under particular fertilization practices such as surface application. anhydrous application to dry sandy soils or application to limed or high pH soils. It is probable that for this reason most of the recent studies of NH₃ volatilization are oriented toward surface application systems. Studies on fertilization of forests are concerned primarily with surface application (Overrein, 1968, 1969; Volk, 1970; Nommik. 1973; Mahendrappa and Ogden, 1973; Mahendrappa and Salonijs, 1974) but these will not be considered further. Similarly. studies with agricultural orientation have been primarily concerned with surface applications of fertilizer materials and have been mainly done under laboratory conditions.

Before a consideration of rates of NH₃ lost from soils that can be derived from the literature. one should be careful to know how this rate was calculated. Several laboratory (Filimonova and Strel'nikova, 1974; Mills *et al.*, 1974) and field (Makarov, 1960; McGarity and Rajaratnam, 1973) experiments serve to illustrate the point. In order to have a comparable rate term throughout the

discussion, the term kg NH₃-N lost/ha/day is used. However, the interval over which this loss is measured will influence the rate expressed on a per day measurement. Suppose ammonium volatilization occurs very rapidly in the first few hours of the measurement duration of a day, then decreases to negligible amounts for the remainder of the day, the maximum instantaneous rate of loss may be very large at first and negligible later on. Calculation of the rate of loss on a per day basis over that day of measurement then, is really an average rate over that period of time that measurement is made. For this reason, the time interval of the measurement of NH₃ loss and subsequent average rate per day calculation is given.

Table 1 gives rates of NH₃ loss under a variety of conditions and rates vary from 0-10.8 kg N/ha/day. When considering a particular soil situation each of these factors must be considered in terms of their effect on the rate of NH₃ loss. Mills *et al.* (1974) showed similar results using NH₄Cl instead of urea and several pH ranges. Besides pH, they superimposed plant growth, which in general reduced the amount of NH₃ lost when compared to a fallow system. This is probably because the plant takes up NH₄-N, reducing the amount available for volatilization. Chao and Kroontje (1964) showed that as water evaporated from a soil in a flow through system, the rate of NH₃ loss increased. They presented the equation

$$N / t = at^b$$

where N/t = ppm NH₃ loss/hr, t = hrs & a and b = constants to show the effect of water saturated and unsaturated are flowing over a soil. The a and b constants were 48.43 and -0.6819. and 10.43 and -0.3714 for unsaturated and saturated atmospheres, respectively. They also derived linear regression equations for four soils in terms of the amounts of NH₃ lost in the form

$$Y = bX + C$$

where Y = ppm NH₃ volatilized. X = NH₃ application rate, b and C = constants. The constants were different for each soil.

The results for three Ontario soils (Table 2) show that liming these soils (initial pH were all near 5) increased NH₃ loss. The rate of loss is probably small because of the long incubation time. however. the total amount of N lost from these systems was relatively small. Despite the inadequacies of the method for measurement for rate calculations. the results do show the influence of soil type and liming rate on the rate of loss. Fenn and Kissel (1976) show a multiple regression equation containing either 3 or 4 variables for NH₃-N loss. The variables included temperature. rate of NH₄⁺-N application, soil CaCO₃ content and time. They did note that the equations tended to overpredict N loss below 0.5% CaCO₃ and underpredicted it at high CaCO₃ amounts. They went on to discuss the limitations of applying their study to field conditions.

The cation associated with the ammonium fertilizer source has been recognized as an important factor affecting the volatilization of ammonia from soil. Urea is readily converted to NH₄⁺ which in turn is readily available for volatilization. Table 3 shows several studies where the

Table 1. Rate of NH₃ loss from soils treated with urea under a variety of laboratory conditions

Treatment	Application (kg N/ha)	Days of test	Rate of NH ₃ lost (kg N/ha/day)		
pH ¹ : 7.5	120	10	6.0		
6.0	"	"	2.0		
5.0	"	"	1.1		
Temperature (° C) ¹ :					
35	120	11	2.6		
15	"	"	1.1		
7	"	"	0.6		
Depth ¹ :					
Surface	120	13	1.5		
3.8 cm	"	"	0.5		
Initial moisture (%)					
A ¹ 37.5	120	14	1.7		
21	"	"	1.0		
1		"	fsl	0.3	
B ² 30	300	10	1.4	9.8	
20	"	"	8.8	10.8	
10	"	"	0.6	4.4	
5	"	"	0.03	0.3	
Application ²	300	28	3.4		
	150	"	1.2		
	50	"	0.2		
	0	"	0		

¹ Earnest and Massey. 1960

² Kresge and Satchell. 1960

Table 2. Rate of NH₃ loss from limed and unlimed Ontario soils incubated for 67 days ¹

Soil	CaCO ₃ treatment	Rate of NH ₃ loss kg N/ha/day
Vaudreuil (s)	0	0.03
	pH 7	0.25
	10%	0.18
St. Thomas (s)	0	0.03
	pH 7	0.08
	10%	0.42
Haldimand (c)	0	0.02
	pH 7	0.05
	10%	0.07

¹ Kowalenko, C.G., unpublished results

Table 3. Rate of NH₃ lost from surface application of various ammonium compounds under laboratory conditions.

Reference	Compound Application	(kg N/ha)	Duration (day)	Rate of NH ₃ loss (kg N/ha/day)	
Fenn and	NH ₄ F	550	4	89	
Kissel, 1973	(NH ₄) ₂ SO ₂	"	"	72	
	(NH ₄) ₂ HPO ₄	"	"	67	
	NH ₄ NO ₃	"	"	24	
	NH ₄ Cl	"	"	24	
	NH ₄ I	"	"	22	
Matocha, 1976	(NH ₂) ₂ CO	600	14	22	
	(NH ₄) ₂ SO ₄	"	"	10	
	SCU ¹ -30	"	"	4	
	SCU-20	"	"	1	
Kresge &	NH ₄ NO ₃ /urea			sl	<u>sil</u>
Satchell, 1960	0/100	354	11	25	26
	45/55	286	"	7	8
	70/30	241	"	4	3

¹ Sulfur coated urea

form of the fertilizer was examined in terms of loss of NH_3 . In each case, this has been directed toward surface application and very high rates of loss can occur. Again, one should realize that the values are average rates over the interval of the measurements, hence the instantaneous maximum rate of NH_3 loss can be exceedingly high.

Very few reports show data for field experiments of NH_3 loss. This is probably because current field methods applied still introduce an artificial situation (eg. absorption systems require a canopy) or measurement of NH_4^+ remaining in a system may include other types of loss. Table 4 shows results of three field experiments, which shows the possible losses. The results show the importance of large instantaneous losses from particular fertilizer practices.

This review of rates of the process of NH_3 volatilization shows that most of the studies have been concerned with a particular fertilizer practice (surface application). With this practice, very large quick losses of NH_3 can occur. On the other hand, small but continuous losses may be possible, but have not been considered to any great extent. It appears safe to assume that intimately mixed NH_4^+ at moderate application rates results in small, almost insignificant, losses which may or may not be continuous (eg. Hooker *et al.*, 1973 - Table 4; Stefanson, 1972). Another area of potentially large NH_3 losses is from manure application (see Lauer *et al.*, 1976) but the diversity of materials and methods of application makes it difficult to relate one study to the other.

(4) Biological denitrification

A large amount of work has been done on biological denitrification but most of it has been laboratory oriented. A number of soil and environmental factors have been shown to influence the rate of the process (Table 5). The results presented have been calculated on a per day basis to allow comparison from one study to the other. An exceedingly wide range of rates of denitrification is evident and this is probably due in large measure to the experimental set up rather than differences in soils used. In most cases shown in the table, conditions were such that denitrification was promoted (eg) addition of a high energy carbon source, anaerobic atmosphere, etc. This was usually done to facilitate the measurements. Each study had specific conditions, therefore comparison from one reference to the other must take these differences into account. Values within a particular study are comparable, therefore, the table should show the effect a particular factor has on the rate of the process. It is evident from the table that two different units are used for the rate and this is due to the authors' view of the process. A good example for the two approaches for rate calculation is shown by Stanford *et al.* (1975b) where zero and first order reaction approaches are calculated and discussed. This study even suggests a second order reaction in certain cases. A valuable discussion on denitrification kinetics by Kohl *et al.* (1976) shows the limitations of good fits of data to mathematical forms. They point out that the mathematical forms may be useful for certain uses (eg) modelling activities but do not prove anything about the reaction mechanism. The report of Sandhu and Moraghan (1972) shows the effect measurement time (even very short time) has on the rate of denitrification, where the rate can vary from 0-99 pg N/g soil/day. Numerous other studies have been done on

Table 4. Rate of NH₃ loss from field systems

Reference	Compound	Application (kg N/ha)	Duration of measurement	Rate of NH ₃ loss (kg N/ha/day)
Makarov, 1960	Urea	variable	variable short term	0 - 1.05
McGarity and Rajaratnam 1973	Urea	118	3 hr periods 24 hr. 14 days	7.68 (max) 2.4 (max) 0.03 (ave)
Hooker et al., 1973	Plowing of native sods	-	160 days	0.006

denitrification but most of these have insufficient information for rate calculation. A new laboratory approach which may have great potential in measuring denitrification rates bears consideration for the future (Bolderston *et al.*, 1976; Yoshinari and Knowles, 1976). The technique enables measurement of N_2O_3 as the gaseous product by blocking N_2O_3 reduction with acetylene. Considerable examination of this technique is required before application.

A laboratory technique which extends denitrification measurements to soil-plant systems has been used to determine the effect of a number of factors on the process (Table 6). This system would not distinguish between gaseous losses from the soil and that from the plant. The previous discussion on plant N suggested that most losses occur during maturity. The 35-40 day experimental did not extend into this apparently critical stage of plant growth for N losses, therefore most of the denitrification would be a result of the soil. The experiments in the majority of cases showed that plants stimulated denitrification. This cannot be compared with a fallow vs cropped condition in the field because in the field situation the fallow would have a different moisture status than a cropped system due to transpiration. The laboratory studies reported in the table would have the moisture contents adjusted to the same percentage.

A number of workers have used another laboratory approach to simulate some field conditions by using continuous flow soil columns (Table 7). Most of these workers have applied zero or first order reaction calculations to examine the rate of the process hence the units are not uniform. McLaren (1976) suggests that microbial numbers be taken into account when calculating rate constants. The diversity of factors which may influence the rate in this experimental system is very large and this makes it difficult to relate the rate constants to field situations. Volz *et al.* (1974) have made an attempt to apply this approach to a particular field situation.

Very few measurements of denitrification in the field have been done because of the analytical problems associated with it. The few results available (Table 8) suggest lower rates than what might be considered from laboratory incubations (Table 1). Similar comments can be made on field measurements as on laboratory measurements (ie) the length of the measurement period can be an important factor in calculating the rate. Two possibilities exist for denitrification namely (1) short bursts of denitrification involving high rates and (2) moderate to small losses over an extended period. In both cases, measurement of the rate of the process is very difficult and no method is entirely satisfactory. It should also be noted that measurement of N_2O_3 flux from a soil (see Table 8) only measures one of the gaseous products of the process, therefore values calculated are probably underestimated.

An alternative of measuring denitrification losses by analysing the gases evolved is through N balance methods. With the advent of isotope methods, N balances can be done with much greater accuracy than relying on non-isotope methods. A large number of ^{15}N balance studies are reported, most of which are laboratory or greenhouse studies. Table 9 displays a number of the balance studies done out in the field using ^{15}N . These results show a range of 0 - 1.43 kg N lost/ ha/day. It must be noted that some studies eg. Shields *et al.* (1973)

Table 5. Laboratory incubation studies showing the effect of a variety of factors on rate of denitrification

Factor	Condition		Rate of denitrification			Reference
			Average	Range	Unit	
Temperature (°C)	30		149	60-257	µgN/g/day	Cooper & Smith, 1963
	25		106	86-132	"	
	20		77	60-98	"	
	35		2.98	2.17-3.79	day ⁻¹	Stanford <i>et al</i> 1975c
	15		0.70	0.45-0.54	"	
	5		0.04	0.03-0.03	"	
	270		8.4	0-32.1	µgN/g/day	
Application rate (µg NO ₃ -N/g soil)	143		9.3	0-55.7	f	Wijler & Delwicke, 1954
	78		18.9	0-48.0	"	
	48		22.9	0-32.2	"	
	300		257	-	µgN/g/day	Cooper & Smith, 1963
	150		226	-	"	
	75		226	-	"	
	37.5		158	-	"	
pH	8.0		-	0-12	µgN/g/day	Focht, 1974
	6.0		-	0-6	"	
	4.35		-	0-3.5	"	
			<u>Aerobic</u>	<u>Anaerobic</u>		
Aerobic- anaerobic cycler (total of 128 days)	2		2	1.77	-	µgN/g/day
	4		4	1.66	-	"
	8		8	1.56	-	"
	16		16	1.45	-	"
	32		32	1.31	-	"
	64		64	1.00	-	"
	128		-	0.50	-	"
	-		1280	0.04	-	"
Soil type	7 soils		-	38-91		Khan & Moore, 1968 Burford & Bremner, 1975
	17 "		19.6	2-55	"	
	6 "		0.38	0.07-0.91	day ⁻¹	Stanford <i>et al.</i> , 1975a
	30 "		13.75	1.87-25.0	"	
	30 "		0.37	0.0264-0.9	µgN/g/day	Stanford <i>et al.</i> , 1975b Stanford <i>et al.</i> , 1975c
	0		0.13	0.04-0.24	day ⁻¹	
Glucose (mgC/g soil) (15°C)	0.3		0.70	0.54-0.86	"	
Cropping (15°C)	Fallow		50	-	µgN/g/day	Bailey, 1976
	Crop		82	-	"	
Sample depth	0-25 cm		0.15	0.09-0.23	"	Dubey and Fox, 1974
	25-50 cm		0.01	0-0.02	"	
	50-125 cm		0	0	"	
Hr. of incubation	0-4		44	30-60	"	Sandhu and Moraghan, 1972
	4-8		28	12-48	"	
	8-24		62	23-99	"	
	24-48		29	15-50	"	
	48-96		4	0-10	"	

Table 6. Sealed growth chamber studies of denitrification in a soil-plant (*Triticum aestivum*) system at 20C constant temperature.

Factor	Condition	Rate of denitrification (kgN/ha/day) ¹		Reference
		Average	Range	
Moisture content	8-36%	2.94	0.15-7.09	Stefanson & Greenland, 1970
	16-24%	1.69	0-6.05	
	28-36%	7.26	0-28.0	
Vegetation	Fallow	1.80	0.10-6.96	Stefanson, 1972a
	Cropped	2.11	0.15-7.15	
N source	NO ₃	2.08	0.76-5.02	Stefanson, 1972b
	NH ₄	0.94	0.54-3.67	
Sample	Undisturbed	0.40	0.25-1.61	Stefanson, 1972c
	Disturbed	0.60	0.47-1.97	
Soil	Pasture	1.97	0.77-4.76	Stefanson, 1973
	Cultivated	0.93	0.24-1.70	
Fertilizer depth	5 cm	1.35	0.45-2.24	Stefanson, 1976
	15 cm	1.35	0.87-2.02	
	25 cm	1.35	0.67-2.24	

¹ Experimental period between 35-40 days

Table 7. Denitrification from continuous flow soil¹ columns

Temperature (°C)	Other Conditions	Average	Range	Units	Reference
20	2 sequential runs	0.0024	0.003-0.0017	hr ⁻¹	Starr <i>et al</i> , 1974
19.5	ambient atmosphere	0.002	-	"	
"	0.5% O ₂	0.019	0.003-0.05	"	
"	5% O ₂	0.001		"	
"	20% O ₂	0.0035	0.001-0.008	"	Misra <i>et al</i> , 1974a, b
34.5	0.5% O ₂	0.035		"	
"	5% O ₂	0.04		"	
"	20% O ₂	0.045		"	
22	zero order calc.	0.056		µgN/g/hr	Doner <i>et al</i> , 1974
"	perfusion		1.2 - 2.3 x10 ⁻⁵	ppm/hr cm ⁻³	Ardakani <i>et al</i> , 1975
"	100 ppm NO ₃ after various conditioning	0.2		µgN/g/hr	Doner, 1975

¹ Soils were either Hanford sl or Columbia sil.

Table 8. Estimations of gaseous N loss from field systems

Method	Soil		Rate of denitrification (kg N/ha/day)	Reference
N ₂ O flux measurement	Urrbrae (f si)		0.01-0.09	Burford & Millington, 1968
Gas measurements	Chernozem & Day time		0.0004-0.0065	Borisova et al, ¹ 1972
	Podzol Entire day		0.0004-0.0215	
N balance	Chernozem		0.00376	
	Podzol		0.0029	
N ₂ O flux measurement	Chernozem		0.695	Burford and Stefanson, 1973
	Podzol		0.363	
	Permanent pasture		0.010	
NO ₃ changes in ponded soil	Hanford (56) Aug	2-7	1.45	Volz <i>et al</i> , 1975a
		7-9	2.47	
		9-13	0.70	
		13-15	2.31	
N ₂ O and ¹⁵ N ₂ flux measurement	Yolo (1)		0-12	Rolston <i>et al</i> , 1976

¹ Measurements during June through August

show almost complete recovery of ^{15}N applied and in many cases it is a result of the experimental set up which may or may not relate to practical methods. In the example given, glucose was added at such a rate that N was immobilized leaving little or no NO_3^- available for denitrification. This may be similar to adding N fertilizer to high straw situation which Myers and Paul (1971) did show reduced denitrification. Other cautions must be given when considering these results for denitrification. First, it has to be assumed that unrecovered N is denitrification. Other types of losses, such as volatilization, chemical decomposition may also be included which may result in overestimation of the rate of this particular process. On the other hand, ^{15}N only gives information on the labelled input (ie) loss due to the fertilizer N and will not include denitrification of unlabelled soil N. This would result in underestimation of the rate of denitrification from the soil. Isotope N balances have been done on forested situations and are not discussed here except to mention a couple of reviews (Gessel *et al.*, 1973; Knowles, 1975).

Long term balance sheets using non-isotope methods will include both soil and fertilizer N. Allison (1955; 1965) and Kundler (1970) reviewed the literature available on N balance sheets and concluded that 5-30% (average 15%) of the nitrogen in a soil system is lost, most likely due to denitrification. Table 10 summarized the information available on rates of denitrification from Allison's review. The author of the review discussed the limitations of such an approach. Other balance sheets are probably available since this review, but considering the limitations of the method little additional information can be added. Two specific cases of rates that can be calculated from non-isotope balance sheets suggest very high rates of denitrification (eg) minimum of 3.9 and 717 kg N lost /ha/day from Wallingford *et al* (1974) and Avorimelech and Ravek (194) reports, respectively. These are cases where large losses of N by denitrification would be expected.

In conclusion, one can see that although considerable work has been done on denitrification and other gaseous N losses most studies have considered the process from a qualitative rather than quantitative aspect. Rates shown in the tables were largely calculated from the data provided in the report. Many other reports did not have sufficient information to make rate calculations. The comments in the text showed limitations of available data (eg) timing of measurement especially, and if this area is to be understood more fully then experiments designed with this specifically in mind must be done.

MINERALIZATION

Mineralization of soil nitrogen is the conversion of organic nitrogen to inorganic nitrogen. The first inorganic N compound formed is $\text{NH}_4^+\text{-N}$, but in most soils nitrification (oxidation of $\text{NH}_4^+\text{-N}$ to $\text{NO}_3^-\text{-N}$) proceeds very rapidly such that $\text{NH}_4^+\text{-N}$ does not accumulate. The result then is decrease of the organic N with a corresponding increase in $\text{NO}_3^-\text{-N}$ rather than ammonium. The term ammonification is often used as the conversion of organic N to $\text{NH}_4^+\text{-N}$ and nitrification is used for the further oxidation of $\text{NH}_4^+\text{-N}$. Mineralization is used synonymous to ammonification in some of the literature but often is used as a term including ammonification and nitrification. In this discussion mineralization is used as the term for formation of inorganic N from the organic N which may or may not be further oxidized. It

should also be remembered the inorganic N can be converted to organic N (immobilization concurrent to mineralization) therefore measurements of inorganic N changes are usually values for the net effect of these processes.

Similar to denitrification, mineralization is difficult to quantify, not only because changes of inorganic N are a net result of mineralization and immobilization but other gains (N fixation, precipitation, etc.) and losses (denitrification, volatilization, leaching, etc.) must also be considered. In a plant system, it is almost impossible to separate the contribution of mineralized soil N from N extracted from deep in the profile to plant N. It is for this reason that considerably more information is available from laboratory than from field studies.

Since almost all of the N in soil is organically bound and since plants are not able to absorb significant amounts, mineralization is an important process in soil-plant systems. In order to assess the adequacy of the soil to supply N to a crop, the initial level of inorganic N and the rate of mineralization have to be considered. The most difficult part of this exercise is to determine the rate of mineralization. There has been an early and continuing interest in incubation systems which give information on mineralization rates (Harmsen and van Schreven, 1955; Bremner, 1965; Keeney and Bremner, 1966, a, b 1967). There have been many variations of the incubations with some attempts to simplify the procedure by forming a chemical test (Jenkinson, 1968). A major problem associated with this procedure is relating the mineralization rate to field conditions. More recently this concept has been pursued at greater depth with modifications to take into consideration temperature (Stanford *et al.*, 1973a, 1975), moisture content (Stanford and Epstein, 1974) and length of incubation (Stanford *et al.*, 1974 besides soil effects (Stanford and Smith, 1972). These values have also been related to plant growth (Stanford *et al.*, 1973b). Table 11 shows results of this type from several studies. Comparisons are difficult because of differences in incubation methods, duration of test, type of storage, etc. Use of this test has gained various responses in application to fertility assessment. In Ontario, for example, an incubation system pioneered by Eagle and Matthews (1958) was eventually abandoned because of lack of confidence. It is also difficult to relate these incubation rates to rates that occur in the field. Values obtained in this way are usually high with short term incubations in the laboratory often relating to mineralization throughout an entire growing season (Harnsen and van Schreven, 1955). Mineralization potentials give relative rather than actual N-supplying capacities of soils (Stanford, 1973) .

Simplicity is an important factor in choosing a method of evaluating mineralization potentials of soils, therefore, short incubations under constant moisture content and temperature are desirable. However, comparison with field observations have suggested that field situations are much more complex. Laboratory studies have shown that environmental factors such as freezing-thawing, wetting-drying, partial sterilization, fluctuating temperatures and moisture-temperature interactions (Mack, 1963; Agarwal *et al.*, 1971; Biederbeck and Campbell, 1973; Shields *et al.*, 1974; Kowalenko and Cameron, 1976) and soil factors such

Table 9. Denitrification rate calculated from ¹⁵N balance study under field conditions

Crop	Duration (days)	Rate of denitrification (kgN/ha/day)	Reference
Corn	730	0.07-0.16 ¹	Owens, 1960
Corn	56	0.48	Firth <i>et al</i> , 1973
	84	0.49	
Oats	Assume 100	0.16-0.20	Zamyatina <i>et al.</i> , 1968
"	" "	0.16-0.50	Zamyatina, 1971
"	" "	0.17	Koren'kov <i>et al</i> , 1975
Wheat	Assume 120	0.03-0.33	Myers & Paul, 1971
Wheat	0-28	0	
	28-70	0.16	
	70-105	0.47	
	0-105	0.22	
Fallow	0-28	0	Craswell & Martin, 1975
	28-70	0.40	
	70-105	0.21	
	0-105	0.22	
Barley	0-85	0.48	
Fallow	0-43	0.63	
	43-85	0.75	
	85-159	0.09	Kowalenko (unpublished data)
	159-316	0.15 ²	
	316-511	0.06 ²	
Sorghum	142	0.06-0.30 ³	Westerman <i>et al</i> , 1972
"	58	0.41	Kissel <i>et al</i> , 1976
	149	0.13	
Sudan grass	56	0.24-0.72	Carter <i>et al</i> , 1967
	70	1.10 ²	
Pasture			
- autumn fertilization	0-28	0-1.43 ⁴	
	0-84	0.06-0.47	Simpson, 1968
- winter fertilization	0-28	0.02-0.88	
	0-84	0.03-0.53	
Pasture			
- prairie-like soil	56	0.34	
- solodic "	301	0.002	Vallis <i>et al</i> , 1973
- yellow & gleyed podzolic soil	56	0.002	

¹ Rate increased with increasing level of irrigation

² Some evidence of leaching losses

³ Rate increased with increasing rate of fertilization

⁴ High rate of loss may include volatilization of NH₃

Table 10. Denitrification rate from long term balance sheets (Allison, 1955)

Place	Duration (yrs)	Number of treatments	N not accounted for (kgN/ha/yr)	
			Range	Average
I Lysimeter experiments				
Ithaca, NY	15	2	45-49	47
" "	8	4	4-50	24
Geneva, NY	16	2	12-52	32 ¹
Windsor, Conn.	10	11	[+12] ¹ 8-72	24
Riverside, Calif.	15	6	1-36	24
II Field experiments				
Rothamsted, Eng.	60	4	[+2] ¹ 4-160	55
New Brunswick, NJ	40	10	49-87	66
Overall range			1-160	
kgN/ha/day			0.003-0.44	

¹ Net gain of N

² Average included net gain

Table 11. Some mineralization potentials of soils using laboratory incubation methods.

Number of soils	Duration (wks)	Temperature (°C)	Rate Range (Ave)	Reference
5	1	35	49-92(70) µgN/g/wk	Eagle & Matthews, 1958
10	1	30	16-58 (32) " "	Keeney & Bremner, 1967
"	3	30	9-30 (18) " "	
53	3	25	3-37 (12) " "	Jenkinson, 1968
10	2	"	-2-64 (23) " "	Lathwell et al, 1972
"	16	"	2-16 (7) " "	
39	30	35	¹ 0.054-0.009wk ⁻¹	Stanford & Smith, 1972

¹ Rate constant

as pH, nitrogen and carbon fractions etc. (Hague & Walmsley, 1972; Dancer *et al*, 1973; Mundra *et al*, 1973; Cornfield, 1952; Cornforth, 1971) can influence mineralization rates. Carbon amendments will alter the balance of mineralization-immobilization (Black and Ritz, 1972; Ahmad *et al*, 1972, 1973), which is a very important consideration in systems cultivating straw or other organic residue into the soil. The nature of the organic material, particularly the C/N ratio is critical in the mineralization-immobilization balance. Ordinary chemical methods can only measure the net result of mineralization- immobilization processes, but tracer techniques have the advantage of distinguishing more clearly the relative importance of the two opposing transformations. A variety of laboratory studies using tracer techniques have greatly added to our understanding of these processes (Hiltbold *et al.*, 1950; Janson, 1958; Broadbent, 1966; Moore and Russell, 1970; Wojcik-Wojtkowiak, 1972; McGill *et al*, 1975; Chichester *et al*, 1975).

Experience with field and laboratory information has resulted in the generalization that 1 to 3% of the soil N is mineralized during the growing season (Bremner, 1965a,b; Woodruff, 1949; Slater and Green, 1933; Bartholomew and Kirkham, 1960). This type of information has limited usefulness when specific cases are being considered and particularly when rate information is desired for specific reasons or parts thereof. Table 12 shows estimated average daily rates of mineralization that can be calculated from field data provided. It should be remembered the kind of problems associated with this type of calculation (ie) losses and gains make a difference as well as the season and length of time between measurements. However, the data does show a range between 0 and 3.84 kgN released/ha/day. Tracer techniques, whether using ¹⁵N, Multiple labelling or radio-carbon dating, has and will add greatly to understanding the dynamics of organic matter degradation in soils (Clark and Paul, 1970; Shields and Paul, 1973).

Another approach to the consideration of N mineralization is the effect that cultivation has on organic N. In general, cultivation results in a decline of the organic nitrogen and in order for the N to be lost as NO₃, N must be mineralized. It appears that the cultivation increases net mineralization the soil from the virgin equilibrium level until a new lower equilibrium level has been reached (Bartholomew and Kirkham, 1960; Stevenson, 1965). There has been recurring interest in this aspect in western Canada (Alberta Soil Sci. Workshop, 1976; Martel and Paul, 1974; Ferguson and Gorby, 1971; Bishop and Atkinson, 1954; Campbell *et al*, 1975) and few studies in eastern Canada (Ho-Yen *et al.*, 1954).

More recently, mathematical simulation activities have had to consider this process and its rate in the soil and many approaches are possible (Morel, 1970; Dutt *et al*, 1972; Beek and Frissel, 1973; Hagin and Amberger, 1974; Greenwood *et al*, 1974; Campbell *et al*, 1974; Russel, 1975; Parnas, 1975; Slavnina and Pashveva, 1976). These as yet have theoretical beginning and have yet to be adequately tested particularly under field conditions.

Table 12. Average daily rate of mineralization estimated from field studies

Location	Soil	Duration of measure	Mineralization (kgN/ha/day)		Reference
			Average	Range	
New York	-	15 yr.	0.12	0.10-0.14	Allison, 1955
"	-	16 "	0.26	0.16-0.36	
Connecticut	-	10 "	0.03	0-0.11	
"	-	10 "	0.09	0-0.21	
California	-	15 "	0.26	0.15-0.39	
England	-	59 "	0.03	0-0.08	
New Jersey	Limed	40 "	0.12	0.02-0.19
	Unlimed	40 "	0.16	0.06-0.21	
Saskatchewan	Chern. Blk.	191 days	0.41	0.14-0.64	Shields et al, 1973
"	Br.	1 yr.	0.21	-	Rennie et al, 1974
S. Ontario	Maple woodlot	2 yr.	1.75	3.84-0.64	Ellis, 1974
	Pasture	"	1.51	-	
	Pine plantation	"	0.43	-	
	Old-field	"	0.56	0.96-0.64	
E. Ontario	Orthic Gleysol	85 days	0.93	0.77-1.10	Kowalenko, unpublished

PLANT UPTAKE

Another very important N process in the nitrogen cycle of soil systems is plant uptake. A great deal of work has been done on this subject both from a plant physiological (Baldwin, 1975; Baldwin *et al.*, 1972; Bassioni, 1972; Brewster and Tinker, 1972; Clarke and Barley, 1968; Hillel *et al.*, 1975; Nielsen, 1972; Pate, 1973; Philips *et al.*, 1976; NaNagara *et al.*, 1976) and soil-modelling (Caassen and Barber, 1976; Greenwood *et al.*, 1974; Seligman *et al.* 1975) orientation. This review is not intended to cover all aspects of nitrogen uptake rate but rather to present sufficient data to get a general picture of the situation. Emphasis is placed on field data and on selected crops (ie) corn, soybean, tobacco, wheat, rye, tomato and potato which were grown in the plots involved in this project. Several general comments can be made at the outset about a review of this type. First, few field studies have the rate of N uptake as the primary objective and the rates have been calculated from data presented. Many studies potentially have suitable data for this calculation except that one or two bits of information are not recorded (ie) precise length of time between samplings, N content of all plant parts, etc. The other prevalent feature of these reports are that most studies of plant N in the field do not include the contribution of the roots. This is probably because of the extreme difficulty of getting an accurate measure of the root mass and its N concentration. Often approximations are made of the contribution of root N to the total N of the plant (Viets, 1965; Stanford, 1973). This review then attempts to show N uptake rate using data of changes of plant N over specified time intervals. Many of the studies only include above ground plant parts and if a true rate is desired, an approximation of plant root N over that period can be added. Where possible, references on root N are suggested and possibly solution culture values may be useful.

(1) Corn

Table 13 shows representative uptake rates of N by corn and the data shows a relatively wide range of rates and that these rates can vary depending on the period of growth. The data in table 13 does not include the nitrogen in the root and if desired the root N may be estimated by considering root development patterns (Follett *et al.*, 1974; Mengel and Barber, 1974; Warnacke and Barber, 1974) and the % N in roots at these stages. A management practice that may influence the uptake on a hectare basis would be planting density and Table 14 shows results of uptake rate based on a single plant basis. This shows the rate can vary considerably and at certain stages of growth there may be a net loss of N (this may be due to loss of pollen, etc - Viets, 1965).

Not all of the plant N is removed from the soil system upon harvest and the proportion that remains depends on the management system. For example, corn may be only grown for the grain and the stover is worked back into the soil. In that case, about 50-61% of the above ground plant N will be removed with the grain (Chaudhary *et al.*, 1975) and the remainder becomes part of the soil system. A much greater amount of N is removed from the system if corn is used for silage.

Table 13. Average daily rate of N uptake by above ground parts of corn calculated on a per hectare basis

Time	Uptake rate-Av.(range) kg N/ha/day		Reference
1971	0.93	(0.72-1.13)	Chaudhary <i>et al.</i> , 1975
1972	1.04	(0.90-1.16)	
June 1 - July 29	1.49	(0.37-2.29)	Hanway, 1962
July 1 - Sept 22	0.80	(0.05-1.24)	
Unspecified	3.15		Firth <i>et al.</i> , 1973

Table 14. Rate of N uptake by corn calculated on a per plant basis

Days after seeding	Uptake rate (mgN/plant/day)	Reference
0-12	1	
12-19	6	
19-26	38	
26-33	81	
33-40	93	
40-47	96	
47-54	159	Bar-Yosef and Kafkafi, 1972 ¹
54-61	-21	
61-68	-46	
68-75	11	
75-83	-7	
<u>83-90</u>	-18	
Av.	30	
34-49	77 (52-96)	
49-76	47 (12-74)	NaNagara <i>et al.</i> , 1976
<u>76-97</u>	<u>37 (21-46)</u>	
34-97	51 (32-68)	
1970 0-14 (after silking)	31	
14-28 " "	17	Beauchamp <i>et al.</i> , 1976 ¹
1971 0-28 " "	26	

¹ Does not include accumulation of N in roots.

(2) Potatoes

Ezeta and McCollum (1972) provide a review table of N uptake rate by tops and tubers with a range of 1.7-4.2 kg N/ha/day. Reports by Soltanpour (1969) and Lorenz *et al.*(1974) suggest lower rates of uptake, with 0.20-1.25 kg N/ha/day calculated from their data. In each of these reports the roots were not included hence root development data and root N concentration are needed to estimate the contribution of this part to the total plant N (DeRoo and Waggoner, 1961; Ezeta and McCollum, 1972; Li and Soyre, 1975; Lesczynski and Tanner, 1976).

Unlike corn, a portion of the underground plant part (tuber) is removed entirely from the soil plant system, hence tuber N is lost from the field. Tuber N accounts for about 71% of the total N taken up (Soltanpour, 1969).

(3) Soybeans

It is difficult to know how to approach N uptake rate of soybeans since they are capable of fixing N by symbiotic process. This means that part of the N uptake may be from the air as well as from the soil, when the symbiotic process is proceeding. Table 15 presents uptake data and although nodulated and nonnodulated values are shown, it is difficult to know the exact distribution of N uptake between soil and air sources. Eck and Davis (1971), Hanway and Weber (1971a) and Mitchell and Russell (1971) may be useful references to estimate the contribution of roots to N uptake rate calculations.

Between 65-89% of the above ground plant-N is seed N at harvest (Hanway and Weber, 1971b; Ham *et al.*, 1975).

(4) Tobacco

Table 16 presents data on N uptake rate of tobacco considering the entire plant. In a field system, N uptake rate is important from transplant to harvest. Planting density may be important in considering uptake on a hectare basis so uptake rate on a single plant basis is included. The results show a wide range - rates dependent on the plants and environmental/fertilizer conditions as well as stage of growth.

A majority of the above ground plant N is removed at harvest but about 25% of the plant N will remain as roots (Takahashi, 1974).

(5) Tomato

There appears to be little work on N uptake by the entire tomato plant under field conditions probably because of the convenience of leaf N analysis to determine the nutrient status and greater interest in greenhouse growth conditions. Table 17 shows data on N uptake considering only above ground parts. More information on N uptake rate by tomatoes may be gathered from greenhouse and other non-field studies with suitable

precautions (Cannell *et al.*, 1963; Ajayi *et al.*, 1970; Shelton, 1976). At harvest about 61% of the above ground plant N is present in the fruit (Hester, 1938).

Table 15. Rate of N uptake by above ground parts of soybean

Rate of N uptake (kgN/ha/day)		Reference
2.02		Hanway and Weber, 1971b Hammond <i>et al.</i> , 1951
1.72 (4.9 max.)		
<u>Non nod.</u>	<u>Nod.</u>	
0.49-2.67	0.68-2.12	Ham <i>et al.</i> , 1975

Table 16. Rate of N uptake by tobacco on hectare and single plant basis

Time (days)	Rate of uptake gN/plant/day	Reference
July 10 - July 24	0.15 (0.12-0.19)	Zartman, Phillips &
July 24 - Aug. 7	0.16 (0.05-0.12)	Atkinson, 1976
Aug. 7- Aug. 21	-0.01 (-0.05-0.01)	
Aug. 21 - Aug. 28	0.04 and 0.11	
Aug.21 - Sept. 6	0.10 and 0.10	
65 after transplant	0.17	Takahashi, 19
93 " "	0.22	
	<u>kgN/ha/day</u>	
31-45 after transplant	3.2 (2.5-4.1)	
45-59 " "	3.4 (1.1-6.4)	Zartman, Phillips &
59-73 " "	0.06 (0-0.2)	Leggett, 1976
73-83 " "	1.8 (0.9-2.3)	

Table 17. Rate of N uptake by tomato plants (above ground parts)

Time (days)	Uptake rate (gN/plant/day)	Reference
30	0.02	
60	0.12	Hester, 1938
90	<u>0.36</u>	
Av	0.16	

(6) Wheat

Table 18 shows the range of daily uptake rate by wheat as effected by stage of growth and environmental/fertilizer conditions. Most of the data includes above ground plant parts only, but the report of McNeal et al. (1966) supplemented by knowledge of root development (Troughton, 1962; Welbank *et al.*, 1974) may make the results more meaningful in terms of the total N uptake.

Wheat grain may account for 72-82% of the above ground plant N at harvest.

(7) Rye

The rate of N uptake is rather difficult to assess if it is plant in the fall. This type of planting would result in a long dormant period over winter where the uptake calculation over the growing period would become meaningless. There appears to be little, if any, field oriented data where N uptake by fall rye can be calculated. This probably also reflects the relatively lower economic value of this cereal. Rumberg and Sneva (1970) have some field data of N uptake but only in the flowering stage where there was an apparent loss. Their nutrient solution culture experiment shows 0.013 and 0.004 g N uptake/plant/day for 11 and 23 successive periods. Grain N may account for 29% of the total plant N at harvest.

Table 18. Rate of N uptake by wheat using only above ground plant parts and total plant

Time after planting	Uptake rate (kgN/ha/day)	Reference
0-20	1.49	
20-33	3.90	
33-42	3.71	
42-54	2.92	McNeal <i>et al.</i> , 1966 ¹
54-64	0.88	
64-84	0.20	
<u>84-94</u>	<u>1.42</u>	
0-94	1.86	
210	0.77 Fuehring, 1969
1956-60	1.01	
1961-65	0.82	Widdowson and Penny, 1972
1966-1970	0.78
June 16 - July 2	1.59	
July 2 - July 15	1.90	
July 15 - July 28	1.93	Spratt, 1974
July 28 - Aug. 12	-1.18	
Aug. 12 - Aug. 28	1.04
0-42	0.58 (0.50-0.65)	
0-70	1.09 (0.91-1.43)	Craswell and Strong 1976
0-119	0.87 (0.72-0.97)	

References

GASEOUS LOSS OF N

(1) Chemical decomposition of oxidized inorganic N forms

- Allison, T.E. 1966. The fate of nitrogen applied to soils. *Adv. Agron.* 18, 219-258.
- Bollag, J.-M., S. Drzymala and L.T. Kardos. 1973. Biological versus chemical nitrite decomposition in soils. *Soil Sci.* 116, 44-50.
- Bremner, J.M. and D.W. Nelson. 1968. Chemical decomposition of nitrite in soils. 9th Intern. Congr. Soil Sci. 2, 495-503. Broadbent, F.E. and F. Clark. 1965. Denitrification. In *Soil nitrogen*. W.V. Bartholomew and F.E. Clark (eds). Agron. Series 10. Am. Soc. Agron., Madison pp. 344-359.
- Chao, T.T. 1967. Inorganic nitrogen transformation through the oxidation and reduction of sulfur compounds. *Soil Sci. Soc. Am. Proc.* 31, 485-490.
- Nelson, D.W. and J.M. Bremner. 1969. Factors affecting chemical transformations of nitrite in soils. *Soil Biol. Biochem.* 1, 229-239.
- Nelson, D.W. and J.M. Bremner. 1970. Role of soil minerals and metallic cations in nitrite decomposition and chemodenitrification in soils. *Soil Biol. Biochem.* 2, 1-8.
- Nommik, N. and J. Thorin. 1972. A mass spectrometric technique for studying the nitrogenous gases produced on the reaction of nitrite with raw humus. *Agrochimia* 16, 319-322.
- Pang, P.E., E.M. Otto and R.A. Hedlin. 1975a. Effects of pH and nitrifier population on nitrification of band-applied and homogeneously mixed urea nitrogen in soils. *Can. J. Soil Sci.* 55, 15-21.
- Pang, P.E., E.M. Cho and R.A. Hedlin. 1975b. Effects of nitrogen concentration on the transformation of band-applied nitrogen fertilizers. *Can. J. Soil Sci.* 55, 23-27.
- Russ, J.O. and R.L. Smith. 1965. Chemical reactions of nitrites in acid soil. *Soil Sci. Soc. Am. Proc.* 29, 267-270.
- Van Cleemput, O. and L. Baert. 1976. Theoretical consideration on nitrite self-decomposition reactions in soils. *Soil Sci. Soc. Am. J.* 40, 322-324.
- Woldendorp, J.W. 1968. Losses of soil nitrogen. *Stiktof* 1962 (No. 12), 32-46.

(2) Gaseous "leakage" during nitrification and plant growth

- Craswell, E.T. and A.E. Martin. 1975. Isotopic studies of the nitrogen balance in cracking clay. I. Recovery of added nitrogen from soil and wheat in the glasshouse and gas lysimeter. *Aust. J. Soil Res.* 13, 43-52.
- Daigger, L.A., D.H. Sander and G.A. Peterson. 1976. Nitrogen content of winter wheat during growth and maturation. *Agron. J.* 68, 815-818.
- Rumberg, C.B. and F.A. Sneva. 1970. Accumulation and loss of nitrogen during growth and maturation of cereal rye (*Secale cereale*). *Agron. J.* 62, 311-313.

- Viets, F.G.Jr. 1965. The plants need for and use of nitrogen. In Soil nitrogen. W.V. Bartholomew and F.E. Clark (eds). Agron. Series 10. Am. Soc. Agron., Madison pp. 503-549.
- Yoshida, T. and M. Alexander 1970. Nitrous oxide formation by *Nitrosomonas europaea* and heterotrophic microorganisms. Soil Sci. Soc. Am. Proc. 34, 880-882.

(3) Volatilization as NH₃

- Allison, F.E. 1955. The enigma of soil nitrogen balance sheets. Adv. Agron. 8, 213-250.
- Allison, F.E. 1965. The fate of nitrogen applied to soils. Adv. Agron. 18, 219-258.
- Chao, T.-T. and W. Kroontje. 1964. Relationship between ammonia volatilization, ammonia concentration and water evaporation. Soil Sci. Soc. Am. Proc. 28, 393-395.
- Earnest, J.W. and H.T. Massey. 1960. The effects of several factors on volatilization of ammonia formed from urea in the soil. Soil Sci. Soc. Am. Proc. 24, 87-90.
- Fenn, L.B. and D.E. Kissel. 1973. Ammonia volatilization from surface application of ammonium compounds on calcareous soils. I. General theory. Soil Sci. Soc. Am. Proc. 37, 855-859.
- Fenn, L.B. and D.E. Kissel. 1976. Ammonia volatilization from surface applications of ammonium compounds on calcareous soils: IV. Effect of calcium carbonate content. Soil Sci. Soc. Am. Proc. 39, 631-633.
- Filimonov, D.A. and R.A. Strel'nikova. 1974. Gaseous losses of ammonia following surface application of urea. Sov. Soil Sci. 6, 426-432.
- Harmsen, G.W. and G.J. Kolenbrander. 1965. Soil inorganic nitrogen. In Soil nitrogen. W.V. Bartholomew and F.E. Clark (eds) Agron. Series 10 Am. Soc. Agron., Madison pp. 43-92.
- Hooker, M.L., G.A. Peterson, and D.H. Sander. 1973. Ammonia nitrogen losses from simulated plowing of native sods. Soil Sci. Soc. Am. Proc. 31, 247-249.
- Kresge, C.B. and D.P. Satchell. 1960. Gaseous loss of ammonia from nitrogen fertilizers applied to soils. Agron. J. 52, 104-107.
- Lauer, D.A., D.R. Bouldin and S.D. Klausner. 1976. Ammonia volatilization from dairy manure spread on the soil surface. Environ. Qual. 5, 134-141.
- Mahendrappa, M.K. and E.D. Ogden. 1973. Patterns of ammonia volatilization from a forest soil. Plant Soil 38, 257-265.
- Mahendrappa, M.K. and P.O. Salonijs. 1974. Ammonia volatilization from black spruce raw humus treated with normal and controlled- release urea. Soil Sci. 117, 117-119.
- Makarov, B.N. 1960. Determination of ammonia from soil. Sov. Soil Sci. (1960), 882-884.
- Matocha, J.E. 1976. Ammonia volatilization and nitrogen utilization from sulfur-coated ureas and conventional nitrogen fertilizers. Soil Sci. Soc. Am. J. 40, 597-601.
- McGarity, J.W. and J.A. Rajaratnam. 1973. Apparatus for the measurement of nitrogen as gas from the field and simulated field environments. Soil Biol. Biochem. 5, 121-131.

- Mills, H.A., A.V. Barker and D.N. Mynard. 1974. Ammonia volatilization from soils. *Agron. J.* 66, 355-358.
- Nommik, H. 1973. Assessment of volatilization loss of ammonia from surface-applied urea on forest soil by N recovery. *Plant Soil* 38, 589-603.
- Overrien, L.N. 1968. Lysimeter studies on tracer nitrogen in forest soil: I. Nitrogen losses by leaching and volatilization after addition of urea-N¹⁵. *Soil Sci.* 106, 280-290.
- Overrien, L.N. 1969. Lysimeter studies on tracer nitrogen in forest soil: II Comparative loss of nitrogen through leaching and volatilization after the addition of urea-, ammonium-, and nitrate-N¹⁵. *Soil Sci.* 107, 149-159.
- Stefanson, R.E. 1972. Soil denitrification in sealed soil-plant systems I. Effect of plants, soil water content and soil organic matter content. *Plant Soil* 37, 113-127.
- Volk, G.M. 1970. Gaseous losses of ammonia from prilled urea applied to slash pine. *Soil Sci. Soc. Am. Proc.* 34, 513-519. Woldendorp, J.W. 1968. Losses of soil nitrogen. *Stikstof (1968)* No. 12, 32-46.

(4) Biological denitrification

- Allison, F.E. 1955. The enigma of soil nitrogen balance sheets. *Adv. Agron.* 8, 213-250.
- Allison, F.E. 1966. The fate of nitrogen applied to soils. *Adv. Agron.* 18, 219-258.
- Ardakani, M.S., L.W. Beker and A.D. McLaren. 1975. Reduction of nitrate in a soil column during continuous flow. *Soil Sci. Soc. Am. Proc.* 39, 270-294.
- Avnimelech, Y. and A. Raveh. 1974. The control of nitrate accumulation in soils by induced denitrification. *Water Res.* 8, 553-555.
- Bailey, L.D. 1976. Effects of temperature and root on denitrification in a soil. *Can. J. Soil Sci.* 56, 79-87.
- Balderston, W.L., B. Sherr and W.J. Payne. 1976. Blockage by acetylene of nitrous oxide reduction in *Pseudomonas perfectomarinus*. *Appl. Environ. Microbiol.* 31, 504-508.
- Borisova, N.N., S.N. Burtseva, V.N. Rodionov and O.K. Kirpaneva. 1972. Determination of nitrogen losses from soil in the form of different oxides and ammonia under field conditions. *Sov. Soil Sci.* 4, 540-546.
- Burford, J.R. and J.M. Bremner. 1975. Relationships between the denitrification capacities of soils and total, water-soluble and readily decomposable soil organic matter. *Soil Biol. Biochem.* 7, 389-394.
- Burford, J.R. and R.J. Millington. 1968. Nitrous oxide in the atmosphere of a red-brown earth. 9th Intern. Congr. Soil Sci. Adelaide 2, 505-511.
- Burford, J.R. and R.E. Stefanson. 1973. Measurement of gaseous losses of nitrogen from soils. *Soil Biol. Biochem.* 5, 133-141.
- Carter, J.N., O.L. Bennett and R.W. Pearson. 1967. Recover of fertilizer nitrogen under field conditions using nitrogen -15. *Soil Sci. Soc. Am. Proc.* 31, 50-56.
- Cooper, G.S. and R.L. Smith. 1963. Sequence of products formed during denitrification in some diverse western soils. *Soil Sci. Soc. Am. Proc.* 27, 659-662.

- Craswell, E.T. and A.E. Martin. 1975. Isotopic studies of the nitrogen balance in a cracking clay. II. Recovery of nitrate ^{15}N added to columns of packed soil and microplots growing in the field. *Aust. J. Soil Res.* 13, 53-61.
- Doner, J.E. 1975. Disappearance of nitrate under transient conditions in columns of soil. *Soil Biol. Biochem.* 7, 257-259.
- Doner, H.E., M.G. Volz and A.D. McLaren. 1974. Column studies of denitrification in soil. *Soil Biol. Biochem.* 6, 341-346.
- Dubey, H.D. and R.H. Fox. 1974. Denitrification losses from humid tropical soils of Puerto Rico. *Soil Sci. Soc. Am. Proc.* 38, 917-920.
- Firth, P., H. Thitipoca, S. Suthipradit, R. Westsalaar and D.T. Beech 1973. Nitrogen balance studies in the central plain of Thailand. *Soil Biol. Biochem.* 5, 41-46.
- Focht, D.D. 1974. The effect of temperature, pH, and aeration on the production of nitrous oxide and gaseous nitrogen-A zero-order kinetic model. *Soil Sci.* 118, 173-179.
- Gessel, S.P., D.W. Cole and E.C. Steinbrenner. 1973. Nitrogen balances in forest ecosystems of the Pacific Northwest. *Soil Biol. Biochem.* 5, 19-34.
- Khan, M.F.A. and A.W. Moore. 1968. Denitrifying capacity of some Alberta soils. *Can. J. Soil Sci.* 48, 89-91.
- Knowles, R. 1975. Interpretation of recent ^{15}N studies of nitrogen in forest systems. In *Forest soils and forest land management*. B. Bernier and C.H. Winget (eds). Proceedings of the 4th N. Am. Forest Soils Conference. Les Presses de l'Université Laval, Quebec.
- Kissel, D.E., S.J. Smith and D.W. Dillow. 1976. Disposition of fertilizer nitrate applied to a swelling clay in the field. *J. Environ. Qual.* 5, 66-71.
- Kohl, D.H., F. Vithayathl, P. Whitlow, G. Shearer and S.H. Chien. 1976. Denitrification kinetics in soil systems. The significance of good fits of data to mathematical forms. *Soil Sci. Soc. Am. Proc.* 40, 249-253.
- Koren'kov, D.A., L.I. Romanyuk, N.M. Varyushkina and L.I. Kirpaneva. 1975. Use of the stable N^{15} isotope to study the balance of fertilizer nitrogen in field lysimeters on sod-podzolic sandy loam soil. *Sov. Soil Sci.* 7, 244-249.
- Kundler, P. 1970. Ausnutzung, festlegung and verluste von dungemittelstickstoff. *Albrecht-Archiv.* 14, 191-210.
- McLaren, A.D. 1976. Rate constants for nitrification and denitrification in soils. *Rad. and Environm. Biophys.* 13, 43-48.
- Misra, C., D.R. Nielsen and J.W. Biggar. 1974a. Nitrogen transformations in soil during leaching. II. Steady state nitrification and nitrate reduction. *Soil Sci. Soc. Am. Proc.* 38, 294-299.
- Misra, C., D.R. Nielsen and J.W. Biggar. 1974b. Nitrogen transformation in soil during leaching: III. Nitrate reduction in soil columns. *Soil Sci. Soc. Am. Proc.* 38, 300-304.
- Myers, R.J.K. and E.A. Paul. 1971. Plant uptake and immobilization of ^{15}N -labelled ammonium nitrate in field experiment with wheat. *Intern. Atomic Energy Agency. Panel Proceedings Series.* pp. 55-64.
- Owens, L.D. 1960. Nitrogen movement and transformations in soils as evaluated by a lysimeter study utilizing isotope nitrogen. *Soil Sci. Soc. Am. Proc.* 24, 372-377.

- Reddy, K.R. and W.H. Patrick, Jr. 1975. Effect of alternate aerobic and anaerobic conditions on redox potential, organic matter decomposition and nitrogen loss in a flooded soil. *Soil Biol. Biochem.* 7, 87-94.
- Rolston, D.E., M. Fried and D.A. Goldhamer. 1976. Denitrification measured directly from nitrogen and nitrous oxide gas fluxes. *Soil Sci. Soc. Am. J.* 40, 259-266.
- Sandhu, M.S. and J.T. Moraghan. 1972. Influence of three chemicals on soil biological activity. *Comm. Soil Sci. Plant Anal.* 3, 439-447.
- Shields, J.A., E.A. Paul and W.E. Lowe. 1973. Turnover of microbial tissue in soil under field conditions. *Soil Biol. Biochem.* 5, 753-764.
- Simpson, J.R. 1968. Losses of urea nitrogen from the surface of pasture soils. *Trans. 9th Intern. Congr. Soil Sci., Adelaide* 2, 459-466.
- Stanford, G., S. Dzienia and R.A. Vander Pol. 1975c. Effect of temperature on denitrification rate in soils. *Soil Sci. Soc. Am. Proc.* 39, 867-870.
- Stanford, G., J.O. Legg, S. Dzienia and E.C. Simpson, Jr. 1975a. Denitrification and associated nitrogen transformations in soils. *Soil Sci.* 120, 147-152.
- Stanford, G., R.A. Vander Pol and S. Dzienia. 1975b. Denitrification rates in relation to total and extractable soil carbon. *Soil Sci. Soc. Am. Proc.* 39, 284-289.
- Starr, J.L., F.E. Broadbent and D.R. Nielsen. 1974. Nitrogen transformations during continuous leaching. *Soil Sci. Soc. Am. Proc.* 38, 283-289.
- Stefanson, R.C. 1972a. Soil denitrification in sealed soil-plant systems. I. Effect of plants, soil water content and soil organic matter content. *Plant Soil* 37, 113-127.
- Stefanson, R.C. 1972b. Soil denitrification in sealed soil-plant systems. II. Effect of water content and form of applied nitrogen. *Plant Soil* 37, 129-140.
- Stefanson, R.C. 1972c. Soil denitrification in sealed soil-plant systems. III. Effect of disturbed and undisturbed soil samples. *Plant Soil* 37, 141-149.
- Stefanson, R.C. 1973. Evolution patterns of nitrogen oxide in sealed soil-plant systems. *Soil Biol. Biochem.* 5, 167-169.
- Stefanson, R.C. 1976. Denitrification from nitrogen fertilizer placed at various depths in the soil-plant system. *Soil Sci.* 121, 353-363.
- Stefanson, R.C. and D.J. Greenland. 1970. Measurement of nitrogen and nitrous oxide evolution from soil-plant systems using sealed growth chambers. *Soil Sci.* 109, 203-206.
- Vallis, I., E.F. Henzell, A.E. Marten and P.J. Ross. 1973. Isotopic studies on the uptake of nitrogen by pasture plants V. ^{15}N balance experiments in field microplots. *Aust. J. Agric. Res.* 24, 693-702.
- Volz, M.G., L.W. Belser, M.S. Ardakani and A.D. McLaren. 1975a. Nitrate reduction and associated microbial populations in a ponded Hanford sandy loam. *J. Environ. Qual.* 4, 99-102.
- Wallingford, G.W., L.S. Murphy, W.L. Powers and H.L. Manges. 1975 Denitrification in soil treated with beef-feedlot manure. *Comm. Soil Sci. Plant Anal.* 6, 147-161.

- Westerman, R.L., L.T. Kurtz and R.D. Hauck. 1972. Recovery of ¹⁵N labelled fertilizers in field experiments. *Soil Sci. Soc. Am. Proc.* 36, 82-86.
- Wijler, J. and C.C. Delwiche. 1954. Investigations on the denitrifying process in soil. *Plant Soil* 5, 155-169.
- Yoshinari, T. and R. Knowles. 1976. Acetylene inhibition of nitrous oxide reduction by denitrifying bacteria. *Biochem. Biophys. Res. Comm.* 69, 705-710.
- Zamyatina, V.B. 1971. Nitrogen balance studies using ¹⁵N-labelled fertilizers. In *Nitrogen-15 in soil-plant studies*. Inter. Atomic Energy Agency. Panel Proceedings Series. pp. 33-45.
- Zamyatina, V.B., N.I. Borisova, N.M. Varyushkina, S.V. Burtzeva and J.I. Kireaneva. 1968. Investigations on the balance and use of ¹⁵N-tagged fertilizer nitrogen by plants in soils. 9th Intern. Congr. Soil Sci., Adelaide, 513-521.

MINERALIZATION

- Agarwal, A.S., B.R. Singh and Y. Kanehiro. 1971. Soil nitrogen and carbon mineralization as affected by drying-rewetting cycles. *Soil Sci. Soc. Am. Proc.* 35, 96-100.
- Ahmad, Z., H. Kai and T. Harada. 1972. Effect of nitrogenous forms on immobilization and release of nitrogen in soil. *J. Fac. Agr., Kyushi Univ.* 17, 49-65.
- Ahmad, Z., Y. Yahiro and H. Kai. 1973. Factors affecting immobilization and release of nitrogen in soil and chemical characteristics of the nitrogen newly immobilized. IV. Chemical nature of the organic nitrogen becoming decomposable due to the drying of soil. *Soil Sci. Plant Nutr.* 19, 287-298.
- Alberta Soil Science Workshop. 1976. Proceedings of western Canada nitrogen symposium. Soil and Feed Testing Lab., Edmonton.
- Allison, F.E. 1955. The enigma of soil nitrogen balance sheets. *Agron.* 8, 213-250.
- Bartholomew, W.V. and D. Kirkham. 1960. Mathematical descriptions and interpretations of culture induced soil nitrogen changes. 7th Intern. Cong. Soil Sci. Madison, Wisc. Comm. III 2, 471-477.
- Beek, J. and M.J. Frissel. 1973. Simulation of nitrogen behavior in soils. Centre for Agricultural Publishing and Documentation, Wageningen. 55 pp.
- Biederbeck, V.O. and C.A. Campbell. 1973. Soil microbial activity as influenced by temperature trends and fluctuations. *Can. J. Soil Sci.* 53, 363-376.
- Bishop, R.F. and H.J. Atkinson. 1954. Changes in organic matter and certain other soil constituents in forty years of cropping under irrigation. *Can. J. Agr. Sci.* 34, 177-180.
- Black, A.L. and L.L. Reitz. 1972. Phosphorus and nitrate-nitrogen immobilization by wheat straw. *Agron. J.* 64, 782-785.
- Bremner, J.M. 1965a. Organic nitrogen in soils. In *Soil nitrogen*. W.V. Bartholomew and F.E. Clark (eds). *Agron. Series 10* Am. Soc. Agron., Madison. pp. 93-149.
- Bremner, J.M. 1965b. Nitrogen availability indexes. In *Methods of soil analysis Part 2. Chemical and microbiological properties*. C.A. Black (ed) *Agron. Series 9*. Am. Soc. Agron. Madison. pp. 1324-1345.
- Broadbent, F.E. 1966. Interchange between inorganic and organic nitrogen in soils. *Hilgardia* 37, 165-180.

- Campbell, C.A., V.O. Biederbeck and W.G. Hinman. 1975. Relationship between nitrate in summer-fallowed surface soil and some environmental variables. *Can. J. Soil Sci.* 55, 213-223.
- Campbell, C.A., W. Nicholaichuk and F.G. Warder. 1975. Effects of a wheat-summer-fallow rotation on subsoil nitrate. *Can. J. Soil Sci.* 55, 279-286.
- Chichester, F.W., J.D. Legg and G. Stanford. 1975. Relative mineralization rates of indigenous and recently incorporated ¹⁵N-labelled nitrogen. *Soil Sci.* 120, 455-460.
- Clark, F.E. and E.A. Paul. 1970. The microflora of grassland. *Adv. Agron.* 22, 375-435.
- Cornfield, A.A. 1952. The mineralization of the nitrogen in soils during incubation: Influence of pH, total nitrogen, and organic carbon contents. *J. Sci. Food Agric.* 3, 343-349.
- Coenforth, I.S. 1971. Nitrogen mineralization in West Indian soils. *Exptl. Agric.* 7, 345-349.
- Dancer, W.S., L.A. Peterson and G. Chesters. 1973. Ammonification and nitrification of N as influenced by soil pH and previous N treatments. *Soil Sci. Soc. Am. Proc.* 37, 67-69.
- Dutt, G.R., M.J. Shoffer and W.J. Moore. 1972. Computer simulation model of dynamic bio-physicochemical processes in soils. Dept. of Soils, Water and Engineering. University of Arizona, Tuscon. Tech. Bull. 196.
- Eagle, D.J. and B.C. Matthews. 1958. Measurement of nitrate-supplying power of soils by an incubation method and correlation with crop yield response. *Can. J. Soil Sci.* 38, 161-170.
- Ellis, R.C. 1974. The seasonal pattern of nitrogen and carbon mineralization in forest and pasture soils in southern Ontario. *Can. J. Soil Sci.* 54, 15-28.
- Ferguson, W.S. and B.J. Gorby. 1971. Effect of various periods of seed-down to alfalfa and brome grass on soil nitrogen. *Can. J. Soil Sci.* 51, 65-73.
- Greenwood, D.J., J.T. Wood and T.J. Cleaver. 1974. A dynamic model for the effects of soil and weather conditions on nitrogen response. *J. Agric. Sci. Camb.* 82, 455-467.
- Hagin, J. and A. Amberger. 1974. Contribution of fertilizers and manures to the N- and P-load of waters. A computer simulation. Technion-Israel Institute of Technology. Soils and Fertilizers Laboratory. Haifa, Israel pp. 119
- Hague, I. and D. Walmsley. 1972. Incubation studies on mineralization of organic sulphur and organic nitrogen. *Plant Soil* 37, 255-264. Harmsen, G.W. and D.A. van Schreven. 1955. Mineralization of organic nitrogen in soil. *Adv. Agron.* 7, 299-398.
- Hiltbold, A.E., W.V. Bartholomew and C.H. Werkman. 1950. The use of tracer techniques in the simultaneous measurement of mineralization and immobilization of nitrogen in soil. *Soil Sci. Soc. Am. Proc.* 15, 166-173.
- Ho-Yen, B.O., L.B. MacLeod, W.A. DeLong and H.A. Stepler. 1954. Apparent effects of management on the organic matter, nitrogen and phosphorus contents of a Brown Forest soil. *Can. J. Agr. Sci.* 34, 507-513.
- Jansson, S.L. 1958. Tracer studies on nitrogen transformations in soil with special attention to mineralization-immobilization relationships. *Kungl. Lantbrukshogskolans Annaler* 24, 100-361.
- Jenkinson, D.S. 1968. Chemical tests for potentially available nitrogen in soil. *J. Sci. Fd. Agric.* 19, 160-168.

- Keeney, D.R. and J.M. Bremner. 1966a. Comparison and evaluation of laboratory methods of obtaining an index of soil nitrogen availability. *Agron. J.* 58, 498-503.
- Keeney, D.R. and J.M. Bremner. 1966b. Characterization of mineralizable nitrogen in soils. *Soil Sci. Soc. Am. Proc.* 30, 714-719.
- Keeney, D.R. and J.M. Bremner. 1967. Determination and isotope-ratio analysis of different forms of nitrogen in soils: 6. Mineralizable nitrogen. *Soil Sci. Soc. Am. Proc.* 31, 34-39.
- Kowalenko, C.G. and D.R. Cameron. 1976. Nitrogen transformations in an incubated soil as affected by combinations of moisture content and temperature and absorption-fixation of ammonium. *Can. J. Soil Sci.* 56, 63-70.
- Lathwell, D.J., H.D. Dubey and R.H. Fox. 1972. Nitrogen supplying power of some tropical soils of Puerto Rico and methods for its evaluation. *Agron. J.* 64, 763-766.
- Mack, A.R. 1963. Biological activity and mineralization of nitrogen in three soils as induced by freezing and drying. *Can. J. Soil Sci.* 43, 316-324.
- Martel, Y.A. and E.A. Paul. 1974. Effects of cultivation on the organic matter of grassland soils as determined by fractionation and radiocarbon dating. *Can. J. Soil Sci.* 54, 419-426.
- McGill, W.B., J.A. Shields and E.A. Paul. 1975. Relation between carbon and nitrogen turnover in soil organic fractions of microbial origin. *Soil Biol. Biochem.* 7, 57-63.
- Moore, A.W. and J.S. Russell. 1970. Changes in chemical fractions of nitrogen during incubation of soils with histories of large organic matter increase under pasture. *Aust. J. Soil Res.* 8, 21-30.
- Morel, R. 1970. Propositions nouvelles relatives au probleme de l'azote organique dans un sol en equilibre azote. *Ann. Agron.* 21, 197-210.
- Mundra, M.C., G.S. Bhandari and O.P. Srivastava. 1973. Studies on the mineralization and immobilization of nitrogen in soil. *Geoderma* 9, 27-33.
- Parnas, H. 1975. Model for decomposition of organic material by microorganisms. *Soil Biol. Biochem.* 7, 161-169.
- Rennie, D.A., E.A. Paul and L.E. Johns. 1974. Isotope tracer-aided research on the nitrogen cycle in selected Saskatchewan soils. In *Effects of agricultural production on nitrates in food and water with particular reference to isotope studies*. Intern. Atomic Energy Agency, Vienna pp. 77-90.
- Russell, J.S. 1975. A mathematical treatment of the effect of cropping system on soil organic nitrogen in two long-term sequential experiments. *Soil Sci.* 120, 37-44.
- Salter, R.M. and T.C. Green. 1933. Factors affecting the accumulation and loss of nitrogen and organic carbon in cropped soils. *Agron. J.* 25, 622-630.
- Shields, J.A. and E.A. Paul. 1973. Decomposition of ¹⁴C-labelled plant material under field conditions. *Can. J. Soil Sci.* 53, 297-306.
- Shields, J.A., E.A. Paul, and W.E. Lowe. 1973. Turnover of microbial tissue in soil under field conditions. *Soil Biol. Biochem.* 5, 753-764.
- Shields, J.A., E.A. Paul, and W.E. Lowe. 1974. Factors influencing the stability of labelled microbial materials in soils. *Soil Biol. Biochem.* 6, 31-37.
- Slavnina, T.P. and G.Y. Pashneva. 1976. Simulation of nitrification and ammonification in a gray forest soil. *Sov. Soil Sci.* 8, 186-192.
- Stanford, G. 1973. Rationale for optimum nitrogen fertilization in corn production. *J. Environ. Qual.* 2, 159-166.

- Stanford, G., J.N. Carter and S.J. Smith. 1974. Estimates of potentially mineralizable soil nitrogen based on short-term incubations. *Soil Sci. Soc. Am. Proc.* 38, 99-102.
- Stanford, G. and E. Epstein. 1974. Nitrogen mineralization-water relations in soils. *Soil Sci. Soc. Am. Proc.* 38, 103-107.
- Stanford, G., M.H. Frere and D.H. Schwaninger. 1973. Temperature coefficient of soil nitrogen mineralization. *Soil Sci.* 115, 321-323.
- Stanford, G., M.H. Frere and R.A. Vander Pol. 1975. Effect of fluctuating temperatures on soil nitrogen mineralization. *Soil Sci.* 119, 222-226.
- Stanford, G., J.D. Legg and S.J. Smith. 1973. Soil nitrogen availability evaluations based on nitrogen mineralization potentials and uptake of labeled and unlabeled nitrogen by plants. *Plant Soil* 39, 113-124. Stanford, G. and S.J. Smith. 1972. Nitrogen mineralization potentials of soils. *Soil Sci. Soc. Am. Proc.* 36, 465-472.
- Stevenson, F.J. 1965. Origin and distribution of nitrogen in soils. In *Soil nitrogen*. W.V. Bartholomew and F.E. Clark (eds) *Agron. Series 10*. Am. Soc. Agron., Madison. pp. 1-42.
- Wojcik-Wojtkowiak, D. 1972. The transformation of nitrogen and carbon in the soil during humification of straw labelled with N¹⁵. *Plant Soil* 36, 261-270.
- Woodruff, C.M. 1949. Estimating the nitrogen delivery of soil from the organic matter determination as reflected by Sanborn Field. *Soil Sci. Soc. Am. Proc.* 14, 208-212.

PLANT UPTAKE

- Baldwin, J.P. 1975. A quantitative analysis of the factors affecting plant nutrient uptake from some soils. *J. Soil Sci.* 26, 195-206.
- Baldwin, J.P., P.B.H. Tinker and P.H. Nye. 1972. Uptake of solutes by multiple root systems from soil. II. The theoretical effects of rooting density and pattern on uptake of nutrients from soil. *Plant Soil* 36, 693-708.
- Bassioni, N.H. 1972. On the mechanism of nitrate uptake by plant roots. Effect of other anions on nitrate uptake. *Agrochimica* 16, 519-522.
- Brewster, J. L. and P.B.H. Tinker. 1972. Nutrient flow rates into roots. *Soils Fert.* 35, 355-359.
- Caassen, N. and S.A. Barber. 1976. Simulation model for nutrient uptake from soil by a growing plant root system. *Agron. J.* 68, 961-964.
- Clarke, A.L. and K.P. Barley. 1968. The uptake of nitrogen from soils in relation to solute diffusion. *Australian J. Soil Res.* 6, 75-92. Greenwood, D.J., J.T. Wood and T.J. Cleaver. 1974. A dynamic model for the effects of soil and weather conditions on nitrogen response. *Agric. Sci. Camb.* 82, 455-467.
- Hillel, D., C.G.E.M. Van Beek and H. Talpaz. 1975. Microscopic scale model of soil-water and salt movement to plant roots. *Soil Sci.* 120, 385-399.
- NaNagara, T., R.G. Philips and J.E. Leggett. 1976. Diffusion and mass flow of nitrate-nitrogen into corn roots grown under field conditions. *Agron. J.* 68, 67-72.
- Nielsen, N.E. 1972. A transport kinetic concept of ion uptake from soil by plants. II. The concept and some theoretical considerations. *Plant Soil* 37, 561-576.

- Pate, J.S. 1973. Uptake, assimilation and transport of nitrogen compounds by plants. *Soil Biol. Biochem.* 5, 109-119.
- Philips, R.E., T. NaNagar, R.E. Zartman and J.E. Leggett. 1976. Diffusion and mass flow of nitrate-nitrogen to plant roots. *Agron. J.* 68, 63-66.
- Seligman, H.G., H. van Keulen and J. Goudriaan. 1975. An elementary model of nitrogen uptake and redistribution by annual plant species. *Decologia (Berl)* 21, 243-261.
- Stanford, G. 1973. Rationale for optimum nitrogen fertilization in corn production. *J. Environ. Qual.* 2, 159-166.
- Viets, F.G., Jr. 1965. The plant's need for and use of nitrogen. In *Soil nitrogen*. W.V. Bartholomew and F.E. Clark (eds.) *Agron. Series 10*. Am. Soc. Agron., Madison pp. 503-549.

CORN

- Bar-Yosef, B. and U. Kafkafi 1972. Rates of growth and nutrient uptake of irrigated corn as affected by N and P fertilization *Soil Sci. Soc. Am. Proc.* 36, 931-936.
- Beauchamp, E.G., L.W. Kannenberg and R.B. Hunter 1976. Nitrogen accumulation and translocation in corn genotype following silking. *Agron. J.* 68, 418-422.
- Chaudhry, T.N., V.K. Bhatnagar and S.S. Prihar 1975. Corn yield and nutrient uptake as affected by water-table depth and soil submergence. *Agron.* 745-749.
- Firth, P., H. Thitipoca, S. Suthipradit, R. Witselaor, and D.F. Bedch. 1973. Nitrogen balance studies in the Central Plain of Thailand. *Soil Biol. Biochem.* 5, 41-46.
- Follett, R.F., R.R. Allmaros and G.A. Reichman 1974. Distribution of corn roots in sandy soil with a declining water table. *Agron. J.* 66, 288-292.
- Hanway, J.J. 1962. Corn growth and composition in relation to soil fertility. II. Uptake of N, P, and K and their distribution in different plant parts during the growing season. *Agron. J.* 54, 217-222.
- Mengel, D.B. and S.A. Barber 1974. Development and distribution of the corn root system under field conditions. *Agron. J.* 66, 341-344. NaNagara, T., R.E. Phillips and J.E. Leggett 1976. Diffusion and mass flow of nitrate-nitrogen into corn roots grown under filed conditions *Agron.* 67-72.
- Warncke, D.D. and S.A. Barber 1974. Root development and nutrient uptake by corn growth in solution culture. *Agron. J.* 66, 514-516.

POTATOES

- De Roo, H. C. and P.E. Waggoner 1961. Root development of potatoes. *Agron. J.* 53, 15-17.
- Ezeta, F. N. and R.E. McCollum 1972. Dry-matter production, and nutrient uptake and removal by *Solanum andigean* in the Peruvian Andes. *Am. Potato J.* 49, 151-163.
- Lesczynski, D.B. and C.B. Tanner 1976. Seasonal variation of root distribution of irrigated, field-grown Russet Burbank potato. *Am. Potato J.* 53, 69-78.
- Li, P.H. and K.D. Sayre 1975. The protein, non-protein and total nitrogen in *Solanum tuberosum* Appr. *andigena* potatoes. *Am. Potato J.* 52, 341-350.
- Lorenz, O.A., B.L. Weir and J.C. Bishop 1974. Effect of sources of nitrogen on yield and nitrogen absorption of potatoes. *Am. Potato J.* 51, 56-65.

Soltanpour, P.N. 1969. Accumulation of dry matter and N, P, K by Russet Burbank, Dromonte and Red McClure potatoes. *Am. Potato J.* 46,111-119.

SOYBEANS

- Eck, H.V. and R.G. Davis 1971. Profile modification and root yield, distribution, and activity. *Agron. J.* 63, 934-937.
- Ham, G.E., J. E. Jiener, S.D. Evans, R.D. Frazier and W.W. Nelson 1975. Yield and composition of soybean seed as affected by N and S fertilization. *Agron. J.* 67, 293-297.
- Hammond, L.C., C.A. Blanck and A.G. Norman 1951. Nutrient uptake by soybeans on two Town soils. *Iowa Agr. Exp. Sta. Res. Bull.* 384.
- Hanway, J.J. and C.R. Webber 1971a N,P. and K percentages in soybean (*Glycine max* (L). Merrill) plant parts. *Agron. J.* 63, 286-290.
- Hanway, J.J. and C.R. Weber 1971b. Accumulation of N, P, and K by soybean (*Glycine max* (L). Merrill) plants. *Agron. J.* 63, 406-408.
- Mitchell, R.L. and W.J. Russell 1971. Root development and rooting patterns of soybean (*Glycine max* (L). Merrill) evaluation under field conditions. *Agron. J.* 63, 313-316.

TOBACCO

- Zartman, R.E., R.E. Phillips and J.E. Leggett 1976. Comparison of simulated and measured nitrogen accumulation in Burley tobacco. *Agron. J.* 68, 406-410.
- Zartman, R.E., R.E. Phillips and W.O. Atkinson 1976. Tillage and nitrogen influence on root densities of Burley tobacco. *Tobacco Sci.* 20, 136-139. Takahashi, 1974 Studies on the control of nicotine content in tobacco. *Tobacco Sci.*

TOMATO

- Ajayé, O. D.A. Maynard and A.V. Barker 1970. The effects of potassium on ammonium nutrition of tomato (*Lycopersicon esculentum* Mill). *Agron. J.* 62, 818-821.
- Cannell, G.H., F.T. Bingham, J.C. Lingle and M.J. Garber 1963. Yield and nutrient composition of tomatoes in relation to soil temperature, moisture and phosphorus levels. *Soil Sci. Soc. Am. Proc.* 27, 560-565.
- Hester, J.B. 1938. The absorption of nutrients by the tomato plant at different stages of growth. *Am. Soc. Hort. Sci. Proc.* 36, 720-722.
- Shelton, J.E. 1976. An evaluation of sulfur-coated area as a preplant total season supply for trellised tomatoes. *Soil Sci. Soc. Am. Proc.* 40, 126-129.

WHEAT

- Craswell, E.T. and W.M. Strong 1976. Isotopic studies of the nitrogen balance in a cracking clay. III Nitrogen recovery in plant and soil in relation to the depth of fertilizer addition and rainfall. *Aust. J. Soil Res.* 14,75-83.
- Tuehring, H.D. 1969. Irrigated wheat on a calcareous soil as affected by application of nitrogen, phosphorus, potassium, and zinc: I Yield, composition, and number of heads. *Agron. J.* 61, 591-596.

- McNeal, F.H., M.A. Berg and C.A. Watson 1966. Nitrogen and dry matter in five spring wheat varieties at successive, stages of development. *Agron. J.* 58,605-608.
- Spratt, E.D. 1974. Effect of ammonium and nitrate forms of fertilizer- -N and their time of application on utilization of N by wheat. *Agron. J.* 66, 57-61.
- Troughton, A. 1962. The roots of temperate cereals (wheat, barley, oats and rye). Mimeographed Publication No. 2/1962. Commonwealth Bureau of Pastures and Field Crops. Hurby, Berkshire
- Welbank, P.J., M.J. Gibb, P.J. Taylor and E.D. Williams 1974. Root growth of cereal crops. Rothamsted Experimental Station Report for 1973. Harpendon 66 pp.
- Widdowson, F.V. and A. Penny 1972. Yields and N, P and K contents of the crops grown in the Rothamsted Reference Experiment, 1956-70. Rothamsted Report for 1972. Part 2 pp. 111-130.

RYE

- Rumburg, C. B. and F.A. Sneva 1970. Accumulation and loss of nitrogen during growth and maturation of cereal rye (*Secak cereale*). *Agron. J.* 62, 311-313.